

11-18-2018

## An Intensive, Large-Scale Batch Culture System to Produce the Calanoid Copepod, *Acartia tonsa*: Appendix A

Brie L. Sarkisian

*University of Southern Mississippi*, Brie.Sarkisian@usm.edu

Jason T. Lemus

*Fish and Wildlife Research Institute, Florida*, jason.lemus@myfwc.com

Angelos Apeitos

*University of Southern Mississippi*, angelos.apeitos@usm.edu

Reginald B. Blaylock

*University of Southern Mississippi*, reg.blaylock@usm.edu

Eric Saillant

*University of Southern Mississippi*, Eric.Saillant@usm.edu

Follow this and additional works at: <https://aquila.usm.edu/datasets>



Part of the [Marine Biology Commons](#)

---

### Recommended Citation

Sarkisian, Brie L.; Lemus, Jason T.; Apeitos, Angelos; Blaylock, Reginald B.; and Saillant, Eric, "An Intensive, Large-Scale Batch Culture System to Produce the Calanoid Copepod, *Acartia tonsa*: Appendix A" (2018). *Research Data*. 1.

<https://aquila.usm.edu/datasets/1>

This Article is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in Research Data by an authorized administrator of The Aquila Digital Community. For more information, please contact [Joshua.Cromwell@usm.edu](mailto:Joshua.Cromwell@usm.edu).

1 Supplementary appendices to accompany:

2 Sarkisian, B. L., Lemus, J.T., Apeitos, A., Blaylock, R.B., and Saillant, E. A. 2018. An intensive,  
3 large-scale batch culture system to produce the calanoid copepod, *Acartia tonsa*. Aquaculture  
4 DOI: 10.1016/j.aquaculture.2018.11.042


5 Appendix A. Microalgae culture

6 Algae production is carried out in a batch culture system in a room adjacent to the  
7 copepod production system. A batch algae system is the most cost-effective method to provide  
8 for high copepod demands and a small volume footprint (150 L, 6 trillion total cells day<sup>-1</sup>).  
9 Tahitian–strain *Isochrysis lutea*, CCMP 1324 is used for the main source of algae to feed  
10 copepod cultures. Algae production is carried out in 19 L plastic bags set on racks with  
11 fluorescent lights between the racks. Light intensity is held at 12,000 – 17,000 lux by use of 32  
12 watt, 2,900 lumen, 4,100 Kelvin mercury-fluorescent bulbs (GE 66349 - F32T8/SPP41/ECO,  
13 General Electric, Boston, MA) over a 24-h photoperiod. The bags are inoculated with 2 L of  
14 algae (in exponential phase at a minimum density of 30 x 10<sup>6</sup> cells mL<sup>-1</sup>) and 16 L of seawater.  
15 Bags are fed F/2 Algae Food (Fritz Aquatics, Mesquite, TX), 2.5 mL of part A to 2.5 mL part B  
16 solution every other day. After a ten to twelve day grow-out period, the bags are harvested as  
17 feed for the copepods at a density of 40 - 50 x 10<sup>6</sup> cells mL<sup>-1</sup>.

18 For the years 2014 and 2015, *T-Iso* feedings were supplemented with *Rhodomonas lens*,  
19 Pascher & Ruttner, 1913, CCMP 739 (*Rhodo*) was produced in a separate room. This algae  
20 requires a specific photoperiod (10 h light : 14 h dark cycle using the same techniques as for T-  
21 Iso), so it was cultured in another building. *Rhodo* bags were inoculated and supplied with  
22 nutrient media the same way as *T-Iso*, reaching a density of 15 - 20 x 10<sup>6</sup> cells mL<sup>-1</sup>. On any  
23 given day, only a total of 25 L of *Rhodo* (0.4 trillion total cells) was available to supplement  
24 copepod feedings.

25

26 Appendix B. Egg production from a single tank over three trials in 2013, 2014, and 2015. Bars  
27 represent an average of daily egg production over a seven-day collection period  $\pm$  SD. Average  
28 daily egg production across years is represented by a curved line. The procedure to derive daily  
29 production and averages is the same as that described in section 3.3 but the single tank was  
30 counted separately.



31

32

33

34