Gulf and Caribbean Research

Volume 2 | Issue 3

January 1968

A Supplement to the Fauna and Flora of Horn Island, Mississippi

E. Avery Richmond

Gulf Coast Research Laboratory

Follow this and additional works at: <https://aquila.usm.edu/gcr>

Part of the [Marine Biology Commons](#)

Recommended Citation

Richmond, E. 1968. A Supplement to the Fauna and Flora of Horn Island, Mississippi. *Gulf Research Reports* 2 (3): 213-254.

Retrieved from <https://aquila.usm.edu/gcr/vol2/iss3/1>

DOI: <https://doi.org/10.18785/grr.0203.01>

This Article is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in Gulf and Caribbean Research by an authorized editor of The Aquila Digital Community. For more information, please contact Joshua.Cromwell@usm.edu.

**A Supplement to the
Fauna and Flora of Horn Island,
Mississippi**

by

E. Avery Richmond

Gulf Coast Research Laboratory

TABLE OF CONTENTS

	Page
List of Figures.....	216
I. Introduction	219
II. Description of the Island.....	221
III. Historical Data.....	222
IV. Animal and Plant Lists.....	222
Fauna	225
Flora	246
References (additional since 1962).....	250
Corrigenda of "The Fauna and Flora of Horn Island, Mississippi" (Gulf Research Reports, Volume 1, No. 1, pp. 59-106, 1962)	253

LIST OF FIGURES

	Page
Figure 1. <i>Panicum amarum</i> Elliott — Bitter Grass.....	217
Figure 2. <i>Uniola paniculata</i> Linnaeus — Sea Oats.....	217
Figure 3. From Sound (Section 18).....	218
Figure 4. Looking West from Section 24.....	218
Figure 5. Winter fences before hurricane (Gulf side).....	227
Figure 6. Winter fences after hurricane (Gulf side).....	227
Figure 7. Typical scene of higher elevations in the interior of Horn Island (Slash Pine, Rosemary, Baccharis and Saw Palmetto).....	241
Figure 8. Tracks of <i>Procyon lotor</i> in sand.....	241
Figure 9. From Gulf to North (Section 29).....	251
Figure 10. Looking West—last of trees—near Section 24.....	251
Figure 11. <i>Euphorbia cordifolia</i> (Ell.) Small — Spurge.....	252
Figure 12. <i>Opuntia humifusa</i> Raf. — Rafinesque's Prickly Pear	252

Figure 1. *Panicum amarum* Elliott — Bitter Grass

Figure 2. *Uniola paniculata* Linnaeus — Sea Oats

Figure 3. From Sound (Section 18)

Figure 4. Looking West from Section 24

I. INTRODUCTION

Since the publication of "The Fauna and Flora of Horn Island, Mississippi" (Richmond 1962) search for unrecorded species present on this island, or in the waters surrounding it, has been continuing by myself and others. The premise that all of the plants and animals living in a given area can never be located completely is naturally well-founded. However, it seems advisable to continue the search for unlisted species of Horn Island.

Many unrecorded specimens have been collected and it seems proper at this time to make a supplementary list available to the public, particularly those interested in fields of science and nature. Knowing what is present on the island doubtless will tend to stimulate biological studies in various fields of endeavor.

The card catalogues of the Gulf Coast Research Laboratory Museum have been carefully examined by me through the courtesy of Mr. C. E. Dawson, Curator. Dr. Gordon Gunter kindly made some of his unpublished data of 1956 and 1957 available. Many visits to Horn Island have been made by the writer and others. In fact all members or students from the Laboratory have contributed in some way or other and, when known, credit is given to each one. The new contributors since the issue of my 1962 paper include W. I. Anderson, recently deceased, of the Shearwater Pottery Company. His expert knowledge of bird life has been especially helpful. David Peterson, manager of the Gulf Island National Wildlife Refuges since he replaced John H. Walther, K. E. Myers and Jerry E. Blackard, respectively managers of previous years, has been extremely cooperative. Frank Carroll, who has been associated with these managers, has assisted me materially. Marvin Phillips, formerly with the United States Army and stationed on the island, accompanied me on many trips during 1944 and 1945.

As usual, the facilities of Cornell University, the Academy of Natural Sciences of Philadelphia, the United States National Museum and the American Museum of Natural History have been available to me for identification. W. J. Gertsch of the American Museum identified the spiders. T. E. Bowman (United States National Museum), has checked a number of crustaceans as has Douglas Farrell of the Gulf Coast Research Laboratory. R. B. Channell (Vanderbilt University), Roland M. Harper (Alabama University) recently deceased, and E. T. Wherry (University of Pennsylvania) verified the names of many plants. Dr. Harper reports that *Pinus clausa* (Engelm.) Vasey, the spruce-pine of Florida, although present in Alabama, is otherwise restricted to Florida. A single specimen is growing on Horn Island according to Dr. Channell and is listed in my original paper (1962). Dr. Harper stated in 1928 that *Pinus clausa* is confined to the stationary dunes of pure white sand on the coast of Baldwin County, Alabama. There is no doubt that many of the plants originally established on Horn Island have disappeared and also that many others are constantly getting a foothold there. For instance, the cypress has gone and the chinaberry tree has arrived.

Again credit is extended to the specialists who kindly gave their time to verify or name the species contained in the lists given in

Chapter IV. My 1962 paper thanked many too briefly but I will not repeat the entire list. Many additional specialists have been interviewed and their courtesy is appreciated. Included are the following scientists: W. A. Connell, D. R. Davis, R. C. Froeschner, J. L. Herring, R. W. Hodges, J. M. Kingsolver, T. J. Spilman and Richard E. White of the United States National Museum; M. D. Leonard, retired, formerly with the Shell Oil Company; Miss E. Van Tassell (formerly at Catholic University, Washington, D. C.); Robert C. Graves (Bowling Green State University, Ohio) and L. L. Pechuman (Insect Curator, Cornell University).

Henry Yost, deceased, and J. Cowger of the Pest Control Laboratory, United States Department of Agriculture, at Gulfport, Mississippi, gave me some supplies and valuable suggestions. My retired friend, Henry Dietrich, former Curator of Entomology at Cornell University, identified many beetles, especially carabids. James E. Böhlke checked my determinations of a number of fresh water fishes. Leona W. Richmond typed most of my preliminary notes and took some very excellent photographs in connection with this study.

James S. Franks of the University of Mississippi and the Gulf Coast Research Laboratory has collected the fishes of the lagoons and ponds of Horn Island through the courtesy of the Fish and Wildlife Service of the U. S. Department of the Interior. He was aided financially by a modest N. S. F. grant obtained by the Gulf Coast Research Laboratory. A number of species, recently collected by him, have been added to my original list.

Dr. A. E. Schuyler helped particularly in the accurate determination of the sedges. H. W. Wilkens of Reading, Pennsylvania, called my attention to several plants not previously listed. Wilkens, visiting the Gulf Coast Research Laboratory in the spring and fall of 1965, did some collecting on the mainland but did not go to Horn Island. He reports that Richard L. Diener, formerly of Keesler Air Force Base, collected fifteen specimens on the island. Some of his specimens were not previously listed. E. P. Bicknell (1899) and E. E. Sherff (1933) each described a new species of plant from Horn Island. The species were collected by S. M. Tracy. As yet, I have not located *Coreopsis corninsularis* Sherff.

II. DESCRIPTION OF THE ISLAND

Horn Island is briefly described by Richmond (1962). As is well known, its conformity is constantly changing. This is particularly noticeable along the shoreline and from the changes in the shape and size of the lagoons and ponds. Evidently the southeastern portion is losing its foundation to the Gulf, whereas the northwestern area is gradually building up. This is in keeping with Richard R. Priddy's (1965) premise that the changes are due to the Mississippi River's power to move the barrier islands by "long shore" currents. These currents discharge southeastward and are diverted northward towards the islands and then westward. The result, he states, is that the islands are moving westward and to some extent southward.

The strong root system of the slash pine holds the central region (East and West) more or less in a stable equilibrium. Wax myrtle, yaupon, groundsel, rosemary, bitter beachgrass (Fig. 1) and sea oats (Fig. 2) doubtless have helped materially.

The dunes on the south side took a terrific beating in the fall of 1965. An attempt had been made by the Fish and Wildlife Service management to anchor those areas by bulldozing the sand into seemingly protective dunes. It was hoped that such a method, together with the use of winter fences (Fig. 5) would tend to alleviate the onslaught of the wind and Gulf waters. No real success was attained and the southern shore area was definitely flattened (Fig. 6). Most of the higher natural dunes held fast. LaGorce (1915) writes most interestingly about the ravages created by the Atlantic Ocean on our eastern coast.

During the passage of hurricane Betsy, heavy rains fell on lower Mississippi from the ninth to twelfth of September 1965. This rain, coupled with winds varying from 70 to 90 miles per hour, caused much shifting of the sands and the recently constructed dunes were leveled off. Raccoons, deer and other animals from all the islands were drowned and a number were washed up on the shore of the mainland. Much miscellaneous debris was dumped on the island. Tropical sea-bean seeds and a ten-foot palm tree bole were found near the area of the former army barracks.

The late W. I. Anderson, who was a well-known naturalist, artist and one of the owners of the Shearwater Pottery of Ocean Springs, attained one of his lifelong ambitions when he weathered this hurricane. He was wont to frequent the island for weeks at a time, as he communed in solitude with nature. Many ideas, emanating from his observations, were used in his pottery designs. At the time that Betsy was on a rampage, he was camping on his favorite so-called abode. When the water rose to his armpits, he tied the painter of his small boat about his waist and moved away from the rising waters to a higher level on one of the larger dunes. Darkness fell and he spent the night and following day safely in the leeward of the higher dunes until the storm abated. He saw his favorite pig washed away and drowned.

On 10 July 1966, an exhibition of some of Anderson's paintings of Horn Island scenes, plants and animals was staged at a motel in Pascagoula, Mississippi.

The tropical storm Debbie left a record of 16.85 inches of rainfall in downtown Mobile on September 29th and 30th of 1965. I have no information of its effect on Horn Island.

III. HISTORICAL DATA

Although Claiborne (1880) did not write about Horn Island, he did write quite extensively concerning Dauphin Island and Ship Island, which lie very close to Horn. Unfortunately his Volume II was destroyed by fire.

Guyton (1952) in a school textbook entitled "Our Mississippi" tells about Pierre LeMoyne d'Iberville's journey from France via Santo Domingo to Ship Island. On February 10, 1699, the expedition arrived. There, they were met by friendly Biloxi Indians. The Biloxis informed d'Iberville relative to a "near-by" mainland and especially about a large river to the west. Leaving his fleet anchored, he and his younger brother, Bienville, toured the coast and located the Mississippi River. Later they returned to Ship Island and at Old Biloxi (Ocean Springs) built Fort Maurepas on April 8, 1699. My previous paper (1962) states that Indians stayed "close to the mainland." Such is evidently incorrect in view of the presence of the Biloxis on Ship. Since the Gulf islands were separated only by relatively narrow expanses of water, it seems very likely that the Indians did roam on Horn. However, no evidence has been revealed concerning such peregrinations. As Thompson (1964) stated, "Horn Island is a blurred page in history." In 1965 Thompson again wrote about Horn Island and used some illustrations. The lighthouse, which was washed away in 1906, was pictured. Another unmanned lighthouse was erected on the north side of Petit Bois. The Pascagoula and Moss Point Chronicle (1965) published an interesting article about these lighthouses.

Since 1945, all military buildings noted in my first paper have been removed or destroyed by the weather elements. Only the power-house chimney in the operations area remains standing. In 1963 the Fish and Wildlife Service erected a 20 x 40 feet crew cabin on Horn Island near the area where the former headquarters of the military establishment was located during World War II. This cabin was erected six feet above a concrete flooring which supports the beams of the building. In 1965, a garage with a storeroom was built to house the vehicles and to supply extra storage space.

Transportation to and from the Island was made available by the Gulf Coast Research Laboratory or the Fish and Wildlife Service. The Laboratory trawler, HERMES, and the 30-foot SKIMMER of the Wildlife Service were used. Travel on the island was made on foot or by a Refugees jeep.

IV. ANIMAL AND PLANT LISTS

"There is not a property in nature but a mind is born to seek and find it, for it is not the plants or the animals, innumerable as they are, nor the whole magazine of material nature that give the sum power, but the infinite applicability

of these things in the hands of thinking man, every new application being equivalent to a new material."

—R. W. Emerson (1879) 1891

As in my 1962 paper, the species listed include only those animals and plants which were actually collected or sighted on Horn Island or in the waters around it out to a few fathoms of depth. The listings herein include only the records acquired from 1962 to 1966 inclusive.

A black-light trap was used occasionally instead of a New Jersey light trap for taking insects. However, its use did not prove entirely satisfactory. Hand-nets, pans, picking up and cupping proved more successful. In 1963, Kent Meyer ran the trap for several days during early April and two lots of collections were forwarded to me. An immediate kill was not obtained and most of the more delicate insects were ruined beyond identification. Cyanide was used as the killing agent. One other collection was made by Jerry Blackard on 14 April 1965. Ethyl acetate was used as the killing agent and the results proved more successful. Due to various climatic and labor difficulties, no further runs were made until June 1966. Now the use of 70% isopropyl alcohol and ethyl acetate keeps the collection in better shape.

The observance of a Barn Owl and the Coypu (*Nutria*) on the island in 1966 was called to my attention by David Peterson, J. S. Franks and others. Tracks of the River Otter were reported by Boyd Kynard and the otter is apparently quite common although I do not list it. Just when these animals moved in is not known. W. T. Anderson saw animals on drifting logs, boxes, etc. from time to time. Bond (1966) briefly discusses the transportation of small animals, invertebrates and birds by vegetative rafts on ocean waters. Hogs raised on the island before 1940 are still present. Trapping has reduced them somewhat.

Ruby-throated hummingbirds migrated through the Island in the Spring of 1964. They were very numerous and for a number of days fed quite commonly on purple thistle. In 1965, I arrived at the island very early in hopes of seeing their activity but no such migration occurred. None were reported in 1966. The nests of the Brown Pelicans on the North Islands were seriously depleted by adverse weather conditions in 1960 but lately the species appears to be on the increase. Several flocks were observed in 1966 and a colony of more than one hundred were seen using a group of spoil bank islands in the Sound just south of Pascagoula. However, their present nesting place is unknown.

A number of recently collected plants have been given to the Academy of Sciences in Philadelphia, as well as to the Bailey Hortorium of Ithaca, New York.

Aside from standard books and texts, publications by the following authors and institutions have been helpful in the studies of the animals and plants encountered in the Southern Mississippi area—Jenkins (1933), Behre (1950), Radford, Ahles and Bell (1964), Ralph Smith *et al.* (1964), Taylor (1960) and Williams (1964). Attention should be called to Richard Lane's (1957) excellent presentation of the history and activities of the Gulf Coast Research Laboratory.

No concentrated effort has been made to consider the protozoans, algae, mosses or ferns but some algae and ferns are listed. Humm and Caylor (1957) wrote extensively on the "Summer Marine Flora of Mississippi Sound" but nothing was specifically collected near Horn Island. F. Drouet of the Academy of Natural Sciences of Philadelphia identified several species of algae.

R. B. Channell's manuscript on "Vegetation of the West End of Horn Island" was most interesting and helpful to me and to members of the teaching staff at the Laboratory. He has also collected many species of algae on trips to Horn Island with his classes during recent years and has kindly given me their names. These species are listed.

Pictures in this paper will perhaps give the reader a broader vision of the plants and characteristic scenes encountered on Horn Island (Figs. 1-12).

An examination of Table No. 1 shows 1,568 species of plants and animals collected and identified from 1944 through 1966, 468 species since 1961. Naturally the insects predominate in numbers (712 species). The fishes follow with 156 species. Two hundred four species of plants are recorded.

TABLE 1
Number of Identified Members of the Fauna and Flora
Horn Island Records (1944-1966)

ANIMAL PHYLA AND MAJOR SUBDIVISIONS	FAMILIES	GENERA	SPECIES
Lower Invertebrates (through Annelida)	28	30	30
Mollusca	56	88	110
Arthropoda (Classes)			
Merostomata	1	1	1
Crustacea	46	71	94
Myriapoda	1	1	1
Insecta	157	478	712
Arachnida	15	24	28
Echinodermata	6	8	11
Prosopygia	1	1	1
Enteropneusta	1	1	1
Urochorda	1	1	1
Cephalochorda	1	1	1
Craniata (Classes)			
Pisces	64	114	156
Amphibia	2	5	10
Reptilia	10	16	23
Aves	38	131	176
Mammalia	6	6	8
Animals	434	977	1364
Plants	77	147	204
Total Collections	511	1124	1568

FAUNA

Phylum COELENTERATA

Class Hydromedusae

Hydractinidae

⁵*Hydractinia echinata* Fleming Spiny Polymorphic Hydroid

Class Scyphomedusae

Pelagidae

⁵*Chrysaora quinquecirrha* Desor Jellyfish

Rhizophysaliidae

Physalia pelagica Bosc Portuguese Man-of-War

Phylum PLATYHELMINTHES — Flatworms

Class Turbellaria — Planarians

Bdellouridae

Bdelloura candida (Girard) Planaria

Phylum ANNELIDA — Segmented Worms

Class Chaetopoda

Chaetopteridae

⁵*Chaetopterus* sp.

Megascolecidae

Pontedrilus bermudensis Beddard

Class Hirudinea — Leeches

Piscicolidae

Myzobdella lugubris Leidy Leech

Phylum MOLLUSCA — Mollusks

Class Pelecypoda — Clams

Teredinidae — Ship Worms

⁵*Teredo* sp. Ship Worm

Order Nudibranchia

Corambidae

Corambella baratariae Harry Barataria Nudibranch

Dorididae

Doris verrucosa Linnaeus Verucose Slug

Class Gastropoda — Snails (Univalves)

Fissurellidae

Diadora cayenensis Lamarck Cayenne Keyhole Limpet

Vitrinellidae

Cyclostremella humilis Bush Humble Cyclostremella

Melongenidae — Large Whelks

⁶*Busycon contrarium*

Strombidae

Strombus alatus Gmelin Florida Fighting Conch

Class Cephalopoda — Cuttlefishes

Loliginidae

Doryteuthis plei (de Blainville) Squid

Loligo pealei LeSueur Peale's Squid

Lolliguncula brevis de Blainville Short Squid

Phylum ARTHROPODA — Segmented Animals

Class Crustacea — Crustaceans

Subclass Copepoda

Clausidiidae

⁴*Clausidium* sp.

Lernaeidae

Lernaenicus radiatus (LeSueur)

Pontellidae

¹*Anomalocera ornata* Sutcliffe

Subclass Cirripedia

Lepadidae

¹*Lepas anatifera* Linnaeus Goose Barnacle

¹*Lepas pectinata* Spengler Pectinate Goose Barnacle

Subclass Malacostraca

Order Amphipoda

Ampeliscidae

Ampelisca holmesi Pearse

Ampithoidae

³*Ampithoe longimanus* Smith

Cymadusa filosa Savigny

Atylidae

Atylus minikai (A. C. Walker)

³*Atylus* sp.

Bateidae

³*Batea* sp.

Caprellidae

Caprella carolinensis Mayer

Carolina Caprella

³*Hemiaegena minuta* Mayer

Corophiidae

³*Corophium acherusicum* Costa

³*Erithonius brasiliensis* (Dana)

Figure 5. Winter fences before hurricane (Gulf side)

Figure 6. Winter fences after hurricane (Gulf side)

<i>Gammaridae</i>		
<i>Carinogammarus mucronatus</i> (Say)		
<i>Haustoriidae</i>		
³ <i>Haustorius mexicanus</i> Bousfield		
<i>Oedicerotidae</i>		
³ <i>Monoculodes edwardsi</i> Holmes		
<i>Photidae</i>		
³ <i>Microprotopus raneyi</i> Wigley		
<i>Talitridae</i> — Sandfleas		
³ <i>Orchestia grillus</i> (Bosc)		
³ <i>Orchestia platensis</i> Krøyer		
³ <i>Talorchestia</i> sp. No. 1		
³ <i>Talorchestia</i> sp. No. 2		
Order Isopoda		
<i>Cymothoidae</i>		
<i>Livoneca ovalis</i> (Say)		
<i>Idotheidae</i>		
<i>Erichsonella attenuata</i> (Harger)		
<i>Sphaeromidae</i>		
<i>Ancinus depressus</i> (Say)		
Order Stomatopoda		
<i>Squillidae</i>		
<i>Lysiosquilla excavatrix</i> Brooks		Mantis Shrimp
<i>Lysiosquilla scabricauda</i> (Lamarck)		Mantis Shrimp
Order Decapoda		
Suborder Macrura		
<i>Alpheidae</i> — Snapping Shrimp		
⁵ <i>Crangon</i> (<i>Alpheus</i>) <i>heterochelis</i> (Say)		
<i>Hippolytidae</i>		
<i>Hippolyte zostericola</i> (Smith)		Eel-Grass Shrimp
<i>Tozeuma carolinense</i> Kingsley	Carolina	Eel-Grass Shrimp
<i>Palaemonidae</i>		
<i>Palaemonetes vulgaris</i> Say		Common Grass Shrimp
<i>Penaeidae</i>		
⁵ <i>Trachypeneus constrictus</i> (Stimpson)		
<i>Ogyrididae</i>		
<i>Ogyrides alphaerostis</i> (Kingsley)		Ogyrides
<i>Processidae</i>		
<i>Processa</i> sp.		Processa
Suborder Brachyura		
<i>Calappidae</i>		
<i>Calappa sulcata</i> Rathbun		Box Crab

<i>Grapsidae</i>		
⁶ <i>Sesarma cinereum</i> (Bosc)		Wharf Crab
<i>Majidae</i> (Maiidae)		
¹ <i>Metoporphaphis calcarata</i> (Say)	"Calcareous"	Spider Crab
<i>Portunidae</i>		
<i>Ovalipes quadulpensis</i> (de Saussure)		Lady Crab
¹ <i>Portunus gibbesii</i> (Stimpson)		Swimming Crab
<i>Xanthidae</i>		
<i>Panopeus herbstii</i> (H. Milne Edwards)		Herbst's Mud Crab
⁵ <i>Panopeus</i> sp.		Mud Crab
Suborder Anomura		
<i>Albuneidae</i>		
<i>Albunea gibbesii</i> (Stimpson)		Sand Crab
<i>Lepidopa benedicti</i> Schmitt		Sand Crab
<i>Callianassidae</i>		
<i>Callianassa islagrande</i> Schmitt		Grand Isle Calianassa
<i>Paguridae</i>		
<i>Clibanarius vittatus</i> (Bosc)		Striped Hermit Crab
<i>Paguristes hummi</i> Wass		Hermit Crab
<i>Pagurus impressus</i> (Benedict)		Impressed Hermit Crab
<i>Petrochirus bahamensis</i> (Herbst)		Red Hermit Crab
<i>Porcellanidae</i>		
<i>Polyonyx gibbesi</i> (Haig)		Gibb's Polyonyx

¹Species collected by E. A. Richmond (1962-1966): ²H. J. Bennett; ³Douglas Farrell; ⁴Philip J. Phillips; ⁵Gordon Gunter—unpublished data; ⁶David Peterson. Remainder taken from a Gulf Coast Research Laboratory list.

CLASS INSECTA (HEXAPODA) — INSECTS

Odonata — Dragonflies, Damselflies

Lestidae

Lestes sp.

Libellulidae

Tarnetrum corruptum Hagen

Orthoptera — Grasshoppers, Crickets, Roaches, et al.

Gryllidae

Nemobius fasciatus (DeG.)

Hemiptera — True Bugs

Coreidae

Alydus pilosulus (Herrich-Schaeffer)

Hydrometridae

Hydrometra martini Kirkaldy

Lygaeidae
Pachybrachius servillei (Guerin)
Miridae
Psallus seriatus (Reuter)
Nepidae
Ranatra australis Hungerford
Notonectidae
Notonecta indica Linnaeus
Pentatomidae
Banasa dimidiata (Say)
Reduviidae
Rasahus hamatus (Fab.)
Homoptera — Aphids, Leafhoppers, et al.
Acanaloniidae
Acanalonia latifrons (Walker)
Aphidiidae
Aphis gossypii Glover
Prociphilus sp.
Cercopidae
Prosapia bicincta (Say)
Cicadellidae
Draeculacephala bradleyi Van Duzee
Draeculacephala portola Ball
Draeculacephala producta (Walker)
Tylozygus fasciatus (Walker)
Flatidae
Ormenoides venusta (Melichar)
Membracidae
Cyrtolobus tuberosa (Fairmaire)
Spissistilus festinus (Say)
Neuroptera — Lacewing-flies, antlions, et al.
Ascalaphidae
Ululodes hageni Weele
Chrysopidae
Chrysopa oculata Say
Myrmeleontidae
Paranthacclisis hageni (Bks)
Lepidoptera — Butterflies, moths, skippers
Suborder Heterocera, Superfamily Noctuoidea
Arctiidae — Tiger Moths
Apantesis nais Drury

Diacrisia virginica (Fab.)
Hyphantria cunea (Drury)
 Eucleidae — Slug Caterpillars
Euclea sp.
Sibine stimulea Clem.
 Olethreutidae
Bactra verutana verutana Zeller
 Phalaenidae (Noctuidae) — Noctuids
Acronicta tritona (Hbn.)
Euagrotis sp.
Euthisanotia unio Hbn.
Feltia subterranea (Fab.)
Leucania sp.
Orthodes crenulata Bth.
Tarachidia candefacta (Hbn)
 Geometridae — Geometrids
Semiothisa sp.
 Megalopygidae
Lagoa crispata Packard
 Pyrallidae
Diatraea sp.
Nomaphila noctuella (D. & G.)
 Gelechiidae
Aroga coloradella (Bsk.)
Dichomeris ligulella (Hbn.)
Filatima sp.
 Blastobasidae
Holcocera sp.
 Yponomeutidae
Atteva punctella (Cramer)
 Diptera — Flies, Mosquitoes
 Bibionidae
Philia orbata (Osten Sacken)
 Bombyliidae
Anthrax tigrina (DeGeer)
 Ceratopogonidae
Atrichopogon sp. (complex)
 Chironomidae
Chironomus sp.
 Dolichopodidae
Condyllostylus chrysoprasi (Walker)

Ephydridae

Dimecoenia spinosa (Loew.)

Setacea sp.

Sciomyzidae

Pherbellia nana (Fallen)

Stratiomyidae

Hermetia illucens (Linnaeus)

Tachinidae

Archytas apicifer (Walker)

Sitophaga sp.

Therevidae

Furcifera sp.

Coleoptera — Beetles

Alleculidae

Hymenorus densus LeConte

Hymenorus distinctus Fall

Anobiidae

Ernobius granulatus LeConte

Petalium seriatum Fall

Tricorynus sp. nr. *gravis* LeConte

Anthicidae

Tomoderus sp.

Vacusus laetus Laf.

Bostrichidae

Amphicerus cornutus Pallus

Lichenophanes armiger (LeConte)

Stephanopachys rugosus (Olivier)

Buprestidae

Taphrocerus sp.

Taphrocerus schaefferi Nic. and Weiss

Byrrhidae

Limnichites sp.

Carabidae

Agonoderus lineola (Fab.)

Agonoderus partiaris Say

Agonoderus pauperculus Dej.

Agonum cincticollis Say

Bembidion contractum Say

Bradycellus rupestris Say

Chlaenius laticollis Say

Chlaenius niger Rand.

Dyschirius erythrocerus Lec.

Euphorticus pubescens Dej.
Loxandrus sp.
Selonophorus sp.
Tachys sp.
Tetragonoderus fasciatus Hald.

Cerambycidae

Arhopalus rusticus nubilus (LeConte)
Leptostylus sp. (probably *knulli* Fisher)

Chrysomelidae

Altica amoena Horn
Altica chalybea Illiger
Altica rufa Illiger
Chrysomela scripta Fab.
Graphops curtipennis Melsh.
Lema trilineata Olivier
Rhadopterus sp.
Strabala rufa (Illiger)

Cicindelidae

Cicindela hamata lacerata Chd.
Cicindela hamata monti Vaurie

Coccinellidae

Ceratomegilla maculata DeGeer (*Coleomegilla maculata fus-*
cilabris Muls. = an aberration)
Chilocorus cacti Linnaeus
Chilocorus tripustulatus DeGeer
Hyperaspis signata Olivier

Colydiidae

Bothrideres geminatus (Say)

Cucujidae

Ahasverus rectus (DeGeer)

Curculionidae

Anthonomus sp.
Baris sp.
Hyperodes sp.
Listronotus blandus Henderson
Perigaster obscura LeConte
Sphenophorus necydaleides (Fab.)

Dytiscidae

Bidessus sp.
Copelatus glyphicus (Say)
Desmopachria grana (LeConte)

Elateridae

Blauta cribraria (Germar)
Conoderus amplicollis (Gyll.)
Conoderus falli Lane
Glyphonyx sp.
Hemicrepidius decolorata Say
Ischiodontus soleatus (Say)
Ischiodontus schwarzi Becker
Ischiodontus simplex (LeConte)
Lanelater sallei LeConte
Melanotus fissilis (Say)
Neotrichophorus carolinensis Schaeffer

Helodidae

Cyphon variabilis Thunb.
Scirtes tibialis Guerin

Heteroceridae

Heterocerus pallidus Say
Heterocerus pusillus Say

Hydrophilidae

Enochrus consors LeConte
Enochrus consortus Green

Lycidae

Celetes basalis LeConte

Monommidae

Hyporhagus punctulatus punctulatus Thomson

Melyridae

Collops floridanus Schaeffer
Collops sp.

Noteridae

Sulphisellus puncticollis Cresson

Omophronidae

Omophron sp.

Ostomidae

Temnochila virescens (Fab.)

Scarabaeidae

Ataenius cognatus LeConte
Ataenius gracilis Melsh.
Ataenius simulator Harold
Aphodius granarius Linnaeus
Diplotaxis bidentata LeConte
Euparia?

Parastasia brevipes (LeConte)

Phyllophaga dispar (Burm.)

Strategus julianus Burm.

Scolytidae

Xyleborus xylographus Say

Silphidae

Silpha surinamensis Fab.

Staphylinidae

Carpelinus sp.

Hesperobium sp.

Lathrobium simplex LeConte

Lobrathium sp.

Philonthus cunctans Horn

Tenebrionidae

Gondwanocrypticus obsoletus (Say)

Cybotus estriatus (LeConte)

Epitragodes tomentosus (LeConte)

Hymenoptera — Bees, Wasps, Ants, et al.

Braconidae — Braconids

Lysiphlebus testaceipes (Cresson)

Microplitis varicolor Viereck

Rogas laphygmae Viereck

Scoliidae — Scoliids Wasps

Campsomeris plumipes fossulana (Fab)

CLASS ARACHNIDA*

Araneida — Spiders

Dictynidae — Dictynids

Dictyna sublata Hentz

Drassidae — Drassids

Gnaphosa sericata Koch

Linyphiidae — Sheet-web Weavers

Ceraticelus similis Banks

Erigone autumnalis Emerton

Araneidae — Typical Orb-weavers

Araniella displicata (Hentz)

Argiopidae — Orb-weavers

Tetragnatha caudata Emerton

Tetragnatha pallescens (Cambridge)

Tetragnatha sp.

*—Collected by Richmond and deposited in the American Museum of Natural History, N. Y.

Thomisidae — Crab-spiders

Misumenops celer Hentz

Tibellus duttonii Hentz

Clubionidae — Clubionids

Meriola decepta Banks

Lycosidae — Wolf-spiders

Arctosa sublata Montgomery

Lycosa antelucana Montgomery

Pardosa pauxilla Montgomery

Pardosa saxatilis Hentz

Attidae — Jumping-spiders

Pellenes coronatus Hentz

Phylum ECHINODERMATA

Class Holothuroidea

Cucumariidae

⁵*Thyone mexicana*

Class Asteroidea

Arbaciidae

¹⁰*Arbacia punctulata* (Lamarck)

Purple Sea Urchin

Phylum PROSOPYGIA

Class Ectoprocta

Vesiculariidae

⁵*Amathia convoluta* Lamouroux

Phylum CHORDATA

Subphylum Cephalochordata (Acrania)

Branchiostomidae

Branchiostoma floridae Hubbs Florida Lancelet, Amphioxus

Subphylum Craniata (Vertebrata)

Class Pisces

Subclass Chondrichthyes — Cartilaginous Fishes

Order Squaliformes

Carcharhinidae — Requiem Sharks

^{1/11}*Carcharinus leucas* (Müller and Henle)

Bull Shark

^{1/11}*Carcharinus limbatus* (Müller and Henle)

Blacktip Shark

¹⁶*Negaprion brevirostris* (Poey)

Lemon Shark

^{1/11}*Scoliodon terraenovae* (Richardson) Atlantic Sharpnose Shark

Sphyrnidae

^{15/16}*Sphyrna tiburo* (Linnaeus)

Bonnethead

Order Rajiformes (Batoidei)		
Pristidae — Sawfishes		
^{12/14} <i>Pristis pectinatus</i> Latham		Common Sawfish
Torpedinidae — Electric Rays		
^{14/15/17} <i>Narcine brasiliensis</i> (Olfers)		Lesser Electric Ray
Rajidae — Skates		
^{6/16} <i>Raja eglanteria</i> Bosc		Skate, Brier Ray
¹⁷ <i>Raja lentiginosa</i> Bigelow and Schroeder		Freckled Skate
Dasyatidae		
¹⁷ <i>Dasyatis americana</i> Hildebrand and Schroeder		Southern Stingray
¹⁷ <i>Gymnura micrura</i> (Bloch and Schneider)		Smooth Butterfly Ray
Subclass Osteichthyes — Bony Fishes		
Order Seminotiformes		
Lepisosteidae		
¹⁶ <i>Lepisosteus oculatus</i> (Winchell)		Spotted Gar
Order Clupeiformes		
Elopidae — Ladyfishes		
¹⁶ <i>Elops saurus</i> Linnaeus		Ladyfish
Megalopidae — Tarpons		
¹⁶ <i>Megalops atlantica</i> Valenciennes		Atlantic Tarpon
Clupeidae — Herrings		
^{1/16} <i>Dorosoma petenense</i> (Günther)		Threadfin Shad
Engraulidae		
¹⁶ <i>Anchoa mitchilli</i> (Valenciennes)		Bay Anchovy
Order Cyprinodontiformes		
Cyprinodontidae — Killifishes — Top Minnows		
^{15/17} <i>Adinia xenica</i> (Jordan and Gilbert)		Diamond Killifish
^{15/17} <i>Fundulus confluentus</i> Goode and Bean		Marsh Killifish
¹⁶ <i>Fundulus pulvereus</i> (Evermann)		Bayou Killifish
Order Gadiformes		
Gadidae — Codfishes		
¹⁶ <i>Urophycis floridanus</i> (Bean and Dresel)		Southern Hake
Order Gasterosteiformes		
Fistulariidae — Cornetfishes		
⁴ <i>Fistularia tabacaria</i> Linnaeus		Cornetfish
Syngnathidae — Pipefishes, Seahorses		
⁶ <i>Syngnathus scovelli</i> (Evermann and Kendall)		Scovell's Pipefish

Order Perciformes

Serranidae — Sea Basses

¹⁷ <i>Centropristes ocyurus</i> (Jordan and Evermann)	Bank Sea Bass
⁴ <i>Centropristes philadelphicus</i> (Linnaeus)	Rock Sea Bass
¹ <i>Diplectrum arcuarium</i> Ginsburg	Sandfish
⁹ <i>Mycteroperca bonaci</i> (Poey)	Black Grouper
⁸ <i>Mycteroperca phenax</i> (Jordan and Swain)	Scamp
⁶ <i>Serraniculus pumilio</i> Ginsburg	Pigmy Sea Bass

Lutjanidae — Snappers

¹⁷ <i>Lutjanus campechanus</i>	Red Snapper
^{1/10} <i>Lutjanus synagris</i> (Linnaeus)	Lane Snapper

Priacanthidae

¹⁷ <i>Pristigenys alta</i> (Gill)	Short Bigeye
--	--------------

Pomatomidae — Bluefishes

^{1/4} <i>Pomatomus saltatrix</i> (Linnaeus)	Bluefish
--	----------

Carangidae — Jacks

⁴ <i>Alectis crinitis</i> (Mitchill)	Threadfish
⁴ <i>Caranx hippos</i> (Linnaeus)	Common Jack
¹⁶ <i>Caranx crysos</i> (Mitchill)	Blue Runner
⁴ <i>Chloroscombrus chrysurus</i> (Linnaeus)	Bumper

Pomadasyidae — Grunts

^{1/14/ 16} <i>Orthopristis chrysopterus</i> (Linnaeus)	Pigfish
---	---------

Sciaenidae — Drums

¹¹ <i>Menticirrhus americanus</i> (Linnaeus)	Southern Kingfish
¹⁶ <i>Stellifer lanceolatus</i> (Holbrook)	Star Drum
¹⁷ <i>Cynoscion arenarius</i> Ginsburg	Sand Sea Trout
¹⁷ <i>Cynoscion nothus</i> (Holbrook)	Silver Sea Trout

Ephippidae — Spadefishes

¹² <i>Chaetodipterus faber</i> (Broussonet)	Spadefish
--	-----------

Pomacentridae — Damselfishes

^{7/8} <i>Abudefduf saxatilis</i> (Linnaeus)	Sergeant Major
--	----------------

Labridae — Wrasse

¹⁷ <i>Halichoeres radiatus</i> (Linnaeus)	Puddingwife
¹¹ <i>Halichoeres caudalis</i> (Poey)	Painted Wrasse
^{13/16} <i>Hemipteronotus novacula</i> (Linnaeus)	Razorfish

Scombridae — Tunas, Mackerels

⁴ <i>Scomberomorus maculatus</i> (Mitchill)	Spanish Mackerel
--	------------------

Xiphiidae — Swordfishes

¹⁴ <i>Xiphias gladius</i> Linnaeus	Swordfish
---	-----------

Eleotridae — Sleepers

^{15/16} <i>Eleotris pisonis</i> (Gmelin)	Spinycheck Sleeper
---	--------------------

Gobiidae — Gobies

^{4/17} <i>Gobioides broussonneti</i> Lacépède	Violet Goby
¹⁶ <i>Gobionellus boleosoma</i> (Jordan and Gilbert)	Darter Goby
¹ <i>Gobiosoma longipala</i> Ginsburg	Naked Goby
¹⁷ <i>Gobiosoma</i> sp.	Goby
¹⁶ <i>Gobiosoma bosci</i> (Lacépède)	Naked Goby
¹ <i>Gobiosoma robustum</i> Ginsburg	Naked Goby
¹⁶ <i>Microgobius gulosus</i> (Girard)	Large-mouthed Goby
¹⁵ <i>Evorthodus lyricus</i> (Girard)	Lyre Goby

Scorpaenidae — Rockfishes, Scorpionfishes

¹⁷ <i>Scorpaena dispar</i> Longley and Hildebrand	Hunchback Scorpionfish
¹¹ <i>Scorpaena grandicornis</i> Cuvier	Lionfish
^{11/17} <i>Scorpaena plumieri</i> Bloch	Scorpion Fish

Triglidae

¹⁷ <i>Prionotus martis</i> Ginsburg	Barred Searobin
¹⁷ <i>Prionotus rubio</i> Jordan	Blackfin Searobin
^{16/17} <i>Prionotus scitulus</i> Jordan and Gilbert	Slender Searobin
¹⁷ <i>Prionotus tribulus</i> Cuvier	Bighead Searobin
¹⁷ <i>Prionotus</i> sp.	Searobin

Blenniidae — Combtooth Blennies

⁴ <i>Hypsoblennius hentzi</i> (LeSueur)	Feather Blenny
⁴ <i>Hypsoblennius ionthas</i> (Jordan and Gilbert)	Freckled Blenny

Ophidiidae — Cusk-eels

¹ <i>Lepophidium graellsii</i> (Poey)	Blackedge Cusk-eel
¹ <i>Ophidion</i> sp.	Cusk-eel
¹ <i>Ophidion welshi</i> (Nichols and Breder)	Crested Cusk-eel
^{15/16} <i>Otophidium</i> sp.	Cusk-eel

Sphyraenidae — Barracudas

^{7/16} <i>Sphyraena barracuda</i> (Walbaum)	Great Barracuda
^{15/16} <i>Sphyraena guachancho</i> Cuvier	Guachanche

Order Pleuronectiformes

Bothidae — Lefteye Flounders

¹⁷ <i>Ancyllopsetta quadrocellata</i> Gill	Ocellated Flounder
⁴ <i>Citharichthys macrops</i> Dresel	Spotted Whiff
^{16/17} <i>Etropus crossotus</i> Jordan and Gilbert	Fringed Flounder
⁴ <i>Syacium gunteri</i> Ginsburg	Gunter's Flounder

Soleidae

¹⁷ <i>Achirus lineatus</i> (Linnaeus)	Lined Sole
^{16/17} <i>Trinectes maculatus</i> Bloch and Schneider	Broad Sole, Hogchoker

Order Echeineiformes

Echeneidae — Remoras

¹⁷ <i>Echeneis naucrates</i> Linnaeus	Sharksucker
--	-------------

Figure 7. Typical scene of higher elevations in the interior of Horn Island (Slash Pine, Rosemary, Baccharis and Saw Palmetto)

Figure 8. Tracks of *Procyon lotor* in sand

Order Gobiesociformes

Gobiesocidae — Clingfishes

^{15/16}*Gobiosoma strumosus* Cope Skilletfish

Order Tetraodontiformes

Balistidae — Filefishes, Triggerfishes

^{4/17}*Aluterus schoepfi* (Walbaum) Orange Filefish

¹⁷*Balistes capricornis* Gmelin Gray Triggerfish

^{5/17}*Monacanthus ciliatus* (Mitchill) Fringed Filefish

^{1/17}*Monacanthus hispidus* (Linnaeus) Common Filefish

Tetraodontidae — Puffers

¹⁷*Sphaeroides nephelus* (Goode and Bean) Southern Puffer

Ostraciidae — Trunkfishes

^{2/11}*Lactophrys quadricornis* (Linnaeus) Cowfish

Order Batrachoidiformes

Batrachoididae — Toadfishes

²*Opsanus beta* (Goode and Bean) Toadfish, Oyster Dog

^{15/16}*Porichthys porosissimus* (Cuvier) Atlantic Midshipman

Order Lophiiformes

Antennariidae — Frogfishes

^{13/16}*Antennarius radiosus* Garman Singlespot Frogfish

⁴*Histrio histrio* (Linnaeus) Sargassum Fish

Ogcocephalidae — Batfishes

¹⁷*Ogcocephalus vespertilio* (Linnaeus) Longnose Batfish

⁶*Ogcocephalus* sp. Batfish

The numbers preceding the species indicate by whom the collection was made. Numbers 1, 2 and 3 are identical with the numbers used in my 1962 paper. The remainder refer to other collectors or references as follows: 4. Gulf Coast Research Laboratory; 5 C. E. Dawson; 6. H. Shoemaker (1955); 7. A. Myrberg; 8. H. T. Boschung; 9. Rohr, Ford and Brewster; 10. W. Demoran; 11. J. Y. Christmas; 12. W. Walley; 13. Gordon Garwood; 14. Wayne Watkins; 15. James B. Ward; 16. James Franks; 17. Gordon Gunter—unpublished data.

Class Amphibia — The Frogs, Toads, Salamanders

Hylidae

Hyla crucifer crucifer Wied.

Hyla femoralis Latreille

Hyla gratiosa LeConte

Pseudacris nigrata nigrata LeConte

Pseudacris ornata (Holbrook)

Sceloporus undulatus undulatus Latreille

Class Reptilia — The Reptiles

Testudinata — Turtles

Chelydridae

³ <i>Chelydra serpentina serpentina</i> (Linnaeus)	Common Snapping Turtle
---	------------------------

Class Aves — Birds

Colymbidae — Grebes

⁵ <i>Colymbus auritus</i> Linnaeus	Horned Grebe	W
---	--------------	---

Ardeidae — Herons and Bitterns

³ <i>Ardea occidentalis occidentalis</i> Audubon	Great White Heron	T
⁵ <i>Botaurus lentiginosus</i> (Baclett)	American Bittern	S
⁵ <i>Ardeola ibis ibis</i> (Linnaeus)	Cattle Egret	S
^{1/3} <i>Dicromanassa rufescens rufescens</i> (Gmelin)	Reddish Egret	P
^{3/5} <i>Florida caerulea caerulea</i> (Linnaeus)	Little Blue Heron	S
¹ <i>Ixobrychus exilis exilis</i> (Gmelin)	Eastern Least Bittern	W

Anatidae — Swans, Geese and Ducks

^{3/5} <i>Anas strepera</i> Linnaeus	Gadwell	W
⁵ <i>Charitonetta albeola</i> (Linnaeus)	Bufflehead	W
⁵ <i>Glaucionetta clangula americana</i> (Bonaparte)	American Goldeneye	W
³ <i>Lophodytes cucullatus</i> (Linnaeus)	Hooded Merganser	W
⁵ <i>Melanitta perspicillata</i> (Linnaeus)	Surf Scoter	M
⁵ <i>Mergus merganser merganser</i> Cassin	American Merganser	W
⁵ <i>Perissonetta collaris</i> (Donovan)	Ring-necked Duck	W
⁵ <i>Spatula clypeata</i> (Linnaeus)	Shoveler	M

Rallidae — Gallinules and Rails

⁴ <i>Laterallus jamaicensis jamaicensis</i> (Gmelin)	Black Rail	M,W
^{3/5} <i>Porzana carolina</i> (Linnaeus)	Sora Rail	T,W

Charadriidae — Plovers, Turnstones and Surfbirds

^{1/5} <i>Charadrius alexandrinus tenuirostris</i> (Lawrence)	Cuban Snowy Plover	S
---	--------------------	---

Scolopacidae — Woodcocks, Snipes and Sandpipers

³ <i>Erolia maritima</i> (Brünnich)	Purple Sandpiper	T
^{1/5} <i>Limnodromus griseus griseus</i> (Gmelin)	Eastern Dowitcher	T

Laridae — Gulls and Terns

³ <i>Rissa tridactyla tridactyla</i> (Linnaeus)	Atlantic Kittiwake	T
⁵ <i>Thalasseus sandvicensis acuflavidus</i> (Cabot)	Cabot Tern	S

Columbidae — Pigeons and Doves

³ <i>Columba livia livia</i> Gmelin	Rock or Domestic Pigeon	W
⁵ <i>Zenaida asiatica asiatica</i> (Linnaeus)	White-winged Dove	T

Tytonidae

⁶ <i>Tyto alba pratincola</i> (Bonaparte)	Barn Owl	P
--	----------	---

Caprimulgidae — Goatsuckers			
^{1/3/5} <i>Caprimulgus carolinensis</i> Gmelin	Chuck-will's Widow	W	
³ <i>Caprimulgus vociferus vociferus</i> Wilson	Eastern Whippoorwill	T	
Trochilidae — Hummingbirds			
^{3/5} <i>Archilochus colubris</i> (Linnaeus)	Ruby-throated Hummingbird	M,T	
Picidae — Woodpeckers			
³ <i>Dendrocopos pubescens pubescens</i> (Linnaeus)	Southern Downy Woodpecker	P	
⁵ <i>Melanerpes erythrocephalus</i> (Linnaeus)	Eastern Red-headed Woodpecker	P,V,S	
Tyrannidae — Flycatcher			
³ <i>Empidonax minimus</i> (Baird and Baird)	Least Flycatcher	M	
Corvidae — Crows and Jays			
³ <i>Corvus ossifragus</i> Wilson	Fish Crow	P	
³ <i>Cyanocitta cristata cristata</i> (Linnaeus)	Southern Blue Jay	P	
Paridae — Titmice			
³ <i>Penthestes carolinensis guilloti</i> Oberholser	Louisiana Chickadee	P	
Troglodytidae — Wrens			
³ <i>Telmatodytes palustris palustris</i> (Wilson)	Long-billed Marsh Wren	P	
³ <i>Nannus troglodytes hiemalis</i> (Vieillot)	Eastern Winter Wren	W	
Mimidae — Mockingbirds and Thrashers			
³ <i>Toxostoma rufum rufum</i> (Linnaeus)	Brown Thrasher	M,W	
Vireonidae — Vireos			
³ <i>Vireo sylvia gilva gilva</i> (Vieillot)	Eastern Warbling Vireo	M	
Parulidae — (Compsothlypidae) — Wood Warblers			
³ <i>Dendroica caerulescens</i> (Gmelin)	Black-throated Blue Warbler	M	
³ <i>Seiurus motacilla</i> (Vieillot)	Louisiana Waterthrush	S	
³ <i>Wilsonia citrina</i> (Boddaert)	Hooded Warbler	M,S	
Icteridae — Meadowlarks, Blackbirds and Oriole			
^{1/3/5} <i>Icterus spurius</i> (Linnaeus)	Orchard Oriole	S,M	
⁵ <i>Icterus galbula</i> (Linnaeus)	Baltimore Oriole	M	
³ <i>Sturnella magna argutula</i> Bangs	Southern Meadowlark	W,T	
Thraupidae — Tanagers			
^{1/3/5} <i>Piranga olivacea</i> (Gmelin)	Scarlet Tanager	M	
^{3/5} <i>Piranaga rubra rubra</i> (Linnaeus)	Summer Tanager	M,T	
Fringillidae — Grosbeaks, Finches, Sparrows and Buntings			

- ^{2/1}*Passerina ciris ciris* (Linnaeus) Painted Bunting M,W
³*Zonotrichia albicollis* (Gmelin) White-throated Sparrow M,W

Additions by the following observers to the 1962 list, published in Volume 1 (no.2) of the Gulf Research Reports.

1. E. A. Richmond; 2. J. R. Walther; 3. W. I. Anderson; 4. K. E. Myers; 5. J. Blackard; 6. D. Peterson; 7. F. Carroll.

M = Migrant, P = Permanent resident, S = Summer resident, T = Transient or winter visitant, V = Summer visitant, W = Winter resident.

Class Mammalia — The Mammals

Order Chiroptera — Bats

Vespertilionidae — Vespertilionid — Bats

Myotis lucifugus lucifugus (LeConte) Little Brown Myotis

Order Rodentia — Rodents

Capromyidae — Nutrias and Coypus

Myocastor coypus bronariensis

(E. Geoffrey St. — Hilaire) Coypus

Muridae

Mus musculus (Linnaeus)

House Mouse

FLORA

THALLOPHYTA

Phaeophyta — Brown Algae

Dictyotaceae

²*Dictyota dichotoma* (Hudson) Lamouroux

Dictyota

Ectocarpaceae

Ectocarpus confervoides (Roth) LeJod.

Ectocarpus

²*Ectocarpus siliculosus* (Dillwa.) Lyngb.

Ectocarpus

Sargassaceae

²*Sargassum fluitans* Borgesen

Sargassum

²*Sargassum natans* (Linnaeus)

Sargassum

Charophyta — Brittleworts

Characeae

²*Chara* sp.

Chara

Chlorophyta — Green Algae

Caulerpaceae

²*Caulerpa prolifera* (Forsk.) Lamouroux

Caulerpa

Cladophoraceae

²*Chaetomorpha gracilis* Kützing

Chaetomorpha

² <i>Cladophora blomquistii</i> Aziz and Humm	Cladophora
² <i>Cladophora gracilis</i> (Griffiths) Kütz.	Cladophora
<i>Ulvaceae</i>	
<i>Enteromorpha prolifera</i> (Müll.) J. Ag.	Enteromorpha
<i>Rhodophyta</i> — Red Algae	
<i>Ceramiales</i>	
² <i>Spyridia filamentosa</i> (Wulfen) Harvey	Spyridia
<i>Champiaceae</i>	
² <i>Champia parvula</i> (C. Agardh) Harvey	Champia
<i>Chantransiaceae</i>	
² <i>Acrochaetium seriatum</i> Borgesen	Acrochaetium
<i>Gracilariaceae</i>	
² <i>Gracilaria verrucosa</i> (Hudson) Papenfuss	Gracilaria
<i>Hypneaceae</i>	
² <i>Hypnea musciformis</i> (Wulfen) Lamouroux	Hypnea
<i>Rhodomelaceae</i>	
² <i>Chondria baileyana</i> (Montague) Harvey	Chondria
² <i>Chondria cnicophylla</i> (Melvill) De Toni	Chondria
² <i>Chondria littoralis</i> Harvey	Chondria
² <i>Laurencia poitei</i> (Lamouroux) Howe	Laurencia
<i>Cyanophyta</i> — Bluegreen Algae	
<i>Myxophyceae</i>	
² <i>Dichothrix penicillata</i> Zanardini	Dichothrix
<i>Pteridophyta</i>	
<i>Osmundaceae</i>	
<i>Osmunda regalis</i> Linnaeus	Royal Fern
<i>Spermatophyta</i>	
<i>Gramineae</i>	
<i>Andropogon maritimus</i> Chapman	Seaside Broom Grass
⁵ <i>Aristida spiciformis</i> Elliott	"Spike-like" Awn Grass
<i>Cynodon dactylon</i> (Kuntze)	Bermuda Grass
<i>Distichlis spicata</i> (Linnaeus) Greene	Spike-grass
[*] <i>Echinochla</i> (<i>Panicum</i>) <i>crus-galli</i> (L.) Beauvoir	Barnyard-grass
<i>Cyperaceae</i>	
<i>Elocharis acicularis</i> (L.) R. + S.	
<i>Fimbristylus spadicea</i> (L.) Vahl.	Chestnut-colored Sedge
(Plants previously identified as <i>F. Harperi</i> and <i>F. castanea</i>)	

*—Listed previously but generic name changed 2. Collected by R. B. Channell (1966); 4. collected by S. M. Tracy (1898-1903); 5. collected by D. L. Diener (1953). Unnumbered species collected by Richmond (1962-1966). Deposited in the Bailey Hortorium at Cornell University or at the Philadelphia Academy of Natural Sciences.

are probably *F. spadicea* according to A. E. Schuyler)
⁵*Fuirena scirpoides* Michx. Umbrella-grass

Xyridaceae

Xyris flabelliformis Chapman Flabellate Yellow-eyed grass

Haemodoraceae

Lacnanthes tinctoria (Walt.) Ell. Red Root

Amaryllidaceae

Allium canadense Linnaeus Wild Garlic

Iridaceae

⁴*Sisyrinchium atlanticum* Bicknell Atlantic Blue-eyed Grass

⁴*Sisyrinchium nanum* Bicknell Blue-eyed Grass

Orchidaceae

Spiranthes floridana Wherry Ladies Tresses

Polygonaceae

**Polygonella gracilis* (Nutt.) Meisner Slender Jointweed
 (Same as *Delopyrum gracilis* Nutt.)

Polygonum opelousanum Riddell Opelousas Smartweed

Amaranthaceae

Achyranthes philoxeroides (Mart.) Standley Chaff Flower

Capparidaceae

**Polanisia tenuifolia* Torrey & Gray Caper

Cruciferae

Cakile fusiformis Greene Cakile

Lepidium virginicum Linnaeus Poor-man's Pepper

Rosaceae

Rubus mississippianus Bailey Southern Dewberry
 (Listed previously as *R. trivialis* (Michaux))

Oxalidaceae

Xanthoxalis filipes Small Slender-stalked Oxalis

Euphorbiaceae

Euphorbia cordifolia (Ell.) Small Spurge

Euphorbia sp. Spurge

Malvaceae

Kosteletzyka virginica (L.) Presl. Seashore-mallow

Kosteletzyka althaeifolia (Chap.) A. Gray Seashore-mallow

Cistaceae

Helianthemum arenicola Chapman Sanddune Frostweed

Helianthemum canadense (L.) Michaux Frostweed

Helianthemum georgianum Chapman Rock Rose

Lechea tenuifolia Michaux Rock Rose

<i>Violaceae</i>		
<i>Viola vittata</i> Greene		White Violet
<i>Melastomataceae</i>		
<i>Rhexia alifanus</i> Walt.		Meadow-beauty
<i>Cornaceae</i>		
<i>Svida stricta</i> Lamarck (Small)		Lamarck's Dogwood
<i>Umbelliferae</i>		
<i>Ptilimnium capillaceum</i> (Michaux) Ref.		Mock's Bishop's Weed
<i>Ericaceae</i>		
<i>Vaccinium Darrowi</i> Camp		Darrow's Blueberry
<i>Primulaceae</i>		
<i>Centrunculus minimum</i> Linnaeus		Small's Chaffweed
<i>Convolvulaceae</i>		
<i>Ipomoea sagittata</i> Cav.		Sagittate Morning-glory
<i>Scrophulariaceae</i>		
<i>Linaria floridana</i> Chapman		Florida Toadflax
<i>Lentibulariaceae</i>		
<i>Pinguicula lutea</i> Walter		Yellow Butterwort
<i>Utricularia subulata</i> Linnaeus		Awl-shaped Bladderwort
<i>Compositae</i>		
<i>Cirsium horridulum</i> Elliotti		Purple Thistle
(Forma Elliotti (T. & G.) Fernald)		
⁴ <i>Coreopsis corninsularis</i> Sherff		Horn Island Tick-seed
<i>Erechtites hieracifolia</i> (L.) Raf.		Pilewort, Fireweed
<i>Helenium amarum</i> (Rab.) H. Rock		Sneezeweed

ADDITIONAL REFERENCES

(Since 1962 paper)

- Behre, Ellinor.* 1950. Annotated list of the fauna of the Grand Isle Region 1928-1946. Occasional Papers of the Marine Laboratory, Louisiana State University, Baton Rouge, Louisiana. No. 6: 1-66. (Includes Marine algae and higher plants).
- Bicknell, Eugene P.* 1899. Studies in *Sisyrinchium*. VI. Additional new species from the southern states. Bull. Torrey Botanical Club. 26(12): 605-616.
- Bond, James.* 1966. Eleventh supplement to the checklist of birds of the West Indies (1956). The Acad. Nat. Sci. Phila. pp. 1-13.
- Chronicle, Pascagoula and Moss Point, Miss.* 1965. Lighthouse keeper, part of island lost in 1906 hurricane. August 5: p. 14, figs. 1-2.
- Claiborne, J. F. H.* 1880. Mississippi as a Province, Territory and State. Vol. 1: i-xxix, 1-545. (Vol. 2: burned up).
- Emerson, Ralph Waldo.* 1891. The fortune of the republic. Miscellaneous Riverside Press, Cambridge, Mass. p. 332. (originally published in 1879)
- Guyton, Pearl V.* 1952. Our Mississippi. Steck Co. Austin, Texas. pp. 1-467, i-xi, fig. 5.
- Harper, Roland M.* 1928. Catalogue of the trees, shrubs and vines of Alabama with their economic properties and local distribution. Monograph 9. Economic Botany of Alabama. Part 2: pp. 1-357, figs. 1-66, maps 1-23 (See page 60).
- Humm, Harold J. and R. L. Caylor.* 1957. The summer marine flora of Mississippi Sound. Pub. Inst. of Marine Sci. 4(2): 228-264, fig. 1, pls. I-IX.
- Jenkins, Winchester* 1933. Wildlife of Mississippi from 45 years experience. Reporter Printing Co., Natchez, Miss. pp. 1-155.
- LaGorce, John O.* 1915. The warfare on our Eastern Coast. The Nat. Geog. Mag. XXVIII (3): 195-230, 29 ills. 2 charts.
- Lane, Richard* 1957. Prying secrets from the sea. The Commercial Appeal, Section V (18Aug): pp. 1 and 3, fig. 5.
- Priddy, Richard R.* 1965. Islands in Mississippi Sound are being shoved all around. Clarion-Ledger, Jackson, Miss. 8 July 1965 (Sect. B): 5.
- Radford, Albert E., Harry E. Ahles and C. R. Bell.* 1964. Guide to the Carolinas with distribution in the Southeastern States. Univ. N. C., Chapel Hill. pp. 1-363.
- Richmond, E. Avery.* 1962. The fauna and flora of Horn Island, Mississippi. Gulf Research Reports 1(2): 59-106, figs. 1-27, maps 1, 2, 13.
- Sherff, Earl E.* 1933. Newer or otherwise noteworthy Compositae. Botanical Gazette. 94(3): 589-597.

Figure 9. From Gulf to North (Section 29)

Figure 10. Looking West — last of trees — near Section 24

Figure 11. *Euphorbia cordifolia* (Ell.) Small — Spurge

Figure 12. *Opuntia humifusa* Raf. — Rafinesque's Prickly Pear

- Smith, Ralph, et al. 1964. Keys to marine invertebrates of the Woods Hole Region. Systematics—Ecology Program, Marine Biol. Lab., Woods Hole, Mass. Contrib. no. 11: Pref. i-x, pp. 1-208, pls. 1-28.
- Taylor, W. R. 1960. Marine algae of the eastern tropical and subtropical coasts of the Americas. Univ. of Mich. Press. Univ. of Michigan Studies. Ann Arbor. Scientific Series. Vol XXI: VII-IX, 1-870, pls. 1-80, figs. 1-14.
- Thompson, Ray M. 1964. Know your state. Horn Island—a blurred page of history. Biloxi-Gulfport Daily Herald, 24 April 1964: 16.
- Thompson, Ray M. 1965. Horn Island, Down South on the beautiful Gulf Coast. Vol 15(2): 4-5, illus.
- Williams, Austin B. 1965. Marine decapod crustaceans of the Carolinas. Fishery Bull. U. S. Dept. of Interior, Fish and Wildlife Service, Bur. Comm. Fisheries 65(1): 1-298, figs. 1-252.

CORRIGENDA

Of

"The Fauna and Flora of Horn Island, Mississippi,"
(Gulf Research Reports, Volume 1, No. 2, pp. 59-106, 1962)

- p. 62, line 1: Fields = Field
- p. 70, line 37: *Andara* = *Anadara*
- p. 71, line 36: *Marcrocallista* = *Macrocallista*
- p. 74, line 23: *Malacostra* = *Malacostraca*
- p. 74, line 30: *Taylorchestia longicornis* = *Talorchestia* sp.
- p. 74, line 43: *ocessatus* = *ocellatus*
- p. 75, line 7: *Emargonarta* = *emarginata*
- p. 78, line 4: *Lygus apicalis* Fieber = *Taylorilygus pallidulus* Blanchard
- p. 78, line 19: *Pangaeus bilineatus* (Say) = *Cydnidae*
- p. 78, line 40: *Graminella nigrifrons* = Delete "Probably *fascifrons* (Stål)"
- p. 78, line 43: *Macrosteles divisus* (Uhler) = *M. fascifrons* (Stål)
- p. 80, line 15: *Crambus* sp. = Place in Crambidae
- p. 81, line 11: *canithorax* = *mississippiensis* Hoffman
- p. 83, line 8: (Fall) = (Fall.)
- p. 83, line 39: *Dorilas* sp. = *Pipunculus*
- p. 83, line 46: Genus? = belongs under *Rhagionidae*
- p. 87, line 7: *serialis* = *seriata*
- p. 87, line 10: *Sylvanus* = *Silvanus*
- p. 88, line 26: *Photuris pennsylvanica* (DeGeer) = probably *P. versicolor* (Fab.) per McDermott

- p. 90, line 22: *Dorymymex* = *Dorymyrmex*
- p. 91, line 15: Delete and use *Linyphiidae* - Linyphiids
- p. 92, line 5: Urochorda = Urochordata
- p. 93, line 10: *Carnax* = *Caranx*
- p. 93, line 26: *Micropogen* = *Micropogon*
- p. 94, line 14: *albiguttus* = *albigutta*
- p. 94, line 15: *lethostigmus* = *lethostigma*
- p. 96, line 7: Gannet = *Sulidae*. Insert on p. 95 after *Pelecanidae*
- p. 96, line 43: Black bellied = hyphenate
- p. 97, line 35: Kingbord = Kingbird
- p. 98, line 5: *Hylochichla* = *Hylocichla*
- p. 99, line 23: *Mus* = *Rattus*, Author of species = (Berkenhout)
- p. 100, line 9: Bitter Beachgrass
- Fig. 26, line 2: groundsell = groundsel
- p. 101, line 31: *carymbosa* = *corymbosa*
- p. 102, line 35: Rafinewque's = Rafinesque's
- p. 103, line 12: *stellaria* = *stellaris*
- p. 104, line 2: Thistle = *Actinospermum*
- p. 104, line 4: Groundsell = Groundsel
- p. 104, line 10: recurring = recurving
- p. 105, line 20: 1946 = 1956