6-1979

Leeches of Some Fishes of the Mobile Bay Region

Ernest H. Williams Jr.
University of Puerto Rico

Follow this and additional works at: <https://aquila.usm.edu/goms>

DOI: 10.18785/negs.0301.04

Recommended Citation

Williams, E. H. Jr. 1979. Leeches of Some Fishes of the Mobile Bay Region. *Northeast Gulf Science* 3 (1). Retrieved from <https://aquila.usm.edu/goms/vol3/iss1/4>

This Article is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in Gulf of Mexico Science by an authorized editor of The Aquila Digital Community. For more information, please contact Joshua.Cromwell@usm.edu.

- Gobionellus* (family Gobiidae). Bull. Bingham Oceanogr. Coll., 4(2):1-51.
- . 1953. Ten new American gobiid fishes in the United States National Museum, including additions to a revision of *Gobionellus*. J. Washington Acad. Sci., 43(1):18-26.
- Girard, C. 1858. Notes upon various new genera and new species of fishes, in the museum of the Smithsonian Institution, and collected in connection with the United States and Mexican Boundary Survey: Major William Emory, Commissioner. Proc. Acad. Nat. Sci. Phila., 10(12):167-171.
- Hastings, P.A. 1979. First North American continental record of *Gobionellus pseudofasciatus* (Pisces: Gobiidae). Northeast Gulf Sci., 2(2):140-144.
- Hoese, D. F. 1977. *Signigobius biocellatus*, a new species of sand-dwelling coral reef gobiid fish from the western tropical Pacific. Japanese J. Ich., 23(4):199-207.
- Hoese, H. D. and R. H. Moore. 1977. Fishes of the Gulf of Mexico (Texas, Louisiana, and adjacent waters). Texas A and M Press, xv + 327 p.
- Hubbs, C. L. and K. F. Lagler. 1958. Fishes of the Great Lakes region. Bull. Cranbrook Inst. Sci., 26:x-xi, 1-213. (Revised edition).
- Jordan, D. S. 1919. The genera of Fishes. Pt. II. From Agassiz to Bleeker, 1833-1858. Stanford Univ. Publ., Univ. Ser., 36:155-284.
- Koumans, F. P. 1953. Gobioidae. In: The Fishes of the Indo-Australian Archipelago, by M. Weber and L. F. DeBeaufort. Vol. 10. 423 p.
- Mead, G. W. and J. E. Böhlke. 1958. *Gobionellus stigmalophius*, a new goby from the Gulf of Campeche and the Great Bahama Bank. Copeia, 1958 (4):285-289.
- Meek, S. E. and S. F. Hildebrand. 1928. The marine fishes of Panama. Pt. 3. Field Mus. Nat. Hist., Zool. Ser., 15: xxiii-xxx, 709-1045, pls. 72-102.
- Menon, A. G. K. and N. Govindan. 1977. *Oxyurichthys nijsseni*, a new gobiid fish from Ennore estuary, east coast of India, with a key to the identification of the Indo-West Pacific species of the genus *Oxyurichthys*. Matsya, 2:13-15.
- Nolan, E. J. (ed.). 1913. An index to the scientific contents of the Journal and Proceedings of the Academy of Natural Sciences of Philadelphia. Acad. Nat. Sci. Phila.: v-xiv, 1-1419.
- Pfaff, J. R. 1933. Report on the fishes collected by Mr. Harry Madsen during Professor O. Olufsen's expedition to French Sudan in the years 1927-28. Vidensk. Medd. Naturh. Foren. Kbenhavn, 94:273-315, figs. 1-13.
- Poey, F. 1868. Synopsis piscium Cubensium. Repertoire fisico natural de la isla de Cuba (1866-1868), 2:279-484.
- . 1876. Enumeratio piscium Cubensium. An. Soc. Esp. Hist. Nat., 5:131-218.
- Robins, C. R. and E. A. Lachner. 1966. The status of *Ctenogobius* Gill (Pisces: Gobiidae). Copeia, 1966(4):867-869.
- Vergara R., R. 1978. Nuevo gobido, genero *Gobionellus* (Pisces: Gobiidae), de las aguas costeras de Cuba. Poeyana, 187:1-7.

SHORT PAPERS AND NOTES

LEECHES OF SOME FISHES OF THE MOBILE BAY REGION

Brackish and marine fishes (2221 specimens of 151 species) from the Mobile Bay Region were examined for parasites from March 1969 to August 1973. Fishes were collected using

seines, monofilament gill nets and trammel nets, hook and line, gigs, dip nets, rotenone, trawls, speargun, crab traps, and lift nets. They were either examined shortly after capture or held alive and examined within a day. Leeches were flattened between glass slides and preserved in 5% formalin. Williams and Gaines (1974) listed the species, numbers, and size ranges of the hosts examined. Acanthocephala from these hosts were reported by Williams and Gaines (1974) and isopods by Williams and Williams (1978). Four species of leeches were collected, representing five new host records (Table 1).

Sawyer, Lawler, and Overstreet (1975) noted a normal infestation of one to two adults (or numerous juveniles) per host for *Branchellion ravenelii* (Giard, 1850). The five specimens collected from the lesser electric ray, *Narcine brasiliensis* (Olfers) in the present study were all adults. The new host record (Gulf kill-fish, *Fundulus grandis* Baird and Girard) for *Malmiana philoherma* Sawyer, et. al, 1975 adds a new host family, Cyprinodontidae, to the four previously known host families (Ariidae, Carangidae, Exocoetidae, Sciaenidae). *Malmiana philoherma* seems to occur very rarely on

juvenile pompano from the south beaches of Dauphin Island. Only two specimens were recorded from thousands of pompano seined for mariculture experiments from 1969 to 1972. The body and fins of all these pompano were routinely examined for external parasites. The four leeches from the red snapper, *Lutjanus campechanus* (Poey), were probably *Trachelobdella lubrica* (Grube, 1840). Unfortunately, they were lost in the mail before this identification could be confirmed.

ACKNOWLEDGMENTS

Thanks are extended to the following for help in collecting and loan of facilities and equipment: Drs. Wilmer A. Rogers and Ronald P. Phelps of the Southeastern Cooperative Fish Disease Project, Department of Fisheries and Allied Aquaculture, Auburn University; Dr. John S. Ramsey of the Cooperative Fisheries Unit of Auburn University; Dr. Joseph R. Sullivan of Humboldt State University; Hugh A. Swingle, Wayne E. Swingle, Donald G. Bland, Madison R. Powell, and Walter M. Tatum of the Alabama Marine Resources Laboratory; and Joe Addison

TABLE 1. Leeches collected from some fishes of the Mobile Bay Region.

Parasite			Host					
Species	Number present	Position	Species	Infested		Examined		Collection locality (Date)
				N	TL (cm)	N	TL (cm)	
<i>Branchellion ravenelii</i>	5	ventral skin	<i>Narcine brasiliensis</i> (Olfers)	1	76.3	2	31.3-76.3	Mouth of Perdido Bay (7 October 1972)
<i>Malmiana philoherma</i>	2	skin	<i>Trachinotus carolinus</i> (Linnaeus)	2	6.8-8.3	300	3.2-29.2	South beaches, Dauphin Island (29 July 1969)
	1	skin	<i>Fundulus grandis</i> * Baird and Girard	1	8.8	12	4.3-9.0	South beaches, Dauphin Island (10 June 1970)
<i>Myzobdella lugubris</i>	3	gills	<i>Acipenser oxyrinchus</i> * Mitchell	1	76.0	8	49.0-104.0	Fairhope, Mobile Bay (31 January 1971)
	1	gills	<i>Dorosoma cepedianum</i> * (Lesueur)	1	25.5	6	22.8-28.1	Fairhope, Mobile Bay (31 January 1971)
	1	gills	<i>Dorosoma petenense</i> * (Günther)	1	17.6	10	4.9-20.2	Fairhope, Mobile Bay (31 January 1971)
	1	gills	<i>Ictalurus furcatus</i> * (Lesueur)	1	6.4	1	6.4	The Devil's Channel, Mobile Causeway (9 April 1970)
	3	fins & barbel bases	<i>Ictalurus punctatus</i> (Rafinesque)	1	24.0	1	24.0	Mobile Causeway, Blakeley River (4 December 1969)
	1	branchial cavity	<i>Leiostomus xanthurus</i> Lacepede	1	8.5	70	7.9-25.2	Mississippi Sound North of Dauphin Island (15 July 1969)
	1	skin	<i>Micropogonias undulatus</i> (Linnaeus)	1	7.5	102	6.6-43.7	Mississippi Sound North of Dauphin Island (15 July 1969)
<i>Trachelobdella lubrica</i> (?)	1	skin	<i>Mugil cephalus</i> Linnaeus	1	49.0	186	9.7-59.2	Eastern and Dauphin Island (21 August 1969)
	1	right pectoral fin	<i>Mugil cephalus</i> Linnaeus	1	51.3	186	9.7-59.2	Mobile Causeway, Blakeley River (4 December 1969)
	1	right pectoral fin	<i>Paralichthys lethostigma</i> Jordan and Gilbert	1	25.4	19	22.4-35.5	Mobile Causeway, Blakeley River (4 December 1969)
	4	gills	<i>Lutjanus campechanus</i> (Poev)	1	28.0	18	20.0-44.0	S.E. Snapper Banks off Dauphin Island (11 June 1970)

* New host records.

and Larry Johnson of Alabama Department of Conservation and Natural Resources. Appreciation is expressed to Dr. Eugene M. Bureson, Virginia Institute of Marine Sciences; and Lucy B. Williams, of the Department of Marine Sciences, University of Puerto Rico for reviewing the manuscript.

LITERATURE CITED

- Sawyer, R. T., A. R. Lawler, and R. M. Overstreet. 1975. Marine leeches of eastern United States and the Gulf of Mexico with a key to the species. *J. Nat. Hist.* 9: 633-677.
- Williams, E. H., Jr. and J. L. Gaines, Jr. 1974. Acanthocephala of fishes from marine and brackish waters of the Mobile Bay Region. *J. Mar. Sci., Alabama* 2 (3): 135-148.
- _____ and L. B. Williams. 1978. Cymothid isopods of some marine fishes from the northern Gulf of Mexico. *Northeast Gulf Sci.* 2 (2):122-124.
- Ernest H. Williams, Jr., *Department of Marine Sciences, University of Puerto Rico, Mayaguez, PR 00708.*

RECYCLING FRESHLY SHUCKED OYSTER SHELLS

Oyster shells are extensively used in culture operations to collect oyster spat. Those shells are either dredged (mud shells) or weathered shells from processing plants (St. Amant, 1958). The replanting of oyster shells immediately after shucking was reported by Moore (1897), but is not currently practiced along the Gulf Coast. The purpose of this study was to determine the growth and survival of attached spat on freshly shucked shells after replanting in Biloxi Bay, Mississippi.

MATERIALS AND METHODS

Freshly shucked, Louisiana oyster shells were obtained from two seafood processing plants on the Back Bay of Biloxi on two separate occasions. The shells were held in fiberglass troughs supplied with running bay water until planted five weeks later. The numbers of shells and the numbers of spat were determined for each of the four experimental groups of shells. The spat on each group of shells were measured (greatest umbo to bill dimension) and marked with an identifying number. The numbered, plastic tags were cemented to the spat with Marine Tex® epoxy resin (Travaco Laboratories, Inc., 345 Easter Ave., Chelsea, Mass. 02150). The four experimental groups were planted on the bottom of Biloxi Bay at Point Cadet just below mean low water in 1 m² staked plots during March, 1977. The four experimental plots contained 201, 525, 527, and 532 single valves with 37, 100, 100, and 41 marked spat, respectively. Two, 1m² control plots were also established and received no shells. All plots were cleared of existing shell material prior to treatment. In addition, two experimental plots were covered with four mil polyethylene sheeting prior to the planting of the shells.

RESULTS

Eighty-six of the 278 marked spat (31%; range 20 to 42%) were recovered at the end of ten months; of these 25 or 9% (range 5-24%) of the original marked spat were recovered alive (Table 1). The surviving spat had grown an average of 29.0 mm (2.9 mm/mo.),

Of the original 1,785 shells planted, 1,372 (77%; range 60 to 106%) were recovered from the experimental plots;