4-1981

First Gulf of Mexico Record for *Biancolina brassiacephala* (Amphipoda: Biancolinidae)

P. Steele

Florida Department of Natural Resources

S.B. Collard

University of West Florida

DOI: 10.18785/negs.0402.06

Follow this and additional works at: <https://aquila.usm.edu/goms>

Recommended Citation

Steele, P. and S. Collard. 1981. First Gulf of Mexico Record for *Biancolina brassiacephala* (Amphipoda: Biancolinidae). *Northeast Gulf Science* 4 (2).

Retrieved from <https://aquila.usm.edu/goms/vol4/iss2/6>

This Article is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in Gulf of Mexico Science by an authorized editor of The Aquila Digital Community. For more information, please contact Joshua.Cromwell@usm.edu.

FIRST GULF OF MEXICO RECORD FOR *Biancolina brassiacephala* (AMPHIPODA: BIANCOLINIDAE)

Six specimens of the monotypic family of marine amphipods, Biancolinidae Barnard, 1972, were found attached to rafts of *Sargassum natans* and *S. fluitans* collected in surface neuston tows in the northeastern Gulf of Mexico. Single specimens of *Biancolina brassiacephala* Lowry, 1974 (Figs. 1-7) were collected in each of three tows at Station C near the Florida Middle Ground, 28°36'N, 84°15' W, in June-July 1976, over 27 m of water, and three others were collected in three tows at Station D southwest of Tampa Bay, 27°24'N, 84°07'W, over 96 m of water, in August 1976 (Fig. 8). All specimens were nongravid females. Several other species of amphipods were found with *B. brassiacephala* on *Sargassum* rafts, including *Lestrigonus bengalensis*, *Hemigina minuta*, and *Phoxocephalus* spp.

Biancolina algicola was first established by Della Valle (1893) for specimens collected on algae in the Bay of Naples, Italy, and was later included in the family Ampithoidae by Stebbing (1906). Nichols (1939) expanded the genus to include *B. australis* from Rottnest, Western Australia, and placed it in the Prophliantidae. Ruffo and Wieser (1951) redescribed *B. algicola* from specimens collected from the Mediterranean. Gurjanova (1958) included *Biancolina* in a newly erected family, Eophilantidae. Barnard (1970) added the species *B. mauihina*, found on intertidal *Sargassum* at Oahu, Hawaii. The possession of non-talitroidean characteristics, e.g., rough epidermis and biramous uropod 3, prompted Barnard (1972) to remove *Biancolina* from the Eophilantidae (superfamily Talitroidea) and place it in the newly erected, monotypic family Biancolinidae. Lowry (1974) described *Biancolina brassiacephala* from specimens associated with rafts of *Sargassum*

Figures 1, 2. *Biancolina brassiacephala* female, 2.5 mm: 1. Lateral view; 2. Anteroventral view.

natans and *S. fluitans* collected from the Gulf Stream and Sargasso Sea in the western North Atlantic Ocean off North and South Carolina. Our specimens extend the known range of *B. brassiacephala* into the Gulf of Mexico.

Specimens from the eastern Gulf of Mexico were collected with a 1 m Khalsico Floating Plankton sampler (mesh size = 202 μ m modified to sample the neustonic community from the surface down to 10 cm. Towing time ranged between 17 and

Figures 3-7. *Biancolina brassiacephala* female: **3.** Gnathopod 1; **4.** Uropod 2; **5.** Maxilliped; **6.** Antenna 2; **7.** Pereopod 1.

TABLE 1. Oceanographic and meteorological collection data, Summer, 1976.

Station	Date	Latitude	Longitude	Depth (meters)	Time Set	Time Haul	Sea State	Temperature(°C)	
								Air	Surface
C-087	6/23/76	28°36'N	84°15'W	27	1800	1833	1	38.0	29.5
C-102	7/25/76	28°36'N	84°15'W	27	1202	1305	1	33.4	30.1
C-107	7/26/76	28°36'N	84°15'W	27	0205	0235	1	28.0	29.8
D-128	8/03/76	27°24'N	84°07'W	96	0600	0617	2	29.7	28.0
D-139	8/04/76	27°24'N	84°07'W	96	2000	2018	1	30.0	29.3
D-142	8/05/76	27°24'N	84°07'W	96	0200	0218	1	28.5	29.2

63 minutes at a ship speed of two knots. Physical oceanographic data and meteorological observations were made at the time of collection (Table 1).

Biancolina brassiacephala has been found only on algal rafts of *S. natans* and *S. fluitans*, suggesting a possible method for species dispersal between the north-eastern Gulf of Mexico and the northwest Sargasso Sea. Loop Current circulation over the Florida Middle Ground during the summer of 1976 has been confirmed by Steward (1980). Individuals on rafts of pelagic *Sargassum* could be transported between the northeast Gulf and the northwest Sargasso Sea by the Loop Current, Florida Current and Gulf Stream systems.

ACKNOWLEDGMENTS

Work in the northeastern Gulf of Mexico was carried out under the Bureau of Land

Management Contract No. 08550-CTS-30-SUSIO.

At the Florida Department of Natural Resources Marine Research Laboratory, we are grateful to Messrs. W. G. Lyons, D. K. Camp, F. S. Kennedy, and S. A. Willis for constructive suggestions which improved the manuscript. Special thanks to Ms. L. Tester for SEM photomicrographs and Ms. L. Walker for light photomicrographs.

LITERATURE CITED

- Barnard, J. L. 1970. Sublittoral Gammaridea (Amphipoda) of the Hawaiian Islands. Smithsonian Contrib. Zool. 34: 1-286.
- _____. 1972. The marine fauna of New Zealand: Algae-living littoral Gammaridea (Crustacea: Amphipoda). Mem. New Zealand Oceanogr. Inst. 62: 1-216.
- Della Valle, A. 1893. Gammarina del Golfo Napoli. Fauna Flora Golf. Neapel. 20: 1-948.
- Gurjanova, E. 1958. K fauna amfipod (Amphipoda) Ostrova Nakkuori. Biol. Soviet. Antarct. Exped. 3: 55-56.
- Lowry, J. K. 1974. A new species of the amphipod *Biancolina* from the Sargasso Sea. Trans. Am. Micros. Soc. 83(1): 71-78.
- Nichols, G. E. 1939. The Prophilantidae. A proposed new family of Amphipoda, with descriptions of a new genus and four new species. Rec. S. Austral. Mus. 6: 309-334.
- Ruffo, S., and W. Wieser. 1951. Osservazi-

Figure 8. Sampling stations, June-August 1976.

oni sistematiche ed ecologiche su alcuni Anfipodi delle coste mediterranee italiane. Mem. Mus. Civ. Stor. Nat. Verona 3: 11-30.

Stebbing, T. R. R. 1906. Amphipoda. 1. Gammaridea. Das Tierreich, Vol. 21, xxxix + 806 p.

Steward, R. 1980. Spatial distribution of suspended particulate matter in the northeastern Gulf of Mexico. Unpubl. Masters Thesis, University of South Florida, St. Petersburg, Florida. 100 p.

Steele, P. *Florida Department of Natural Resources, Marine Research Laboratory, 100 Eighth Avenue S. E., St. Petersburg, Florida 33701.*

Collard, S. B. *Department of Biology, University of West Florida, Pensacola, Florida 32504.*