

7-1982

## First Gulf of Mexico Record of *Zu cristatus* (Bonnelli) (Pisces: Trachypteridae)

William J. Eckmayer  
*Alabama Marine Resources Laboratory*

DOI: 10.18785/negs.0502.05

Follow this and additional works at: <https://aquila.usm.edu/goms>

---

### Recommended Citation

Eckmayer, W. J. 1982. First Gulf of Mexico Record of *Zu cristatus* (Bonnelli) (Pisces: Trachypteridae). *Northeast Gulf Science* 5 (2). Retrieved from <https://aquila.usm.edu/goms/vol5/iss2/5>

This Article is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in *Gulf of Mexico Science* by an authorized editor of The Aquila Digital Community. For more information, please contact [Joshua.Cromwell@usm.edu](mailto:Joshua.Cromwell@usm.edu).

**FIRST GULF OF MEXICO  
RECORD OF *Zu cristatus*  
(BONNELLI) (PISCES:  
TRACHYPTERIDAE)**

A scalloped ribbonfish, *Zu cristatus* (Bonnelli), (Fig. 1) was collected on March 30, 1981 by Mr. Elmer Wheeler of Coden, Alabama. Using oyster tongs he captured the fish while it was swimming on the surface at the south end of Portersville Bay between Cat Island and Barron Point (30°18.1'N, 88°12.1'W). The fish has been deposited in the Ichthyological Museum of the Gulf Coast Research Laboratory (GCRL 17184), Ocean Springs, Mississippi and represents the first collection of scalloped ribbonfish in the Gulf of Mexico.

Meristic data taken from this 316 mm SL specimen are given below.

dorsal - 6+129  
anal - absent

pectoral - 11

pelvic - 6

Fin ray counts agree with those listed by Miller and Lea (1972).


Scalloped ribbonfish are pelagic, circumtropical fish with a habitat which ranges from shallow water to a depth of 90 m (Briggs, 1960). Previous specimens have been collected from the mid-Atlantic Ocean (Backus *et al.*, 1965), Mediterranean Sea and both western and eastern Pacific Ocean (Fitch, 1964).

**ACKNOWLEDGMENTS**

I wish to thank Dr. C.E. Dawson for verification of identification and distribution of the scalloped ribbonfish and for photographing the specimen.

**LITERATURE CITED**

Backus, R.H., G.W. Mead, R.L. Haedrich and A.W. Ebeling. 1965. The mesope-



**Figure 1.** Scalloped ribbonfish, *Zu cristatus* (Bonnelli), 316 mm SL captured by Elmer Wheeler in Portersville Bay, Alabama, March 30, 1981. Photograph by C.E. Dawson.

52 Short papers and notes

logic fishes collected during cruise 17 of the *R/V Chain*, with a method for analyzing faunal transects. Bull. Mus. Comp. Zool. 134(5): 139-158.

Briggs, J.C. 1960. Fishes of worldwide (circumtropical) distribution. *Copeia* 1960(3):171-180.

Fitch, J.E. 1964. The ribbonfishes (Family Trachypteridae) of the eastern Pacific, with a description of new species. *Calif. Fish Game* 50 (4):228-240.

Miller, D.J. and R.N. Lea. 1972. Guide to the coastal marine fishes of California. *Calif. Fish Game Fish. Bull* 157:235 pp.

William J. Eckmayer, *Alabama Department of Conservation and Natural Resources Marine Resources Division, Alabama Marine Resources Laboratory, Post Office Box 189, Dauphin Island, AL 36528-0189.*