

Gulf and Caribbean Research

Volume 6 | Issue 4

January 1980

Studies on *Amyloodinium ocellatum* (Dinoflagellata) in Mississippi Sound: Natural and Experimental Hosts

Adrian R. Lawler
Gulf Coast Research Laboratory

Follow this and additional works at: <https://aquila.usm.edu/gcr>



Part of the [Marine Biology Commons](#)

Recommended Citation

Lawler, A. R. 1980. Studies on *Amyloodinium ocellatum* (Dinoflagellata) in Mississippi Sound: Natural and Experimental Hosts. *Gulf Research Reports* 6 (4): 403-413.

Retrieved from <https://aquila.usm.edu/gcr/vol6/iss4/8>

DOI: <https://doi.org/10.18785/grr.0604.08>

This Article is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in *Gulf and Caribbean Research* by an authorized editor of The Aquila Digital Community. For more information, please contact aquilastaff@usm.edu.

STUDIES ON *AMYLOODINIUM OCELLATUM* (DINOFLAGELLATA) IN MISSISSIPPI SOUND: NATURAL AND EXPERIMENTAL HOSTS¹

ADRIAN R. LAWLER

Parasitology Section, Gulf Coast Research Laboratory,
Ocean Springs, Mississippi 39564

ABSTRACT Four species of parasitic dinoflagellates have been found to occur naturally on the gills and fins of Mississippi Sound fishes: *Amyloodinium ocellatum* (Brown 1931) Brown and Hovasse 1946, *Oodinium cyprinodontum* Lawler 1967, and two undescribed species. Sixteen of 43 species of fishes examined had natural gill infections of *A. ocellatum*. Seventy-one of 79 species of fishes exposed to *A. ocellatum* dinospores were susceptible, and succumbed, to the dinoflagellate. Eight did not die even though exposed to numerous dinospores. The most common signs in an infested fish were spasmodic gasping and uncoordinated movements. Trophonts of *A. ocellatum* were found on the gills, skin, fins, eyes, pseudobranchs, membranes of the branchial cavity and around the teeth; and in the lateral line pits, nasal passages, esophagus, and intestine of experimentally infected fishes. The dinoflagellate causes extensive mortalities of fishes held under closed-system mariculture conditions.

INTRODUCTION

Parasitic dinoflagellates were first reported from vertebrates by Brown (1931), who described *Oodinium* (= *Amyloodinium*) *ocellatum* from marine aquarium fishes. Additional references to this species are those by Brown (1934), Nigrelli (1936, 1939, 1940), Brown and Hovasse (1946), Chatton (1952), Porter (1952), Dempster (1955, 1956, 1972), Laird (1956), Simkatis (1958), Højgaard (1962), Paccaud (1962), Chlupaty (1962), Buxton (1962), Graaf (1962), Valenti (1968), Kingsford (1975), Becker (1977), and Lawler (1977a, 1977b, 1979).

Oodinium limneticum, a species from freshwater fishes, was described by Jacobs (1946); additional references are those by Kozloff (1948, 1949), Patterson (1949), and Lewis (1963). Another freshwater species, *O. pillularis* Schäperclaus, 1954, has received considerable attention in Europe (Weiser 1949; Schäperclaus 1951; Hirschmann and Partsch 1953; Reichenbach-Klinke 1954, 1955, 1956, 1961; Schubert 1959; Geus, 1960a, 1960b, 1960c, 1960d; Reichenbach-Klinke and Elkan 1965; and Lucky 1970).

An estuarine species, *O. cyprinodontum*, was described from Virginia cyprinodontids by Lawler (1967). It has been reported by Dillon (1966), Lawler (1967, 1968a, 1968b), Lom and Lawler (1971, 1973), and Williams (1972). Of these species, only *O. pillularis* has not been reported from North America. Unidentified parasitic dinoflagellates from North American fishes have been reported by Overstreet (1968), Lom and Lawler (1971, 1973), and Paperna and Zwerner (1976). *Oodinium cyprinodontum* has been reported from the Gulf of Mexico (Williams 1972); Kingsford (1975) illustrated a trophont from a mangrove snapper (*Lutjanus griseus*) with an *A. ocellatum* infection, presumably from the eastern Gulf of Mexico.

Amyloodinium ocellatum (Brown 1931) causes extensive mortalities of fishes in confined areas (Brown 1931, 1934; Nigrelli 1936, 1939, 1940; Højgaard 1962; Lawler 1977a, 1977b), and has been reported from many species of marine teleosts (Brown 1934; Nigrelli 1936, 1940; Brown and Hovasse 1946; Dempster 1955, 1956; Chlupaty 1962; Graaf 1962; Paccaud 1962; Straughan 1970; Lawler 1979). *Oodinium cyprinodontum* Lawler 1967 has not yet been found to cause fish deaths in aquaria (Lawler, unpublished).

In October 1971, I started trying to control *A. ocellatum* on *Micropogonias undulatus* which were being held for experiments on lymphocystis (Cook 1972). Studies were initiated to ascertain the following: (1) identity of the species of parasitic dinoflagellates naturally present on fishes of Mississippi Sound; (2) the natural hosts of parasitic dinoflagellates in Mississippi waters; (3) those fishes susceptible to *A. ocellatum* in aquaria; and (4) the methods for controlling dinospores and trophonts of *A. ocellatum* in aquaria. The present report includes my findings thus far on the first three items. The results of efforts to control the parasites will be presented in a future report.

MATERIALS AND METHODS

Fishes were collected mainly in Mississippi Sound, from Biloxi Bay to Horn Island, by means of trawls, traps, dip nets, and hook and line. Salinity was determined with a refractometer. Fishes were held alive in water from their place of capture and examined within a day. The fishes were bled by gutting them or cutting their tail to draw the blood away from the gills, thus facilitating examination of the excised gills under reflected light with a dissecting microscope. The gills of fishes collected were examined for natural infections in autoclaved seawater to avoid introduction of fortuitous dinoflagellates. Infected filaments were removed with irridatory scissors and the live trophonts were examined with a compound microscope for the presence of a stigma, stomopode, and other morphological characters.

¹This study was conducted in part in cooperation with the U.S. Department of Commerce, NOAA, National Marine Fisheries Service, under PL 88-309, Project No. 2-262-R.

Manuscript received August 1, 1979; accepted October 10, 1979.

To ascertain the fishes susceptible to *A. ocellatum*, live individuals were placed in several 1- to 100-gallon aquaria with high concentrations of dinospores; the fishes were fed periodically. High levels of dinospores were ensured by periodically introducing susceptible fishes. When the introduced fishes died, they were either left in the tanks so the dinoflagellates could encyst and fall to the tank bottoms, or their excised gills were returned to the tanks. The date and time of introduction for each fish into experimental infection tanks were recorded so that an idea of how fast *A. ocellatum* killed each fish could be determined. However, since dinospore concentrations could not be adequately monitored in a tank, the intensity of attacks could only be estimated by the number of trophonts present per gill filament at the death of the fish. All introduced test fishes were watched closely and were examined at or near death for the presence of dinoflagellates. The date and time of each death were recorded. Trophont morphology and the presence of a stigma were confirmed for each infection.

Three quick ways to check for *A. ocellatum* without excising the gills and killing the host are to: (1) quickly lift the opercule of a fish and examine the gills under a dissecting microscope; (2) immerse and hold down a live fish in a dish of water and examine the eyes and fins; or (3) brush a live fish held in a dish of water and examine the bottom of the dish for dislodged trophonts.

All scientific names of fish hosts are according to Bailey et al. (1970) except *Micropogonias undulatus*, which has undergone a name change (Chao 1978). Measurements are given in microns.

RESULTS AND DISCUSSION

Parasitic Dinoflagellates Recovered

A list of hosts for four parasitic dinoflagellates is presented in Table 1.

Species 1: This species is presently considered to be *Amyloodinium ocellatum* (Brown 1931) Brown and Hovasse, 1946 because: (1) morphology and size of the trophont compare with that described by Brown (1934) and Brown and Hovasse (1946); (2) a red stigma is present; (3) a stomopode ("root-like process" of Brown and Hovasse, 1946; "flagellum" of Nigrelli, 1936) is present; and (4) the stigma persists throughout all stages (Brown and Hovasse 1946). Even though it can be keyed out to *A. ocellatum* using the key presented by Brown and Hovasse (1946), the majority of dinospores differed somewhat in shape and size from those described by Brown (1934), Nigrelli (1936), and Brown and Hovasse (1946). Some spores were more flattened anteriorly-posteriorly than others. Dinospore characteristics are: red stigma in hypocone; epicone smaller than hypocone; one flagellum in a very distinct girdle with an undulating beat, and the sulcal flagellum long with a more or less back and forth beat; body 11.6–15.4 long by 10.4–14.5 wide (N = 10), apparently after the last division. Dinospores free-swimming for at least 15 days. Pyriform trophonts up to 350 long, but most usually 150 or less.

Trophonts nonpigmented, containing starch granules, vacuoles, and a large nucleus, and possessing a red stigma, stomopode, and attachment rhizoids. Cysts held in seawater were allowed to divide and the following observations were made: (1) all the cells do not divide at once, (2) the stigma is retained throughout division stages as noted by Nigrelli (1936), and (3) cells in division stages are in columns as noted by Nigrelli (1936).

Species 2: Spherical to elongate with greenish pigment and reddish granules; holdfast and stomopode not observed. This is possibly a free-living dinoflagellate which adheres to the mucus of fish skin and gills.

Species 3: Larger and more robust than *A. ocellatum*; no pigment or stigma; stomopode present; broad holdfast disc with numerous rhizoids; distal, large nucleus. This is possibly the same species as reported by Overstreet (1968).

Species 4: The fourth species, *Oodinium cyprinodontum* Lawler 1967, has already been reported from the Gulf of Mexico by Williams (1972), from gills of both *Fundulus similis* (Baird and Girard) and *Cyprinodon variegatus* Lacépède collected near Pensacola, Florida, and Mobile Bay, Alabama, respectively. I recovered *O. cyprinodontum* from the same two hosts from various localities (Deer and Horn islands, Davis Bayou) in Mississippi Sound. It was also recovered from another cyprinodontid, *Adinia xenica* (Jordan and Gilbert), a new host record, from Horn Island; of five fish examined, one individual was observed.

Natural Hosts of Amyloodinium ocellatum

Of 99 fishes examined, representing 43 species from 28 families, 16 species of 13 families were found to be infected naturally with *A. ocellatum* on the gills (Table 1). This represents more natural hosts than previously reported. Natural infections generally are very light in Mississippi waters, and apparently cause no deaths. The natural prevalence and intensity of *A. ocellatum* on *Micropogonias undulatus* (Linnaeus) in Mississippi waters will be discussed by Dr. Robin M. Overstreet (in preparation). The natural intensity on fishes I examined generally was low; however, one specimen of *Cynoscion nebulosus* (Cuvier) had 115 *A. ocellatum* on its gills. This number still is far short of the several thousand necessary to kill the host. Three of the hosts from the present study previously have been reported to be susceptible to *A. ocellatum* (Table 1); the rest were reported recently as new host records by Lawler (1979). Nigrelli (1936) assumed that *Chilomycterus schoepfi* (Walbaum) was a natural host which, when introduced into the New York Aquarium closed-circulation tanks, spread the parasite to other fishes. Brown (1934) found that *Holocentrus ascensionis* (Osbeck), *Teuthis bahianus* (= *Acanthurus bahianus* Castelnau), and *Glyphisodon saxatilis* (= *Abudefduf saxatilis* [Linnaeus]?) had very slight infections of *A. ocellatum* when they arrived dead in a consignment of fish from Bermuda. Dempster (1955) found *A. ocellatum* on both the gills and body of *Amphyprion*

TABLE 1.
Mississippi Sound fishes examined for the natural occurrence of parasitic dinoflagellates.*

Species	mm Total Length (TL) ¹	Dinoflagellate species present	Number infected per number examined	Estimated number of dinoflagellates per infected host (gills)
Acipenseridae				
<i>Acipenser oxyrinchus desotoi</i> Vladykov	1,065 SL (standard length)	—	0/1	—
Lepisosteidae				
<i>Lepisosteus osseus</i> (Linnaeus)	1,029	—	0/1	—
Elopidae				
<i>Elops saurus</i> Linnaeus	267	—	0/1	—
Ophichthidae				
<i>Ophichthus gomesi</i> (Castelnau)	273, 458, <u>508</u>	1?	1/3	1
Clupeidae				
<i>Brevoortia patronus</i> Goode	232	—	0/1	—
<i>Dorosoma petenense</i> (Günther)	109	—	0/1	—
Synodontidae				
<i>Synodus foetens</i> (Linnaeus)	<u>206</u> <u>229</u>	3? 1?	1/2 1/2 (2/2)	11 few
Ariidae				
<i>Bagre marinus</i> (Mitchill)	153, <u>157</u> , 163	2?	1/3	several
Batrachoididae				
<i>Porichthys porosissimus</i> (Valenciennes)	163	—	0/1	—
Gadidae				
<i>Urophycis floridanus</i> (Bean and Dresel)	61, 95, 106, 124, 203, 207	—	0/6	—
Cyprinodontidae				
<i>Adinia xenica</i> (Jordan and Gilbert)	<u>29</u> , 29, 30, 30, 39	4	1/5	1
<i>Cyprinodon variegatus</i> Lacépède	<u>41</u> , 43 <u>37</u> <u>47</u>	3 4 1?	1/2 2/3 2/3	2 25 200
<i>Fundulus similis</i> (Baird and Girard)	<u>57</u> <u>?</u> , 39, 82	4 1?	1/3	14 150–200 5
Poeciliidae				
<i>Poecilia latipinna</i> (Lesueur)	37	—	0/1	—
Atherinidae				
<i>Menidia beryllina</i> (Cope)	79	—	0/1	—
Serranidae				
<i>Centropristis ocyurus</i> (Jordan and Evermann)	110	—	0/1	—
<i>Centropristis philadelphia</i> (Linnaeus)	115, 119	—	0/2	—
Carangidae				
<i>Selene vomer</i> (Linnaeus)	113	—	0/1	—
<i>Vomer setapinnis</i> (Mitchill)	123	—	0/1	—
Lutjanidae				
^{2,7} <i>Lutjanus griseus</i> (Linnaeus)	<u>72</u> , <u>104</u>	3?	1/2	1
² <i>Lutjanus analis</i> (Cuvier)	<u>150</u>	1?	1/1	2–3
Gerreidae				
<i>Eucinostomus argenteus</i> Baird and Girard	79, <u>89</u>	1?	1/2	20–30
Sparidae				
<i>Archosargus probatocephalus</i> (Walbaum)	305, <u>337</u>	1	1/2	30–50
<i>Lagodon rhomboides</i> (Linnaeus)	106, <u>104</u>	1	1/2	less than one/filament
Sciaenidae				
<i>Bairdiella chrysura</i> (Lacépède)	50, 65	—	0/2	

TABLE 1 – Continued

Species	mm Total Length (TL) ¹	Dinoflagellate species present	Number infected per number examined	Estimated number of dinoflagellates per infected host (gills)
<i>Cynoscion arenarius</i> Ginsburg	100, 302	–	0/2	–
<i>Cynoscion nebulosus</i> (Cuvier)	<u>170</u> , 187, 222, 223, <u>230</u> , <u>305</u>	1	3/6	1–115
³ <i>Leiostomus xanthurus</i> Lacépède	116, 129, 138, 233, 238	–	0/5	–
<i>Menticirrhus americanus</i> (Linnaeus)	167, <u>195</u> , <u>211</u> , <u>225</u>	1	3/4	1–5
<i>Micropogonias undulatus</i> (Linnaeus)	<u>310</u>	1	1/1	50–100
<i>Sciaenops ocellata</i> (Linnaeus)	296, 407	–	0/2	–
Mugilidae				
⁴ <i>Mugil cephalus</i> Linnaeus	<u>115</u> , 239	1?	1/2	1
Blenniidae				
<i>Hypsoblennius ionthas</i> (Jordan and Gilbert)	<u>47</u>	1?	1/1	1
Gobiidae				
<i>Gobionellus hastatus</i> Girard	160	–	0/1	–
<i>Gobiosoma bosci</i> (Lacépède)	21, <u>28</u> , <u>31</u>	1	2/3	1
Microdesmidae				
<i>Microdesmus longipinnis</i> (Weymouth)	129, 141, 150, 154, 202	–	0/5	–
Trichiuridae				
<i>Trichiurus lepturus</i> Linnaeus	296	–	0/1	–
Stromateidae				
<i>Peprilus alepidotus</i> (Linnaeus)	153	–	0/1	–
<i>Peprilus burti</i> Fowler	25, 73, 76, 83, 84, 86, 91	–	0/7	–
Triglidae				
<i>Prionotus rubio</i> Jordan	<u>55</u> , 85	1	1/2	1
Balistidae				
<i>Monacanthus hispidus</i> (Linnaeus)	65	–	0/1	–
Ostraciidae				
<i>Lactophrys quadricornis</i> (Linnaeus)	<u>88</u>	1?	1/1	2
Diodontidae				
^{2,3,5,6} <i>Chilomycterus schoepfi</i> (Walbaum)	<u>219</u> , 229, 257	1?	1/3	9

*Parasitic dinoflagellate species recovered (see text):

Species 1 – *Amyloodinium ocellatum* (Brown, 1931).

Species 2 – Unidentified green-pigmented form with red granules (contaminate?).

Species 3 – Unidentified form with flagellum but no stigma.

Species 4 – *Oodinium cyprinodontum* Lawler, 1967.

¹The total length of each infected fish is underlined.

Reported as a host for *Amyloodinium ocellatum* (Brown, 1931) by:

²Nigrelli (1940)

³Nigrelli (1936)

⁴Brown (1934)

⁵Brown and Hovasse (1946)

⁶Simkatis (1958)

⁷Kingsford (1975)

percula (Lacépède) that had been collected near Singapore. Simkatis (1958) found *Oodinium* (= *A. ocellatum*?) to infect *Chilomycterus schoepfi* and *Sphoeroides maculatus* (Bloch and Schneider) naturally. Porter (1952) thought that *A. ocellatum* was introduced into aquaria by newly arrived *Amphiprion ocellatum* (= *Amphiprion ocellaris* Cuvier).

Experimental Hosts of *Amyloodinium ocellatum*

Mississippi Sound hosts that are susceptible and succumb to *A. ocellatum* in aquaria are listed in Table 2. To date, 71 species of 39 families have been confirmed to die of *A. ocellatum* in aquaria. Seven had been reported previously as susceptible hosts (Table 2), the rest recently were reported

TABLE 2.
Mississippi Sound fishes experimentally infected by and succumbing to *Amyloodinium ocellatum*.

Host	mm Total Length (TL) (range)	Number Infected
Dasyatidae		
* <i>Dasyatis sabina</i> (Lesueur)	640–740	2
Ophichthidae		
* <i>Myrophis punctatus</i> Lütken	70–168	4
* <i>Ophichthus gomesi</i> (Castelnau)	432–564	2
Clupeidae		
* <i>Harengula jaguana</i> Poey	56	1
Engraulidae		
* <i>Anchoa mitchilli</i> (Valenciennes)	29– 40	5
Synodontidae		
* <i>Synodus foetens</i> (Linnaeus)	37–216	2
Ariidae		
* <i>Arius felis</i> (Linnaeus)	104–228	5
* <i>Bagre marinus</i> (Mitchill)	135	1
Batrachoididae		
* <i>Opsanus beta</i> (Goode and Bean)	25–163	6
* <i>Porichthys porosissimus</i> (Valenciennes)	110–127	2
Gobiesocidae		
* <i>Gobiesox strumosus</i> Cope	57	1
Gadidae		
* <i>Urophycis floridanus</i> (Bean and Dresel)	184–197	3
Exocoetidae		
*Flyingfish (unidentified)	18– 33	4
Cyprinodontidae		
* <i>Fundulus jenkinsi</i> (Evermann)	20– 30	several
Poeciliidae		
* <i>Gambusia affinis</i> (Baird and Girard)	20– 30	several
Syngnathidae		
* <i>Hippocampus erectus</i> Perry	47– 50	2
* <i>Syngnathus louisianae</i> Günther	128–232	6
Percichthyidae		
^{1,2} <i>Morone saxatilis</i> (Walbaum)	88–102	3
Serranidae		
* <i>Centropristis philadelphia</i> (Linnaeus)	97	1
* <i>Epinephelus niveatus</i> (Valenciennes)	164	1
* <i>Serraniculus pumilio</i> Ginsburg	61	1
* <i>Serranus subligarius</i> (Cope)	120	1
Grammistidae		
* <i>Rypticus maculatus</i> Holbrook	205	1
Centrarchidae		
* <i>Lepomis macrochirus</i> Rafinesque	125	1
Carangidae		
¹ <i>Caranx hippos</i> (Linnaeus)	32– 58	2
* <i>Caranx latus</i> Agassiz	118	1
* <i>Chloroscombrus chrysurus</i> (Linnaeus)	50– 55	3
* <i>Oligoplites saurus</i> (Bloch and Schneider)	20–110	7
* <i>Trachinotus carolinus</i> (Linnaeus)	60	1
Lutjanidae		
* <i>Lutjanus campechanus</i> (Poey)	83	1
^{2,6} <i>Lutjanus griseus</i> (Linnaeus)	29–135	7
Lobotidae		
* <i>Lobotes surinamensis</i> (Bloch)	21– 55	6
Gerreidae		
* <i>Eucinostomus argenteus</i> Baird and Girard	36– 56	3
Pomadasyidae		
* <i>Orthopristis chrysoptera</i> (Linnaeus)	187–223	2
Sparidae		
* <i>Archosargus probatocephalus</i> (Walbaum)	89–151	7

TABLE 2 - Continued

Host	mm Total Length (TL) (range)	Number Infected
<i>*Lagodon rhomboides</i> (Linnaeus)	98-137	12
Sciaenidae		
<i>*Bairdiella chrysura</i> (Lacépède)	65-185	14
<i>*Cynoscion arenarius</i> Ginsburg	71-277	6
<i>*Cynoscion nebulosus</i> (Cuvier)	160	1
<i>*Equetus acuminatus</i> (Bloch and Schneider)	70	2
<i>*Larimus fasciatus</i> Holbrook	35- 80	2
¹ <i>Leiostomus xanthurus</i> Lacépède	104-223	11
<i>*Menticirrhus americanus</i> (Linnaeus)	110	1
<i>*Micropogonias undulatus</i> (Linnaeus)	169-196	3
<i>*Sciaenops ocellata</i> (Linnaeus)	?-190	2
Ephippidae		
¹ <i>Chaetodipterus faber</i> (Broussonet)	92	1
Mugilidae		
³ <i>Mugil cephalus</i> Linnaeus	88-110	5
Blenniidae		
<i>*Chasmodes bosquianus</i> (Lacépède)	15- 69	12
<i>*Hypsoblennius hentzi</i> (Lesueur)	40- 57	2
<i>*Hypsoblennius ionthas</i> (Jordan and Gilbert)	?- 86	10
Eleotridae		
<i>*Eleotris pisonis</i> (Gmelin)	94-114	3
Gobiidae		
<i>*Bathygobius soporator</i> (Valenciennes)	85	1
<i>*Gobioides broussonneti</i> Lacépède	413	1
<i>*Gobiosoma bosci</i> (Lacépède)	30- 56	8
<i>*Gobiosoma robustum</i> Ginsburg	46	1
<i>*Microgobius gulosus</i> (Girard)	32- 53	9
Microdesmidae		
<i>*Microdesmus longipinnis</i> (Weymouth)	?-135	4
Scorpaenidae		
<i>*Scorpaena brasiliensis</i> Cuvier	76	1
Triglidae		
<i>*Prionotus roseus</i> Jordan and Evermann	223	1
<i>*Prionotus tribulus</i> Cuvier	102	1
Bothidae		
<i>*Citharichthys spilopterus</i> Günther	84- 92	3
<i>*Etropus crossotus</i> Jordan and Gilbert	95-118	3
<i>*Paralichthys lethostigma</i> Jordan and Gilbert	148-286	4
Soleidae		
<i>*Achirus lineatus</i> (Linnaeus)	34- 41	4
<i>*Trinectes maculatus</i> (Bloch and Schneider)	56-109	13
Cynoglossidae		
<i>*Symphurus plagiata</i> (Linnaeus)	39-124	11
Balistidae		
<i>*Aluterus schoepfi</i> (Walbaum)	46	1
<i>*Monacanthus hispidus</i> (Linnaeus)	76- 79	2
Ostraciidae		
<i>*Lactophrys quadricornis</i> (Linnaeus)	60- 97	2
Tetraodontidae		
<i>*Sphoeroides parvus</i> Shipp and Yerger	50- 74	7
Diodontidae		
^{1,2,4,5} <i>Chilomycterus schoepfi</i> (Walbaum)	126-143	2

Reported as a host by:

* Lawler (1979)

¹ Nigrelli (1936)² Nigrelli (1940)³ Brown (1934)⁴ Brown and Hovasse (1946)⁵ Simkatis (1958)⁶ Kingsford (1975)

as new hosts by Lawler (1979). There is no way of ascertaining if these susceptible fishes had light infections prior to their introduction into tanks containing dinospores. Fish deaths were noted as soon as 12 hours after introduction into tanks with high dinospore concentrations.

The numbers of infected fishes listed in Table 2 represent those hosts confirmed to have died from *A. ocellatum* after examination of fresh material with a compound microscope. Many more individuals of these fishes died from *A. ocellatum* in maintaining dinospore concentrations in tanks, in experimental host studies, and in treatment experiments. Because some fish died before they could be critically examined, only those confirmed to have died from *A. ocellatum* are enumerated in Table 2. Susceptible and locally abundant fishes such as *Arius felis*, *Oligoplites saurus*, *Lutjanus griseus*, *Archosargus probatocephalus*, *Lagodon rhomboides*, *Bairdiella chrysura*, *Leiostomus xanthurus*, *Micropogonias undulatus*, *Chaetodipterus faber*, *Hypsoblennius ionthas*, *Gobiosoma bosci*, *Achirus lineatus*, and *Trinectes maculatus* were employed in maintaining high dinospore concentrations. Their numerous deaths were not recorded.

Other susceptible hosts were reported by the following: Brown (1931, 1934); Schäperclaus (1935; as *Branchiophilus maris*, 1954); Nigrelli (1936, 1940); Brown and Hovasse (1946); Porter (1952); Dempster (1955, 1956); Laird (1956); Simkatis (1958); Amlacher (1961); Graaf (1962); Højgaard (1962); Paccaud (1962); Valenti (1968); Kingsford (1975); and Lawler (1979).

Ever since Brown (1934, p. 583) said that "provided conditions in the tanks are suitable it appears to attack all fish indiscriminately," most subsequent authors have stated that *A. ocellatum* appears "to be non-specific in host selection" (Sindermann 1970). This study, however, indicates that some Mississippi Sound fishes may not be susceptible to *A. ocellatum*, and others may become less susceptible with increasing size. Although definite proof has not yet been obtained, a few specimens of each of nine species of fishes were not found to die of *A. ocellatum* when exposed to dinospores (Table 3). One *Anguilla rostrata* stayed in an infection-tank from November 24, 1971 to June 21, 1972, but did not die from *A. ocellatum*, while several hundred other fishes died in the same tank in as little as 2 days. One *Fundulus grandis* lived in a tank with dinospores from November 24, 1971 to October 18, 1972; another specimen lived in a different infection tank from February 20 to October 24, 1972. Neither specimen died, although many other fishes introduced into the same tanks succumbed to the dinoflagellate. One *Menidia beryllina* (examined alive) had no *A. ocellatum* on its gills, but in the same tank, two stingrays *Dasyatis sabina* (Lesueur), exhibited massive (200 to 500 trophonts per gill filament) infections after the same 6-day exposure. The stingray represents the first record of an elasmobranch being susceptible to *A. ocellatum*. Also, *Opsanus beta* (Goode and Bean) appears to have some resistance to *A. ocellatum*.

Although deaths of *O. beta* can result from exposure to *A. ocellatum* (Table 2), several specimens were never found to harbor the dinoflagellates on their gills when they died from other causes (Table 3). Four specimens of *O. beta* survived until June 25-27, 1972, without any evidence of *A. ocellatum* on their gills, even though 15 *Arius felis* and 8 *Trinectes maculatus* died from *A. ocellatum* in the same tank on May 20. Brown (1934) also noted that some species appear more susceptible to this parasite than others.

TABLE 3.
Mississippi Sound fishes exposed to *Amyloodinium ocellatum* but not succumbing.

Species	mm TL (range)	Number exposed and examined
Anguillidae		
<i>Anguilla rostrata</i> (Lesueur)	?-570	4
Batrachoididae		
<i>Opsanus beta</i> (Goode and Bean)	?-180	5
Cyprinodontidae		
<i>Cyprinodon variegatus</i> Lacépède	30- 50	several
<i>Fundulus grandis</i> Baird and Girard	89-118	3
<i>Fundulus grandis</i> Baird and Girard	-	40
<i>Fundulus similis</i> (Baird and Girard)	62	1
Poeciliidae		
<i>Poecilia latipinna</i> (Lesueur)	35- 54	7
Atherinidae		
<i>Menidia beryllina</i> (Cope)	60- 79	7
Eleotridae		
<i>Dormitator maculatus</i> (Bloch)	82	1
<i>Dormitator maculatus</i> (Bloch)	-	25
Gobiidae		
<i>Gobionellus hastatus</i> Girard	138	1

Twenty lab-reared *Fundulus grandis* Baird and Girard (hatched March 26, 1978) were placed into each of two 5-gallon aquaria (salinity = 15 ppt) on July 10, 1978, and exposed to dinospores. *Amyloodinium ocellatum* cysts were added to the tanks 10 times between July 10 and October 3, 1978. None of the 40 fish so exposed died from *A. ocellatum*.

On October 12, 1977, 25 wild-caught sleepers, *Dormitator maculatus* (Bloch), were placed into a 75-gallon fiberglass tank (salinity = 17 ppt) where high dinospore concentrations had been maintained almost continuously since 1971. On July 26, 1979, eight of the sleepers were still alive; none of those that died were found to carry *A. ocellatum*. During this time period, approximately 310 fish of 15 species died of *A. ocellatum* in the same tank. Two of the sleepers were sacrificed during the study; no *A. ocellatum* were found on the gills of either one.

As noted above, when several species of fish are placed in an infection tank, some species will be affected and others will not. Conversely, if the species not affected is placed alone with dinospores, it may become affected. Also,

Nigrelli (1936) found that *Fundulus heteroclitus* (Linnaeus), when exposed to dinospores in a 2-gallon container, had infections of *A. ocellatum* on the second day; however, none were ever found infected in the New York Aquarium although they were in the presence of infected fishes.

Most of the host fish not affected either produce large amounts of mucus or can tolerate low oxygen levels. Possibly, heavy mucus production prevents attachment or the mucus of some species may contain a repellent.

The number of trophonts per gill filament necessary to kill a host varies with species and size of host. It also varies with size of the trophonts. Some examples are listed in Table 4.

All susceptible hosts reported prior to this study have been teleosts. Two *Dasyatis sabina* were placed in a tank with a high concentration of dinospores. Six days later they were still alive but badly stressed, being sluggish, and exhibiting an irregular spiracle beat (Table 5). Both were examined while still alive and were found to have a massive infection of 200 to 500 small to medium trophonts per gill filament; trophonts also were found in the olfactory organs. More extensive studies need to be made to ascertain all the species of fish that *A. ocellatum* will attack.

Dinospores did not attach to the skin of *Hyla cinerea cinerea* (Schneider) (= green tree frog) held partially submerged in water (salinity = 18 ppt) in a covered dish for 2 days with the free-swimming dinospores present throughout this period. This is significant because dinospores were observed swimming to the frog, bouncing around on the frog's skin, and then swimming off. I also found that a freshwater fish, *Lepomis macrochirus* Rafinesque, held in water with a salinity of 4 ppt, was susceptible to *A. ocellatum*. Trophonts occurred on the skin and gills, 46 on the first left gill arch alone indicating this was not a chance infection.

The signs of severely distressed fish, when they have been heavily infected by *A. ocellatum*, are listed by Brown (1934), Brown and Hovasse (1946), Dempster (1955, 1956), Straughan (1965, 1970), Valenti (1968), Lawler (1977a, 1977b), and others. These are: (1) having little or no interest in food, (2) rapidly gasping for air, (3) congregating near the surface, (4) scratching against objects in the tank, and (5) being sluggish unless prodded or chased by another fish. Brown (1934) was the first to note that different species may act differently when infected. She noted that clown fish either "rose to the top of the tank fluttering aimlessly, or lay at the bottom panting and incapable of co-ordinated movement" (Brown 1934, p. 603), whereas "*Julis pavo*, normally a bottom-living fish, frequently came to the top of the tanks gasping and often showed a curvature of the body," and other species became very sluggish. I have observed all of the above signs, which could also indicate one or more disorders other than infection by *A. ocellatum*. In addition, I note that any abnormal behavior of the fish may indicate an infection by *A. ocellatum*, the most common signs being spasmodic gasping, moving in an

uncoordinated fashion, and squirting water in an effort to backflush gills (Table 5).

TABLE 4.

A few examples of numbers of *Amyloodinium ocellatum* per gill filament and hours until death for various size fishes. Sizes of dinoflagellates were not measured.

Species	mm TL	Hours to death	Number of dinoflagellates per filament
<i>Arius felis</i>	131	91	10- 40
<i>Bagre marinus</i>	135	53	10- 40
<i>Bairdiella chrysur</i>	168, 175, 185	38	20- 30
<i>Bairdiella chrysur</i>	166	31	50+
<i>Caranx latus</i>	118	21	100+
<i>Chaetodipterus faber</i>	92	30	40- 50
<i>Chasmodes bosquianus</i>	62	24	5- 20+
<i>Cynoscion nebulosus</i>	160	42	20- 50
<i>Dasyatis sabina</i>	640, 740	144*	200-500+
<i>Etropus crossotus</i>	112	40	100+
<i>Lagodon rhomboides</i>	105, 131	161	50-100
<i>Leiostomus xanthurus</i>	120	39	20- 50
<i>Leiostomus xanthurus</i>	210	41	100-200
<i>Lutjanus griseus</i>	130	48	20- 30
<i>Mugil cephalus</i>	99	20	40- 50
<i>Mugil cephalus</i>	101, 101, 110	47	20- 50+
<i>Orthopristis chrysoptera</i>	223	39	300
<i>Paralichthys lethostigma</i>	148	52	20- 50
<i>Paralichthys lethostigma</i>	250	71	50-100
<i>Sphoeroides parvus</i>	50	28	75-100
<i>Sphoeroides parvus</i>	65	16	100-200+
<i>Symphurus plagiatus</i>	97	65	20- 30
<i>Trinectes maculatus</i>	56, 58, 65, 69	46	10- 40

*Moribund.

Amyloodinium ocellatum is reported to be primarily a gill parasite of marine fishes (Brown 1934, Brown and Hovasse 1946, Sindermann 1966) which also occurs on the skin. Dempster (1955, 1956) collected some clown fish (*Amphiprion percula*) that had both gills and body covered with *Oodinium* (= *A. ocellatum*?). Aquarists often refer to tegumental infections as "velvet disease." In the present

TABLE 5.
Partial behavior of various fishes when heavily infected by
Amyloodinium ocellatum.

Species	Behavior
<i>Dasyatis sabina</i>	Sluggish; irregular beat of spiracles; snout turned up so water could be pumped easier; swimming at surface with snout out of water; coughing to back-flush gills.
<i>Ophichthus gomesi</i>	Head at top of tank in corner; gasping and squirting water (back-flushing of gills); trying to get out of tank; when weaker sinking to bottom, usually on back.
<i>Anchoa mitchilli</i>	Gasping; jerky, uncoordinated movements around tank; darting to surface and then sinking vertically with tail down.
<i>Arius felis</i>	Gasping; spasmodic beat of opercules; head up and body vertical at top of tank, then sinking vertically; on back or side at bottom gasping.
<i>Archosargus probatocephalus</i> and <i>Lagodon rhomboides</i>	Gasping; spasmodic beat of opercules, with mouth almost never fully closed; scraping on objects in tank, and tank bottom.
<i>Bairdiella chrysura</i>	Gasping; spasmodic beat of opercules; sluggish.
<i>Leiostomus xanthurus</i>	Gasping; spasmodic beat of opercules; vertical at surface and squirting water (back-flushing of gills).
<i>Micropogonias undulatus</i>	Gasping; spasmodic beat of opercules; vertical at surface, head up and squirting water (back-flushing of gills); sinking vertically with tail down.
<i>Hypsoblennius ionthas</i>	Gasping; spasmodic beat of opercules, with mouth almost never fully closed; constant swimming of some specimens; trying to jump out of tank.
<i>Prionotus tribulus</i>	Constant swimming near surface.
<i>Archirus lineatus</i> and <i>Trinectes maculatus</i>	Gasping; spasmodic beat of opercules; sluggish, little or no darting when prodded; can no longer hold onto tank sides utilizing ventral surface, slide or fall to bottom.

study, *A. ocellatum* was found on the gills (filaments, arches, rakers), skin, fins, eyes, pseudobranchs, membranes of the branchial cavity and around the teeth; and in nasal passages, esophagus, and intestines of experimentally infected fishes. It was not unusual to find 200 plus trophonts per filament on experimentally infected fish. The cysts (10 to 50) found in the intestine of *Micropogonias undulatus* were not attached, so it is possible they were ingested after being dislodged from the gills. Brown (1934) found a large collection of *A. ocellatum* cysts free in the stomach of *Julis pavo*. Højgaard (1962) said it can be found in the intestines, and noted that in such a location the parasite probably cannot be controlled effectively with copper. In addition, unidentified cysts were found under the epithelium of gill filaments of three specimens of *M. undulatus* that had been treated with the drug "TetraCare Fungi Stop." Other fishes from experimental tanks had numerous small (N = 6) 8.7–20.3

by 10.2–24.8 cysts in their gills. The cysts had no apparent attachment to the gills and no flagellum, but some had a red stigma. Whether these cysts are a resting stage of *A. ocellatum*, a free-living organism, or a different parasitic organism is unknown. Similar cysts were reported by Schäperclaus (1935, 1954) for *Branchiophilus maris* (= *A. ocellatum*).

Brown (1934) listed a series of conditions which were found in association with the occurrence of *A. ocellatum* on the gills, and said (p. 603), "In cases of heavy infection the numbers of parasites would, in themselves, constitute a mechanical cause of death by obstruction of the passage of water over the gill-filaments." Duijn (1967, p. 57) summarized her findings as follows: "The parasites cause haemorrhages [sic], inflammation and necrosis in the gills, which open [the] way to secondary bacterial infections." Nigrelli (1936) detected some infected fish by a pink-tinted mucus secretion on the surface of the body and thought, "This pink color is due possibly to waste products of the parasites, . . ." No analysis of damage to the host has been conducted for the present study.

Heavily infected gills generally lacked the bright coloration of healthy oxygenated blood and appeared light pink. Various tegumental lesions associated with *A. ocellatum* and secondary infections have also been observed. Lom and Lawler (1973) found that *Amyloodinium* sp. on Cyprinodontidae did extensive damage to epithelial cells by the rhizoids pulling out surface regions into villi-like projections which may be severed off from the cell. They suggested that the stomopode might bring host cell material to the phagocytic region of *Amyloodinium* sp. and concluded that "The extensive damage done by a single trophont to many epithelial cells in which its rhizoids are embedded, as well as avid feeding on large lumps of host cell cytoplasm, explains the high pathogenicity of dinoflagellates for fish." It is possible that the "pink-tinted mucous [sic] secretion" noted by Nigrelli (1936), and the damage noted by Brown (1934) are caused by a similar process.

Mariculture Problems

The occurrence of *A. ocellatum* in closed-system mariculture ventures is a major problem. For example, the Anadromous Fishes Section of the Gulf Coast Research Laboratory lost about 300,000 juvenile (37 to 38 days old) striped bass *Morone saxatilis*, on June 25-26, 1976, because of *A. ocellatum* (see McIlwain 1976b). This was 75 to 80% of the stock on hand. The fish were held in a recirculating system (McIlwain 1976a) consisting of 1000-gallon circular fiberglass tanks containing water (salinity = 5 ppt) which was pumped from a small craft harbor into two holding ponds, and then into the tanks. Apparently the initial infective dinospores were introduced when water was pumped from the harbor into the ponds and then to the tanks. On July 12, 1972, and on July 30, 1975, I examined dying juvenile pompano which were being raised by the

Louisiana Wildlife and Fisheries Commission at Grand Terre, Louisiana; deaths were due to 10 to 30 plus *A. ocellatum* trophonts per gill filament.

Some additional reports of outbreaks of this parasite are those occurring at (1) the Claude Petet Mariculture Center, Gulf Shores, Alabama (May 25 and October 2-3, 1978), involving *Trachinotus carolinus* (Linnaeus), *Sciaenops ocellata* (Linnaeus), *Morone saxatilis* (Walbaum), and *Lutjanus campechanus* (Poey); (2) the National Marine Fisheries Service laboratory at Galveston, Texas (June 16, 1978), involving *Morone saxatilis*; and (3) the Gulf Coast Research Laboratory at Ocean Springs, Mississippi (June 25 and July 13, 1979), involving *Mugil cephalus* Linnaeus and *Aluterus schoepfi* (Walbaum) which were being held for toxicity tests.

ACKNOWLEDGMENTS

National Marine Fisheries Service and State of Mississippi grants to Dr. Robin M. Overstreet provided the support for this study; he also gave many suggestions and critically reviewed the manuscript. The following Gulf Coast Research Laboratory personnel assisted in collecting some of the hosts: Clyde Cannette, Dennis Chew, Sherman Cole, Al Glass, Dr. William Hawkins, Dr. Richard Heard, Thomas Henderson, Tim Hendrickson, Monroe Howell, Walter Langley, Tom Mattis, Dr. Thomas McIlwain, Larry Nicholson, Ronnie Palmer, Thomas VanDevender, and Ned Whatley. Some technical assistance was provided by Steven L. Shepard and W. Jeff Tolbert. Richard S. Waller identified many of the hosts. C. E. Dawson kindly identified *Lutjanus analis*.

REFERENCES CITED

- Amlacher, E. 1961. *Taschenbuch der Fischkrankheiten für Fischereibiologen, Tierärzte, Fischzuchter und Aquarianer*. Veb Gustav Fischer Verlag, Jena. 286 pp.
- Bailey, R. M., J. E. Fitch, E. S. Herald, E. A. Lachner, C. C. Lindsey, C. R. Robins & W. B. Scott. 1970. *A List of Common and Scientific Names of Fishes from the United States and Canada* (Third Edition). Am. Fish. Soc. Spec. Publ. 6. 150 pp.
- Becker, C. D. 1977. Flagellate parasites of fish. Pp. 357-416 in *Parasitic Protozoa*, Vol. 1. Academic Press, Inc., New York, San Francisco, and London.
- Brown, E. M. 1931. Note on a new species of dinoflagellate from the gills and epidermis of marine fishes. *Proc. Zool. Soc. Lond.*, Part 1, pp. 345-346.
- _____. 1934. On *Oodinium ocellatum* Brown, a parasitic dinoflagellate causing epidemic disease in marine fish. *Proc. Zool. Soc. Lond.*, Part 3, pp. 583-607.
- _____. & R. Hovasse. 1946. *Amyloodinium ocellatum* (Brown), a peridinium parasitic on marine fishes. A complementary study. *Proc. Zool. Soc. Lond.* 116(1):33-46.
- Buxton, A. G. 1962. The transportation of fish from near and far. 1^{er} Congrès International d'Aquariologie. *Bull. Inst. Océanogr. (Monaco)*. Numéro spécial 1A:153-163.
- Chao, Labbish Ning. 1978. A basis for classifying western Atlantic Sciaenidae (Teleostei: Perciformes). *NOAA Tech. Rep. NMFS Cir.* 415, 64 pp.
- Chatton, E. 1952. La classe de Dinoflagellés ou Péridiniens. Pp. 309-390 in P.-P. Grasse (ed.), *Traité de Zoologie*. t. 1, fasc. 1.
- Chlupaty, P. 1962. Krankheiten der Korallenfische und ihre Behandlung. 1^{er} Congrès International d'Aquariologie. *Bull. Inst. Océanogr. (Monaco)*. Numéro spécial 1A:81-92.
- Cook, D. W. 1972. Experimental infection studies with lymphocystis virus from Atlantic croaker. Pp. 329-335 in *Proceedings, 3rd Annual Workshop, World Mariculture Society*. St. Petersburg, Florida.
- Dempster, R. P. 1955. The use of copper sulfate as a cure for fish diseases caused by parasitic dinoflagellates of the genus *Oodinium*. *Zoologica* 40(12):133-138.
- _____. 1956. Copper sulfate as a cure for *Oodinium* disease. *Aquarium J.* 27:193-199.
- _____. 1972. A description of the use of copper sulfate as cure for gill disease in marine tropical fish tanks. *Anchor* 6(11):450-452.
- Dillon, W. A. 1966. Provisional list of parasites occurring on *Fundulus* spp. *Va. J. Sci.* 17(1):21-31.
- Duijn, C. van. 1967. *Disease of Fishes*. London Iliffe Books Ltd. 309 pp.
- Geus, A. 1960a. Ein gefährlicher Parasit an Fischen: das Geisseltierchen *Oodinium pillularis*. *Mikrokosmos* 49(9):264-269.
- _____. 1960b. Zur Entwicklung und Lebensweise von *Oodinium pillularis*. *Aquarium und Terrarien-Zeit.* 13(8):247-250.
- _____. 1960c. Zur Entwicklung und Lebensweise von *Oodinium pillularis*. *Aquarium und Terrarien-Zeit.* 13(9):280-283.
- _____. 1960d. Nachträgliche Bemerkungen zur Biologie des fischpathogenen Dinoflagellaten *Oodinium pillularis* Schäperclaus. *Aquarium und Terrarien-Zeit.* 13(10):305-306.
- Graaf, F. de. 1962. A new parasite causing epidemic infection in captive coralfishes. 1^{er} Congrès International d'Aquariologie. *Bull. Inst. Océanogr. (Monaco)*. Numéro spécial 1A:93-96.
- Hirschmann, H. & K. Partsch. 1953. Der "Colisaparasit"—ein Dinoflagellat aus der Oodiniengruppe. *Die Aquarium Terr. Z.* 6:229-234.
- Højgaard, M. 1962. Experiences made in Danmarks Akvarium concerning the treatment of *Oodinium ocellatum*. 1^{er} Congrès International d'Aquariologie. *Bull. Inst. Océanogr. (Monaco)*. Numéro spécial 1A:77-79.
- Jacobs, D. L. 1946. A new parasitic dinoflagellate from fresh-water fish. *Trans. Am. Microsc. Soc.* 65(1):1-17.
- Kingsford, E. 1975. *Treatment of Exotic Marine Fish Diseases*. Palmetto Publishing Co., St. Petersburg, Florida. 90 pp.
- Kozloff, E. N. 1948. "Velvet": A common protozoan infection of aquarium fish. *All-Pets Magazine*. May 1948, pp. 43-44.
- _____. 1949. "Velvet"—A protozoan infection of aquarium fish. *The Aquarist* 14(1):14.
- Laird, M. 1956. Aspects of fish parasitology. Pp. 46-54 in *Proceedings, 2nd Joint Symposium of the Science Society of Malaya and Malayan Mathematical Society*.
- Lawler, A. R. 1967. *Oodinium cyprinodontum* n. sp., a parasitic dinoflagellate on gills of Cyprinodontidae of Virginia. *Chesapeake Sci.* 8(1):67-68.
- _____. 1968a. New host record for the parasitic dinoflagellate *Oodinium cyprinodontum* Lawler, 1967. *Chesapeake Sci.* 9(4):263.
- _____. 1968b. Occurrence of the parasitic dinoflagellate *Oodinium cyprinodontum* Lawler, 1967 in North Carolina. *Va. J. Sci.* 19(4):240.
- _____. 1977a. Dinoflagellate (*Amyloodinium*) infestation of pompano. Pp. 257-264 in C. J. Sindermann (ed.), *Disease Diagnosis and Control in North American Marine Aquaculture. Developments in Aquaculture and Fisheries Science*. Vol. 6. Elsevier Scientific Publishing Company, Amsterdam, Oxford, and New York.
- _____. 1977b. The parasitic dinoflagellate *Amyloodinium ocellatum* in marine aquaria. *Drum and Croaker* 17(2):17-20.

- _____. 1979. North American fishes reported as hosts of *Amyloodinium ocellatum* (Brown, 1931). *Drum and Croaker* 19(1):8-14.
- Lewis, W. M. 1963. *Maintaining Fishes for Experimental and Instructional Purposes*. Southern Illinois University Press, Carbondale. 100 pp.
- Lom, J. & A. R. Lawler. 1971. Mode of attachment and relation to host tissue in two dinoflagellates from gills of cyprinodonts of Virginia. *J. Protozool.* 18(Aug. 1971, Supplement):43-44.
- _____. 1973. An ultrastructural study on the mode of attachment in dinoflagellates invading gills of Cyprinodontidae. *Protistologica* 9(2):293-309.
- Lucky, Z. 1970. Pathological changes with oodiosis of aquarium fish. *Acta. Vet. Brno.* 39(Suppl. 1):57-66.
- McIlwain, T. D. 1976a. Closed circulating system for striped bass production. Pp. 523-534 in *Proceedings, 7th Annual Workshop, World Mariculture Society*. San Diego, California.
- _____. 1976b. Striped bass rearing and stocking program—Mississippi. Completion Report AFCS-5. National Marine Fisheries Service. 121 pp.
- Nigrelli, R. F. 1936. The morphology, cytology and life-history of *Oodinium ocellatum* Brown, a dinoflagellate parasite on marine fishes. *Zoologica* 21(3):129-164.
- _____. 1939. Health of the specimens. Pp. 41-43 in *43rd Annual Report of the New York Zoological Society*.
- _____. 1940. Mortality statistics for specimens in the New York aquarium, 1939. *Zoologica* 25(4):525-552.
- Overstreet, R. M. 1968. Parasites of the inshore lizardfish, *Synodus foetens*, from South Florida, including a description of a new genus of Cestoda. *Bull. Mar. Sci.* 18(2):444-470.
- Paccaud, A. 1962. Essais divers pour enrayer les maladies des Poissons dits "des Coraux" en eau artificielle. 1^{er} Congrès International d'Aquariologie. *Bull. Inst. Océanogr. (Monaco)*. Numéro spécial 1A:57-75.
- Paperna, I. & D. E. Zwerner. 1976. Parasites and diseases of striped bass, *Morone saxatilis* (Walbaum), from the lower Chesapeake Bay. *J. Fish Biol.* 9:267-287.
- Patterson, E. E. 1949. Velvet—A sure cure. *Oregon Aquatic News* 10(9):14.
- Porter, A. 1952. Report of the honorary parasitologist for 1951. *Proc. Zool. Soc. Lond.* 122(2):535-536.
- Reichenbach-Klinke, H. H. 1954. Untersuchungen über die bei Fischen durch Parasiten hervorgerufenen Cysten und deren Wirkung auf den Wirtskörper. *Z. Fisch. N. S.* 3:565-636.
- _____. 1955. Die Artzugehörigkeit der in Mitteleuropa vorkommenden Oodinium-Art und Beobachtungen über ihr Parasitäres Stadium (Dinoflagellata, Gymnodinidae) (English and Italian summaries). *G. Microbiol.* 1(2):106-111.
- _____. 1956. Die Dinoflagellatenart *Oodinium pillularis* Schäperclaus als Bindegewebsparasit von Süßwasserfischen. *G. Microbiol.* 1(4):263-265.
- _____. 1961. Die transkontinentale Verschleppung von Parasiten insbesondere der Wassertiere. *Vchr. Dt. Zool. Ges. Bonn* (1960). *Zool. Anz. Suppl.* 24:410-415.
- _____. & E. Elkan. 1965. *The Principal Diseases of Lower Vertebrates*. Academic Press, London and New York. 600 pp.
- Schäperclaus, W. 1935. *Fischkrankheiten*. Braunschweig, Germany.
- _____. 1951. Der Colisa-Parasit, ein neuer Krankheitserreger bei Aquarienfischen. *Aquar. u. Terrar. Ztschr.* 4(7):169-171.
- _____. 1954. *Fischkrankheiten*. 3. Auflage. Akademie-Verlag, Berlin. 708 pp.
- Schubert, G. 1959. Oodinium—nicht nur ein Hautparasit. *Datz* 1:20.
- Simkatis, H. 1958. *Salt-water Fishes for the Home Aquarium*. J. B. Lippincott Company, Philadelphia and New York. 254 pp.
- Sindermann, C. J. 1966. Diseases of marine fishes. *Adv. Mar. Biol.* 4:1-89.
- _____. 1970. *Principal Diseases of Marine Fish and Shellfish*. Academic Press, London and New York. 369 pp.
- Straughan, R. P. L. 1965. Epidemic disease in marine aquariums. *Salt Water Aquarium* 1(3):42-45.
- _____. 1970. *The Salt-water Aquarium in the Home*. A. S. Barnes and Company, South Brunswick and New York. 360 pp.
- Valenti, R. J. 1968. *The Salt Water Aquarium Manual*. Aquarium Stock Company, New York. 162 pp.
- Weiser, J. 1949. Parasites of fresh-water fish. 2. *Vestnik csl. Zool. spolecnosti* 13(3):364-371.
- Williams, E. H., Jr. 1972. *Oodinium cyprinodontum* Lawler (Dinoflagellida) on *Fundulus similis* (Baird and Girard) and *Cyprinodon variegatus* Lacépède from the Gulf of Mexico. *Ala. Mar. Resour. Bull.* 8:32-33.