7-1984

First Record of *Lophiodes* from the United States Atlantic Coast

Steve W. Ross

North Carolina Division of Marine Fisheries

John H. Caruso

Audubon Park and Zoological Garden

Follow this and additional works at: <https://aquila.usm.edu/goms>

DOI: 10.18785/negs.0701.09

Recommended Citation

Ross, S. W. and J. H. Caruso. 1984. First Record of *Lophiodes* from the United States Atlantic Coast. *Northeast Gulf Science* 7 (1).

Retrieved from <https://aquila.usm.edu/goms/vol7/iss1/9>

This Article is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in Gulf of Mexico Science by an authorized editor of The Aquila Digital Community. For more information, please contact Joshua.Cromwell@usm.edu.

Short Papers and Notes:

FIRST RECORD OF *LOPHIODES* FROM THE UNITED STATES ATLANTIC COAST

In his revision of the lophiid anglerfish genus *Lophiodes*, Caruso (1981) indicated that the three western Atlantic species (*L. beroe*, *L. monodi*, and *L. reticulatus*) are known from the Gulf of Mexico, Caribbean Sea, and Atlantic coast of northern South America. None of the 132 Atlantic specimens examined for that study was collected off the Atlantic coast of the United States. All lophiids examined from that area (30

specimens) were either *Lophius americanus* or *Lophius gastrophysus* (Caruso, 1983).

Recent survey cruises by the National Marine Fisheries Service vessels ALBATROSS IV and DELAWARE II off North Carolina have yielded three specimens of the reticulate goosefish, *Lophiodes reticulatus*: MCZ 59108, ♀, 165mm SL; ALBATROSS IV Cruise 82-11, Station 23; 35° 07'N, 75° 07'W, 101-256m; 17 Sept. 1982. UF 36471, 2♂, 108mm SL; ♀, 134mm SL; DELAWARE II Cruise 82-04, Station 154; 33° 26'N, 76° 59'W, 135m; 9 July 1982 (Fig. 1). The latter two were collected with three juvenile *Lophius gastrophysus* UF 36472,

Figure 1. *Lophiodes reticulatus* Caruso and Suttkus, 1979. Male (right) 108mm SL, female (left) 134mm SL. UF 36471; DELAWARE II Cruise 82-04, Station 154; 33°26'N, 76°59'W, 135 m; 9 July 1982. Note sexual dimorphism of olfactory organs.

74-87mm SL) whose distribution was discussed by Caruso (1983).

These collections represent the first records of *Lophiodes* on the United States east coast. Previously, the closest record of *L. reticulatus* was near the Straits of Florida (FMNH 77264, SILVER BAY Sta. 2390, Caruso and Suttkus, 1979). Of the 30 Atlantic specimens mentioned above, only six were taken south of Cape Hatteras. Apparently little effort has gone into collecting lophiid specimens from this area, and there has been some confusion concerning their identification. We suspect, therefore, that *Lophiodes reticulatus* may be widely distributed south of Cape Hatteras between 50 and 400m.

We thank the following National Marine Fisheries Service personnel for cruise support: T. Azarovitz, D. Flescher, and E. J. Gutherz. S. P. Epperly, North Carolina Division of Marine Fisheries, helped collect the specimens.

LITERATURE CITED

- Caruso, J. H. 1981. The systematics and distribution of the lophiid anglerfishes: I. A revision of the genus *Lophiodes* with the description of two new species. *Copeia* 1981(3): 522-549.
- _____. 1983. The systematics and distribution of the lophiid anglerfishes: II. Revisions of the genera *Lophiomus* and *Lophius*. *Copeia* 1983(1):11-30.
- _____. and R. D. Suttkus. 1979. A new species of lophiid anglerfish from the Western North Atlantic. *Bull. Mar. Sci.* 29(4):491-496.
- Steve W. Ross *North Carolina Division of Marine Fisheries, P.O. Box 769, Morehead City, North Carolina 28557.*
- John H. Caruso, *Audubon Park and Zoological Garden, P.O. Box 4327, New Orleans, LA 70178.*