

Gulf and Caribbean Research

Volume 8 | Issue 1

January 1985

Yield-Per-Recruit of Spotted Seatrout

Richard E. Condrey
Louisiana State University

Gerald Adkins
Louisiana Department of Wildlife and Fisheries

Michael W. Wascom
Louisiana State University

Follow this and additional works at: <https://aquila.usm.edu/gcr>



Part of the [Marine Biology Commons](#)

Recommended Citation

Condrey, R. E., G. Adkins and M. W. Wascom. 1985. Yield-Per-Recruit of Spotted Seatrout. *Gulf Research Reports* 8 (1): 63-67.
Retrieved from <https://aquila.usm.edu/gcr/vol8/iss1/9>
DOI: <https://doi.org/10.18785/grr.0801.09>

This Short Communication is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in *Gulf and Caribbean Research* by an authorized editor of The Aquila Digital Community. For more information, please contact aquilastaff@usm.edu.

SHORT COMMUNICATIONS

YIELD-PER-RECRUIT OF SPOTTED SEATROUT¹

RICHARD E. CONDREY², GERALD ADKINS³
AND MICHAEL W. WASCOM²

²Coastal Fisheries Institute, Center for Wetland Resources, Louisiana State University, Baton Rouge, Louisiana 70803-7503

³Louisiana Department of Wildlife and Fisheries, Bourg, Louisiana 70343

ABSTRACT A von Bertalanffy growth curve,

$$L = 65.47 \text{ cm} (1 - e^{-.2005(t + .4113 \text{ yr})}),$$

is derived from published data on spotted seatrout in the U.S. Gulf of Mexico and used in constructing a yield-per-recruit contour. Maximum yield-per-recruit is approached as F increases above 1 and age of first entry approaches 3.9 years (14.9 in., 1.1 lb). A linear regression is derived relating average size of capture to gill net mesh size (MS in inches),

$$L = 1.97 \text{ in.} + 8.63 \text{ MS},$$

and used along with legal sizes of first harvest to evaluate the impact of current laws in the Gulf states on yield-per-recruit of spotted seatrout.

INTRODUCTION

Spotted seatrout are one of the most important edible finfish in the northern Gulf of Mexico. Despite their pre-eminence, there is a perception that scientific "information [on seatrout] is general and, for the most part, inadequate" to meet management's needs (Lorio and Perret 1980). Current regulations on the size at harvest are not based upon a quantitative consideration of yield-per-recruit and spawner-recruit relationships. Rather, current laws are largely based upon expedient compromises between conflicting user groups (Perret et al. 1980; Merriner 1980).

In this note we present a yield-per-recruit analysis based entirely upon a synthesis of published data. While lacking the rigor of a study based upon its own data collection, this synthesis does offer a needed first look at the effect of current regulatory restrictions on the yield of this increasingly important resource.

RESULTS AND DISCUSSION

Construction of the yield-per-recruit contour

A von Bertalanffy growth curve,

$$L = 65.47 \text{ cm} (1 - e^{-.2005(t + .4113 \text{ yr})}),$$

¹This is contribution No. CFI-84-5, Coastal Fisheries Institute, Center for Wetland Resources, Louisiana State University, Baton Rouge, Louisiana 70803-7503. It results from research supported by the Louisiana Fisheries Initiative.

Manuscript received October 23, 1984; accepted May 18, 1985.

was fitted to size-at-age data (total length in cm) reported for seatrout in U.S. Gulf of Mexico estuaries (Pearson 1929, Klima and Tabb 1959, Moffett 1961, Stewart 1961, Tatum 1980, and Colura et al. 1984) (Figure 1). Mean annual air




Figure 1. Growth rate of spotted seatrout in the northern Gulf of Mexico. Tatum's "Age I+ . . . Age VI+" data are plotted as age 2 through age 7 fish under the assumption that all of the annual growth had occurred when the length-age measurements were made.

TABLE 2

Current size and gill net restrictions on the harvest of spotted seatrout in the northern Gulf of Mexico.

	Florida	Alabama	Mississippi	Louisiana	Texas
Size limit					
Recreational	12 in. (but no size limit in Gulf and Franklin counties).	12 in.	None	None	14 in.
Commercial	12 in. (but no size limit in Gulf and Franklin counties).	Currently prohibited. Formerly 12 in.	12 in.	12 in.	Currently prohibited. Formerly 12 in.
Gill net mesh size (minimum)	Varies by local statutes or general statutes of local application or by rules of the Marine Fisheries Commission that are approved by the Governor and Cabinet.	Currently prohibited. Formerly 1.25 in. in Mobile County and 1.5 in. in Baldwin County.	1.5 in.	1.75 in.	Currently prohibited. Formerly 1.5 in.




Figure 3. A yield-per-recruit contour for spotted seatrout in the northern Gulf of Mexico. Points indicate the entry levels associated with the current or recent Gulf state laws on minimum legal sizes of harvest and of gill net mesh (Table 2).

Effect on yield-per-recruit

A yield-per-recruit contour was computed with Ricker's (1975) expanded form of Beverton's expression using these estimates, Harrington et al.'s (1979) length-weight relationship, and 12 years as an estimate of the maximum attainable age (Figure 3). Sizes of first entry as denoted by legal size limits (Table 2) and average size at entry predicted for gill net mesh limits are denoted for the respective states on the plot.

The fisheries of most concern are in Florida's Gulf and Franklin counties and in Louisiana and Mississippi's recreational fisheries since these fisheries have no legal minimum limits on the size of first harvest. As such any growth-overfishing concerns are superseded by the open nature of these fisheries since they are fully exposed to the potential for spawner-recruit overfishing.

The situation in Louisiana's commercial harvest has been greatly improved by two pieces of recent legislation (Ford 1984). The first reduced Louisiana's gill net mesh from 2.0 in. to 1.75 in., moving the gill net fishery from fish averaging 19.2 in. (6.4 yr, 2.4 lb) to those averaging 17.1 in. (5.0 yr, 1.7 lb). The second increased the minimum legal commercial harvest from 10 in. (2.0 yr, .33 lb) to 12 in. (2.7 yr, .57 lb).

On the other hand, Alabama and, perhaps, Texas have recently moved away from maximum yield per recruit. In both states commercial harvest has been recently prohibited. Before the prohibition the existing regulations targeted the

commercial harvest towards the size of fish which would maximize yield: 3–4 years old, 12–15 in., and 0.6–1.1 lb. Given our current estimate of fishing mortality for Alabama, this prohibition will reduce the overall yield for that state, unless it stimulates an increase in the recreational fishery. A similar pattern might be expected for Texas, although the situation is less clear as we have no direct estimate of fishing mortality for that state.

Since the spotted seatrout fishery has a large recreational component, management may be far more concerned with catch-per-angler-hour and spawner-recruit relationships than with yield-per-recruit. Our analysis suggests, however, that efforts to optimize catch-per-angler-hour and to maintain an adequate spawning biomass may be compatible with efforts to maximize yield-per-recruit. Yield appears to be maximized when spotted seatrout are harvested at 3.9 years. This age represents the second year of spawning activity. As such, management that provides for maximum yield-per-recruit, also reduces the danger of spawner-recruit overfishing (as compared to most current regulations), and enhances the recreational experience through the harvest of larger fish.

ACKNOWLEDGMENTS

We would like to thank the following reviewers for their suggestions: A. M. Bankston, V. Guillory, G. C. Matlock, M. D. Murphy, J. E. Roussel, B. Thompson, G. Waguespack and W. Wiseman.

REFERENCES CITED

- Adkins, G., J. Tarver, P. Bowman, & B. Savoie. 1979. A study of the commercial finfish in coastal Louisiana. *Louisiana Dep. of Wildlife and Fisheries Tech. Bull.* 29. New Orleans, Louisiana.
- Adkins, G. & M. J. Bourgeois. 1982. An evaluation of gill nets of various mesh sizes. *Louisiana Dep. of Wildlife and Fisheries Tech. Bull.* 36. New Orleans, Louisiana.
- Arnoldi, D. 1982. Certain aspects of the life history and habits of the spotted seatrout in Calcasieu Lake, Louisiana. Final report for Dingle Johnson Project Number F-32. Louisiana Dep. of Wildlife and Fisheries, Baton Rouge, Louisiana.
- Colura, R. L., C. W. Porter, & A. F. Maciorowski. 1984. Preliminary evaluation of the scale method for describing age and growth of spotted seatrout (*Cynoscion nebulosus*) in the Matagorda Bay System, Texas. *Tex. Parks Wildl. Dep., Manage. Data Ser. No.* 257. Austin, Texas.
- Ford, T. B. 1984. *Governor's Task Force on Saltwater Finfish Management: Report to the Governor*. Louisiana Dep. of Wildlife and Fisheries, Baton Rouge, Louisiana.
- Harrington, R. A., G. C. Matlock, & J. E. Weaver. 1979. Standard-total length, total length-whole weight, and dressed weight-whole weight relationships for selected species from Texas bays. *Tex. Parks Wildl. Dep., Tech. Ser. No.* 26. Austin, Texas.
- Klima, E. F., & D. C. Tabb. 1959. A contribution to the biology of the spotted weakfish, *Cynoscion nebulosus* (Cuvier), from northwest Florida, with a description of the fishery. *Fla. Board Conserv. Mar. Res. Lab., Tech. Ser.* 30. St. Petersburg, Florida.
- Lorio, W. J. & W. S. Perret. 1980. Biology and ecology of spotted seatrout (*Cynoscion nebulosus* Cuvier). Pages 7–13 in: *Proceedings of the colloquium on the biology and management of red drum and spotted seatrout*. Gulf States Marine Fisheries Commission, Ocean Springs, Mississippi.
- Lorio, W. J., T. Heaton, & O. Dakin. 1980. The relative impact of netting and sport fishing on economically important estuarine species. Mississippi-Alabama Sea Grant Consortium Final Report, MASGP-79-025, Ocean Springs, Mississippi.
- Matlock, G. C., J. E. Weaver, & A. W. Green. 1978. *Trends in finfish abundance in Texas estuaries as indicated by gill nets*. Texas Parks and Wildlife Dep., Coastal Fisheries Branch, Austin, Texas.
- Merriner, J. V. 1980. History and management of the spotted seatrout fishery. Pages 55–61 in: *Proceedings of the colloquium on the biology and management of red drum and spotted seatrout*. Gulf States Marine Fisheries Commission, Ocean Springs, Mississippi.
- Moffett, A. W. 1961. Movement and growth of spotted seatrout, *Cynoscion nebulosus* (Cuvier) in West Florida. *Fla. Board Conserv. Mar. Res. Lab., Tech. Ser.* 36. 35 pp.
- NOAA. 1981. *Climatological data: Annual summary, Florida 1980*. 84(13):2–3.
- . 1983a. *Climatological data: Annual summary, Alabama 1982*. 88(13):11.
- . 1983b. *Climatological data: Annual summary, Texas 1982*. 87(13):31–32.
- Pauly, G. 1979. On the interrelationship between natural mortality, growth parameters and mean environmental temperature in 175 fish stocks. *J. Cons. Int. l'Explor. Mer* 39:175–192.

- Pearson, J. C. 1929. Natural history and conservation of the redfish and other commercial sciaenids on the Texas Coast. *Bull. U.S. Bur. Fish.* 44:129-214.
- Perret, W. S., J. E. Weaver, R. O. Williams, P. L. Johansen, T. D. McIlwain, R. C. Raulerson, & W. M. Tatum. 1980. *Fishery profiles of red drum and spotted seatrout*. Gulf States Marine Fisheries Commission, Ocean Springs, Mississippi.
- Ricker, W. E. 1975. Computation and interpretation of biological statistics of fish populations. *Bull. Fish. Res. Board Can.* 191: 251-259.
- Stewart, K. W. 1961. Contributions to the biology of the spotted seatrout (*Cynoscion nebulosus*) in the Everglades National Park, Florida. Masters thesis, Univ. of Miami, Miami, Florida.
- Tatum, W. M. 1980. Spotted seatrout (*Cynoscion nebulosus*) age and growth: data from annual fishing tournaments in coastal Alabama, 1964-1977. Pages 89-92 in: *Proceedings of the colloquium on the biology and management of red drum and spotted seatrout*. Gulf States Marine Fisheries Commission, Ocean Springs, Mississippi.
- Trent, L., & P. J. Pristas. 1977. Selectivity of gill nets on estuarine and coastal fishes from St. Andrew Bay, Florida. *Fish. Bull.* 75:185-198.

ADDENDUM (in proof)

Since this paper was written, Mississippi and Florida have begun consideration of new regulations that would change the size restrictions in their states. In Mississippi it is probable that state regulations will be changed to make it illegal to sell, offer for sale, or transport for sale in or from the state of Mississippi, spotted seatrout under 14 in. In Florida it is possible that state regulations will be changed to make 14 in. the minimum size limit for spotted seatrout for both commercial and recreational fisheries. The Florida

regulation might or might not be applied statewide. If applied statewide in Florida's recreational and commercial fisheries and applied in Mississippi's commercial fisheries, the 14 in. minimum limit would target the harvest towards the size of fish that would maximize yield-per-recruit in these fisheries. On the other hand, if part of Florida remains exempt from this regulation that part, along with the recreational fisheries in Louisiana and Mississippi, will be fully exposed to the threat of spawner-recruit overfishing.