12-1990

Maximum Total Length and Age of Black Drum *Pogonias cromis* (Osteichthyes: Sciaenidae) Off Texas

Gary C. Matlock
Texas Parks and Wildlife Department

DOI: 10.18785/negs.1102.11

Follow this and additional works at: <https://aquila.usm.edu/goms>

Recommended Citation

Matlock, G. C. 1990. Maximum Total Length and Age of Black Drum *Pogonias cromis* (Osteichthyes: Sciaenidae) Off Texas. *Northeast Gulf Science* 11 (2). Retrieved from <https://aquila.usm.edu/goms/vol11/iss2/11>

This Article is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in Gulf of Mexico Science by an authorized editor of The Aquila Digital Community. For more information, please contact Joshua.Cromwell@usm.edu.

MAXIMUM TOTAL LENGTH AND AGE OF BLACK DRUM

Pogonias cromis

(Osteichthyes: Sciaenidae)

OFF TEXAS

Black drum (*Pogonias cromis*) support economically important fisheries throughout their range (Doerzbacher *et al.* 1988). Reliable stock assessments and sustainable yield predictions are required to sustain these fisheries in the face of increasing recreational and commercial demands. Yield models, such as the Beverton-Holt model (Gulland 1969), require an estimate of maximum length and age. Doerzbacher *et al.* (1988) estimated black drum maximum total length (L_{∞}) in Texas as 798 ± 42 mm by fitting a temperature compensated von Bertalanffy growth model to tag release and recapture data. However, their estimate may have been biased low because their data were skewed toward fish 300 to 700 mm free less than 200 days. Further, L_{∞} is defined mathematically as the absolute maximum attainable length (Vaughn and Kanciruk 1982). While L_{∞} is of interest, the value most often used as a proxy for L_{∞} is the average maximum length ($\bar{L}$) as described by Alverson and Carney (1975) especially for heavily exploited species. This study reports average maximum total length ($\bar{L}$) and age (t_L) that black drum reach in Texas estuarine and Gulf of Mexico waters.

MATERIALS AND METHODS

The $\bar{L}$ that black drum reach in Texas was determined from total length (TL) frequency data following Alverson and Carney's (1975) definition that only 0.5-1.0% of the catch exceeds $\bar{L}$. Length frequencies of fish caught by anglers from privately owned boats fishing in Texas bays and the adjacent Gulf of Mexico during the period 1983-1988 were used to estimate $\bar{L}$ as above. Collection

procedures are described in Osburn and Ferguson (1987). Tagged fish free at least 3 years (Green 1986) were used to determine a range for $\bar{L}$ and its corresponding t_L in the sense of t_L in the Beverton-Holt model (Gulland 1969). The $\bar{L}$ estimates were compared to an estimate made based upon length frequencies of fish caught in trammel nets in bays only during October 1976-1980 ($\bar{L}$). Collection procedures are described in Matlock (1984).

RESULTS AND DISCUSSION

The $\bar{L}$ that black drum typically reach off Texas appears to be about 1000 to 1200 mm. This conclusion is based on the following: 1) until 1988, the state angling record was a fish 1185 mm long based upon the weight reported by Texas Parks and Wildlife Department (1988) and conversion equations in Harrington *et al.* (1979); 2) the newest state angling record is a fish 1300 mm long (Texas Parks and Wildlife Department unpublished data); 3) none of the 4349 black drum caught and landed by private boat anglers in Texas bays and the Gulf of Mexico off Texas and measured by Texas Parks and Wildlife Department (TPWD) were larger than 1200 mm; 99.5% of the fish were ≤ 1020 mm (Figure 1); and 4) three tagged fish recaptured in Texas after 10 years at large reached 686-965 mm (Table 1).

The t_L that black drum typically reach in Texas is at least 13 years. Four black drum recoveries were from fish free 10.42, 10.48, 11.34, and 11.77 years (Table 1). Each fish was at least 1 year old at tagging because their lengths equalled or exceeded the estimated length (<250 mm) of 1-year old fish (Doerzbacher *et al.* 1988, Richards 1973, Simmons and Breuer 1962, Pearson 1929). The t_L estimated from four tag recoveries is about 30% less than that estimated for black drum in Louisiana (43 years) using otoliths (Beckman *et al.*

Figure 1. Total length (mm) frequency and cumulative percentage of 4349 black drum landed by private sport-boat anglers in Texas bays and the adjacent Gulf of Mexico, May 1983 - May 1988.

1988).

The q_L of black drum in Texas bays typically is at least 755 mm. During the period October 1976 - April 1980, 99.5% of the 7244 fish captured in TPWD trammel nets were smaller than 755 mm

(Figure 2). The average maximum size (q_L^*) estimated for bay fish (755 mm) agrees well with the 798 ± 42 mm L_∞ appears to be an underestimate because the estimate of q_L^* for bay fish underestimates q_L for angler-caught bay fish.

Table 1. Summary of data for black drum tagged and released in Texas bays during the 1960's and recaptured at least 3 years later. Blanks indicate no data available. (Data from Green 1986).

At Tagging			At Recapture				
Date	Location	Total length (mm)	Date	Location	Total length (mm)	Estimated age (years) ^a	Time free (years)
7-6-59	Upper Laguna Madre, West of Marker 69	265	12-21-62	Upper Laguna Madre; Marker 96	381	3.84	3.46
7-62	Aransas Bay, Turtle Bayou	210	2-5-66	Aransas Bay; Canal near Carbon Black Plant	480	3.60	3.60
12-6-62	Galveston Bay, Mud Cut	280	11-19-66	Galveston Bay; West Bay Causeway	559	4.43	3.95
7-15-66	Aransas Bay, Turtle Pen	220	7-30-69	Aransas Bay; Copano Bay upper end	470	3.11	3.04
10-14-66	Upper Laguna Madre, East of Marker 89		3-18-77	Lower Laguna Madre; Port Mansfield	965		10.42
10-14-66	Upper Laguna Madre, East of Marker 89	340	7-21-78	Upper Laguna Madre; Starvation Point, Baffin Bay		12.66	11.77
12-13-67	Upper Laguna Madre, South Bird Island	440	6-6-78	Upper Laguna Madre; Marker 115	711	12.06	10.48
12-18-67	Upper Laguna Madre, Yerbia Annis Windmill	355	4-22-79	Upper Laguna Madre; Petronilla Creek	686	12.33	11.34
7-15-68	Upper Laguna Madre, Marker 75	440	1-2-76	Upper Laguna Madre; West of Point of Rocks			7.47

^aAge was estimated by summing the time free and the estimated age at tagging using the TL at tagging and the growth equation of Doerzbacher *et al.* (1988).

Figure 2. Total length frequency and cumulative percentage of 7244 black drum collected in trammel nets in Texas bays, October 1976 - April 1980 (from Matlock 1985).

Over 99.5% of the fish landed by anglers from Texas bays and measured by TPWD were ≤ 1015 mm. Estimates for L_{∞} of black drum in Louisiana and Virginia were 1745 mm (fork length) and 1474 mm (fork length), respectively (Beckman *et al.* 1988, Richards, 1973). Their data bases included many fish >750 mm. Growth data from fish in Texas larger than 750 mm are needed to estimate better the von Bertalanffy parameters.

Table 2. Record size of black drum landed by anglers in the Gulf of Mexico and southern Atlantic Ocean.

State	Weight (kg)	Year caught	Reference
Alabama	24.2	1986	Alabama Department of Conservation and Natural Resources (1988)
Florida	42.2	1957	Roy Williams, Florida Marine Fisheries Commission, Tallahassee, Florida
Georgia	36.7	1983	Henry Ansley, Georgia Department of Natural Resources, Brunswick, Georgia
Louisiana	34.9	1975	Gerald Adkins, Louisiana Department of Wildlife and Fisheries Sciences, Baton Rouge, Louisiana
Mississippi	24.3	1984	Fred Deegan, Mississippi Department of Wildlife Conservation, Long Beach, Mississippi
South Carolina	40.4	1978	Don Hammond, South Carolina Wildlife and Marine Resources Department
Texas	36.7	1988	Texas Parks and Wildlife Department (unpublished data), Austin, Texas

Maximum sizes of black drum occupying the cold temperate waters of the Atlantic Ocean near and north of Cape Hatteras, North Carolina, appear greater than those in the warm temperate waters of the Carolinian Province south of that general area and extending into the Gulf of Mexico. In contrast to records of 24.2 to 42.2 kg from the Carolinian Province states south of North Carolina (Table 2), the record in Delaware is 51.3 kg (International Game Fish Association 1988). Fish larger than 35 kg are commonly caught in the northeast United States (Richards 1973). The estimate of L_{∞} for black drum in Virginia is 1474 mm (Richards 1973). The difference in sizes north and south of Cape Hatteras suggests that there is a zoogeographic variation in the population dynamics of black drum, as has been suggested for several other species (Matlock 1987).

LITERATURE CITED

- Alabama Department of Conservation and Natural Resources. 1988. Alabama saltwater fish records. (Unpublished table). Gulf Shores, Ala. 2 p.
- Alverson, D.L. and M.J. Carney. 1975. A graphic view of the growth and decay of population cohorts. *J. Cons. Int. Explor. Mer.* 36:133-143.
- Beckman, D.W., A.L. Stanley, J.H. Render, and C.A. Wilson. 1988. Age and growth of black drum in Louisiana Gulf

- of Mexico. Coast. Fish. Inst., Center Wetl. Res., La. State Univ., Final Report, MARFIN Program NA87WC-H-06314. 24 p.
- Doerzbacher, J.F., A.W. Green, G.C. Matlock, and H.R. Osburn. 1988. A temperature compensated von Bertalanffy growth model for tagged red drum and black drum in Texas bays. Fish. Res. 6:135-152.
- Gree, L.M. 1985. Fish tagging on the Texas coast, 1950-1975. Tex. Parks Wildl. Dep., Coast. Fish. Branch, Mngmnt. Data Ser. No. 99. 206 p.
- Gulland, J.A. 1969. Manual of methods for fish stock assessment. Part 1. Fish population analysis. FAO Mar. Fish. Sci. 4 154 p.
- Harrington, R.A., G.C. Matlock, and J.E. Weaver. 1979. Standard-total length, total length-wet weight, and dressed-whole weight relationships for selected species from Texas bays. Tex. Parks Wildl. Dep., Tech. Ser. 26. 6 p.
- International Game Fish Association. 1988. World record game fishes. ISSN 0084-2214 ISBN 0-935217-03-07. Fort Lauderdale, Florida. 320 p.
- Matlock, G.C. 1984. A basis for the development of a management plan for red drum in Texas. Ph.D. Thesis, Texas A&M Univ., College Station, Texas. 290 p.
- Matlock, G.C. 1985. Lengths of 24 salt-water fishes caught in trammel nets in Texas bays. Tex. Parks Wildl. Dep., Coast. Fish. Branch, Mngmnt. Data Ser. No. 83. 29 p.
- Matlock, G.C. 1987. Maximum total length and age of red drum off Texas. N.E. Gulf Sci. 9(1):49-52.
- Osburn, H.R. and M.O. Ferguson. 1987. Trends in finfish landings by sport-boat fishermen in Texas marine waters, May 1974 - May 1986. Tex. Parks Wildl. Dep., Coast. Fish. Branch, Mngmnt. Data Ser. 119. 464 p.
- Pearson, J.C. 1929. Natural history and conservation of the redfish and other commercial sciaenids on the Texas coast. Bull. Bur. Fish. XLIV: 129-214.
- Richards, C.E. 1973. Age, growth, and distribution of the black drum (*Pogonias cromis*) in Virginia. Trans. Am. Fish. Soc. 102:584-590.
- Simmons, E.G. and J.P. Breuer. 1962. A study of redfish, *Sclaeonops ocellata* Linnaeus, and black drum, *Pogonias cromis* Linnaeus. Pub. Inst. Mar. Sci., Univ. Tex. 8:184-211.
- Texas Parks and Wildlife Department. 1988. Texas state fish records. PWD-Leaflet-9000-5-3/88. 1 p.
- Vaughn, D.S. and P. Kanciruk. 1982. An empirical comparison of estimation procedures for the von Bertalanffy growth equation. J. Cons. Int. Explor. Mer. 40:211-219.

Gary C. Matlock, Texas Parks and Wildlife Department, 4200 Smith School Road, Austin, Texas 78744

NORTHEAST GULF SCIENCE REVIEWERS

The following scientists have generously contributed their time and expertise toward the publication of quality science. To them we are genuinely grateful.

Dr. William Anderson	Dr. Carter Gilbert	Dr. Glenn Parsons
Mr. Randy Austin	Dr. Churchhill Grimes	Dr. Jonathan R. Pennock
Dr. Steve Branstetter	Dr. Phil Hastings	Dr. Theodore W. Pietsch
Dr. Dick Bray	Dr. Robert W. Hastings	Dr. Susan Rees
Dr. Steve Bortone	Dr. Kenneth Heck	Dr. Daniel Roberts, Jr.
Dr. George Burgess	Dr. Dannie A. Hensley	Mr. Bennie Rohr
Mr. Steven Carey	Dr. Terry Henwood	Dr. Steve Ross
Dr. John H. Caruso	Dr. Dan Holliman	Dr. Anne Rudloe
Dr. Jose Castro	Dr. Jurij Homziak	Dr. William Schroeder
Dr. Mark Chittenden	Dr. Tom Hopkins	Mr. Walter Tatum
Dr. Doug Clarke	Dr. G. Dave Johnson	Dr. Bruce Thompson
Dr. Bruce Coulle	Dr. Robert Jones	Dr. Peter Thomas
Mr. Mike Dardeau	Dr. Joseph J. Kimmel	Dr. Gregory Tolley
Dr. Will Davis	Dr. Mark W. LaSalle	Dr. John Tucker
Mr. C.E. Dawson	Dr. Tom C. Lewis	Dr. William Tyler
Dr. Tom Deardorff	Dr. Robert H. McMichael	Dr. Charlie Wenner
Dr. Doug DeVries	Mr. Don Marley	Dr. Jeff Williams
Mr. John Finucane	Dr. Gary Matlock	Dr. William J. Wiseman, Jr.
Dr. Joseph Fitzpatrick	Dr. James J. Morris	Dr. Stephen Zehr
Dr. John Fitzsimmons	Dr. Dave Nelson	
Dr. Jack Fournie	Mr. Larry Ogren	