Gulf and Caribbean Research

Volume 7 | Issue 4

January 1984

A Preliminary Checklist of Epiphytic and Benthic Marine Diatoms of Louisiana

Robert S. Maples
McNeese State University

Follow this and additional works at: <https://aquila.usm.edu/gcr>

Part of the [Marine Biology Commons](#)

Recommended Citation

Maples, R. S. 1984. A Preliminary Checklist of Epiphytic and Benthic Marine Diatoms of Louisiana. *Gulf Research Reports* 7 (4): 385-388.

Retrieved from <https://aquila.usm.edu/gcr/vol7/iss4/13>

DOI: <https://doi.org/10.18785/grr.0704.13>

This Short Communication is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in Gulf and Caribbean Research by an authorized editor of The Aquila Digital Community. For more information, please contact Joshua.Cromwell@usm.edu.

A PRELIMINARY CHECKLIST OF EPIPHYTIC AND BENTHIC MARINE DIATOMS OF LOUISIANA

ROBERT S. MAPLES

Department of Biological and Environmental Sciences,
McNeese State University, Lake Charles, Louisiana 70609

ABSTRACT A checklist of diatoms collected from the various habitats in coastal salt marshes, estuaries, and nearshore Gulf waters of Louisiana has been compiled for the first time. The checklist includes 193 taxa (species and their varieties) in 38 genera. The largest number of taxa are species of *Navicula* and *Nitzschia* with 46 and 36, respectively. Future studies of the varied habitats along the Louisiana coast should result in many new additions to this preliminary checklist.

INTRODUCTION

The algal flora of coastal Louisiana has been variously studied since the first reports of Featherman (1871, 1872). Kapraun (1974) surveyed the marine benthic green, red, and brown algae found on jetty structures bordering Louisiana's marine passes. *Vaucheria* species were recorded by Pecora (1977, 1980) from coastal salt marsh habitats. Maples (1982) described an edaphic bluegreen algal community associated with a coastal salt panne in southwestern Louisiana. Checklists of marine planktonic diatoms in the nearshore waters of Louisiana are found in papers by Featherman (1871, 1872), Simmons and Thomas (1962), Bamforth (1974), Housley (1976), Fucik and El-Sayed (1979), and Maples (1983a). Studies of marine benthic diatoms have been largely centered in the Mississippi delta region of Louisiana. Cook and Whipple (1982) described the spatial and temporal distribution of edaphic diatom communities along a complex gradient from brackish to saline marshland. Stowe (1982) investigated the distribution of epiphytic diatoms on the culms of *Spartina alterniflora* Loisel. Maples (1983b) reported on the taxonomy, diversity, and similarity of benthic diatom assemblages on the pneumatophores of the black mangrove *Avicennia germinans* (L.) L., in a Louisiana salt marsh.

Kalinsky (1983) compiled a checklist of non-marine algae from Louisiana. Many species found in Louisiana by this

author were not included in Kalinsky's (1983) checklist. An up-to-date list of diatoms in Louisiana's coastal environments was thought to be needed. In light of the reports of Wood (1963) and Medlin (1983) for Texas, and Sullivan (1981) for Mississippi, the present checklist contributes to our knowledge of the geographic distribution of marine diatoms.

Three different papers were used in the preparation of this checklist (Cook and Whipple 1982, Stowe 1982, Maples 1983b) as well as unpublished observations by this author. The present checklist is biased toward the epiphytic diatom flora of coastal salt marshes, but edaphic (sediment-associated) diatom taxa are included. A number of taxa are primarily freshwater forms, but these may be occasionally observed in brackish water habitats. The flora reported herein is of an edaphic or epiphytic nature, and represents marsh, estuarine, and nearshore habitats.

CHECKLIST FORMAT

For the purpose of this checklist, diatoms were considered to constitute the single class Bacillariophyceae. Genera are arranged alphabetically, and species with their varieties are arranged alphabetically within each genus following the format of Hendey's (1974) checklist for British marine diatoms. Species and their varieties collected only from the pelagic alga *Sargassum* are marked with an asterisk.

BACILLARIOPHYCEAE

ACHNANTHES Bory, 1822

- biasoletiana* (Ag.) Grun.
- brevipes* var. *intermedia* (Kütz.) C1.
- curvirostrum* Brun.
- exigua* Grun.
- hauckiana* Grun.
- lanceolata* var. *dubia* Grun.
- lemmermanni* Hust.
- submarina* Hust.
- subsessilis* Kütz.

temperei M. Perag.

AMPHIPRORA Ehrenburg, 1843

- alata* Kütz.
- paludosa* W. Sm.

AMPHORA Ehrenburg, 1831

- angusta* Greg.
- var. *angusta*
- var. *oblongella* Grun.
- bigibba* Grun.
- caroliniana* Giffen
- coeffeiformis* (Ag.) Kütz.
- exigua* Greg.

- granulata* Greg.
laevis var. *perminuta* Grun.
libyca Ehr.
pediculus (Kütz.) Grun.
proteus Greg.
sabyii Salah
tenerrima Hust.
tenuissima Hust.
ventricosa Greg.
ANOMOENEIS Pfitzer, 1871
sphaeophora (Kütz.) Pfitzer
BACILLARIA Gmelin, 1778
paxillifer (Müll.) Hendey
BERKELEYA Greville, 1827
rutilans (Trent.) Grun.
CALONEIS Cleve, 1894
westii (W. Sms.) Hendey
CAMPYLOSIRA Grunow, 1882
alexandrica Salah
CAPARTOGRAMMA Kufferath, 1956
crucicula (Grun.) Ross
COCCONEIS Ehrenburg, 1838
fluviatilis Wallace
placentula Ehr.
var. *placentula*
var. *euglypta* (Ehr.) Grun.
pseudodiruptoides Foged*
scutellum Ehr.
var. *scutellum*
var. *parva* Grun.
CYCLOTELLA Kützing, 1833
atomus Hust.
caspia Grun.
comta (Ehr.) Kütz.
meneghiniana Kütz.
striata (Kütz.) Grun.
stylorum Brightwell
CYLINDROTHECA Rabenhorst, 1859
gracilis (Bréb.) Grun.
CYMATOSIRA Grunow, 1862
belgica Grun.
CYMBELLA Agardh, 1830
pusilla Grun.
DENTICULA Kützing, 1844
subtilis Grun.
DIPLONEIS Ehrenburg, 1840
didyma (Ehr.) Ehr.
interrupta (Kütz.) C1.
var. *interrupta*
var. *caffra* Giffen
pseudovalis Hust.
smithii (Bréb.) C1.
weissflogi (A.S.) C1.
EUNOTOGRAMMA Weisse, 1854
laevis Grun. (*laeve*)
- FRAGILARIA** Lyngbye, 1819
brevistriata Grun.
construens (Ehr.) Grun.
var. *construens*
var. *venter* (Ehr.) Grun.
GOMPHONEMA Agardh, 1824
gracile Ehr.
littorale Hendey
parvulum Kütz.
GRAMMATOPHORA Ehrenberg, 1840
oceanica Ehr.*
GYROSIGMA Hassall, 1845
balticum (Ehr.) Rabh.
beaufortianum Hust.
fascicola (Ehr.) C1.
hummi Hust.
obliquum Boyer
peisonis (Grun.) C1.
LICMOPHORA Agardh, 1827
abbreviata Ag.
cf. *debilis* (Kütz.) Grun.
remulus Grun.*
MASTOGLOIA Thwaites, 1856
acutiuscula Grun.*
binotata (Grun.) C1.
crucicula (Grun.) C1.
dubia Kütz.
erythraea Grun.*
exigua Lewis
jurgensii Ag.
ovalis A.S.*
pumila (Grun.) C1.
pusilla (Grun.) C1.
var. *pusilla*
var. *subcapitata* Hust.*
MELOSIRA Agardh, 1824
nummuloides Ag.
NAVICULA Bory, 1822
abunda Hust.
accomoda Hust.
aequorea Hust.
ammophila Grun.
amphipleuroides Hust.
capitata var. *hungarica* (Grun.) Ross
circumtexta Meister
contenta Grun.
creuzburgensis Krasske
crucicula (W.Sm.) Donk.
cryptocephala Kütz.
cryptolyra Brockman
diserta Hust.
elegans W. Sm.
flanatica Grun.
gregaria Donk.
hudsonis Grun.
hyalinula DeToni

incomposita Hagelstein
 var. *incomposita*
 var. *minor* Hagelstein
lanceolata (Ag.) Kütz.
marina Ralfs
menisculus Schum.
mutica (Hilse) Grun.
 var. *mutica*
 var. *stigma* Patr.
nolens Simonsen
obsoleta Hust.
pavillardi Hust.
peregrina (Ehr.) Kütz.
phyllepta Kütz.
platyventris Meister
comoides (Ag.) Perag.
pseudocrassirostris Hust.
pupula var. *rectangularis* (Greg.) Grun.
salinarum Grun.
salinicola Hust.
schroeteri Meister
spicula (Hickie) C1.
subforcipata Hust.
taraxa Hohn & Hellerm.
tenera Hust.
teneroides Hust.
tripunctata (Müll.) Bory
 var. *tripunctata*
 var. *schizonemoides* (V.H.) Patr.
yarrenis Grun.
zostereti Grun.
NITZSCHIA Hassall, 1845
amphibia Grun.
angularis W. Sm.
apiculata (Greg.) Grun.
bicapitata C1.*
bilobata var. *ambigua* Mang.
brittonia Hagelstein
closterium W. Sm.
communis var. *hyalina* Lund
deblis (Arnott) Grun.
dissipita (Kütz.) Grun.
epithemioides Grun.
fasciculata (Grun.) Grun.
filiformis (W. Sm.) Schutt
frustulum (Kütz.) Grun.
 var. *frustulum*
 var. *perminuta* Grun.
gandersheimiensis Krasske
grana Hohn & Hellerm.
granulata Grun.
hungarica Grun.
lanceolata W. Sm.
lorenziana var. *subtilis* Grun.
minutula Grun.
microcephala Grun.

obtusia W. Sm.
 var. *obtusia*
 var. *nana* Grun.
palea (Kütz.) W. Sm.
panduriformis Greg.
 var. *panduriformis*
 var. *continua* Greg.
perversa Grun.
romana Grun.
romanoides Mang.
scalaris (Ehr.) W. Sm.
sigma (Kütz.) W. Sm.
tryblionella Hantz.
vitrea Norman
 var. *vitrea*
 var. *salinarum* Grun.
OPEPHORA Petit, 1888
 parva (Grun.) Krasske
PARALIA Heiberg, 1863
 sulcata (Ehr.) C1.
PLEUROSIGMA W. Smith, 1852.
 angulatum (Quek.) W. Sm.
 normanii Ralfs
 salinarum (Grun.) Grun.
PSAMMODISCUS Round & Mann, 1980
 nitidus (Greg.) Round & Mann
RHOPALODIA O. Müller, 1895
 gibberula (Ehr.) Müll.
 musculus var. *producta* Grun.
STAURONEIS Ehrenberg, 1984
 amphioxys Greg.
 var. *amphioxys*
 var. *obtusata* Hendey
 legeri Hust.
STRIATELLA Agardh, 1832
 unipunctata (Lyngb.) Ag.*
SURIRELLA Turpin, 1828
 angusta Kütz.
 atomus Hust.
 ovalis Bréb.
SYNEDRA Ehrenberg, 1830
 affinis Kütz.
 fasciculata (Ag.) Kütz.
 var. *fasciculata*
 var. *intermedia* M. Sullivan
 demerarae Grun.
 provincialis var. *tortuosa* Grun.*
 tabulata var. *parva* (Kütz.) Hust.
THALASSIOSIRA Cleve, 1873
 eccentrica (Ehr.) C1.
 leptopus (Grun.) Hasle & Fryxell
TRACHYSPHENIA Petit, 1877
 acuminata Perag.*
TROPIDONEIS Cleve, 1891
 lepidoptera (Greg.) C1.
 vitrea (W. Sm.) C1.

GENERAL SUMMARY

A total of 193 taxa (species and their varieties) representing 38 genera comprise the present checklist. Genera with the largest number of taxa are *Navicula* and *Nitzschia* with 46 and 36, respectively. The next most abundant taxa were species of *Amphora* (15), *Mastogloia* (11), and *Achnanthes* (10). Also well represented are *Cocconeis*, *Cyclotella*,

Diploneis, *Gyrosigma*, and *Synedra* with six taxa each. Eleven taxa were collected only from *Sargassum* in the near-shore waters of southwestern Louisiana.

The present checklist provides important distributional information of an ecologically important group of organisms. Future studies of the varied habitats along the Louisiana coast should result in many new additions to this preliminary checklist.

REFERENCES CITED

- Bamforth, S. S. 1974. Coastal plankton off Grand Isle, Louisiana. *Proc. La. Acad. Sci.* 36:64–69.
- Cook, L. L. & S. A. Whipple. 1982. The distribution of edaphic diatoms along environmental gradients of a Louisiana salt marsh. *J. Phycol.* 18:64–71.
- Featherman, A. 1871. Second annual report of botanical survey of southwest and northwest Louisiana, made during the year 1870. Pages 63–131 in: Annual Report of Prof. D. F. Boyd, Superintendent, Louisiana State University, for year 1871, to the Governor of State of Louisiana. New Orleans, Louisiana.
- . 1872. Third annual report of botanical survey of southwest and northwest Louisiana, made during the year of 1870. Pages 101–161 in: Annual Report of Prof. D. F. Boyd, Superintendent, Louisiana State University, for year 1871, to the Governor of State of Louisiana. New Orleans, Louisiana.
- Fucik, K. W. and S. Z. El-Sayed. 1979. Effect of oil production and drilling operations on the ecology of phytoplankton in the OEI study area. Pages 325–353 in: C. H. Ward, M. E. Bender and D. J. Reish (eds.), *The Offshore Ecology Investigation*, Vol. 65, *Effects of Oil Drilling and Production in a Coastal Environment*. Rice University, Houston, Texas.
- Hendey, N. I. 1974. A revised checklist of British marine diatoms. *J. Mar. Biol. Assoc. U. K.* 54:277–300.
- Housley, H. L. 1976. Distribution, Periodicity, and Identification of the Phytoplankton in the Bay of St. Louis, Mississippi and the Northeastern Gulf of Mexico. Ph.D. Dissertation, University of Southern Mississippi, Hattiesburg. 206 pp.
- Kalinsky, R. G. 1983. Notes on Louisiana algae. II. A checklist of the non-marine algal flora of Louisiana. *Proc. La. Acad. Sci.* 46: 62–96.
- Kapraun, D. F. 1974. Seasonal periodicity and spatial distribution of benthic marine algae in Louisiana. *Contrib. Mar. Sci.* 18: 138–167.
- Maples, R. S. 1982. Bluegreen algae of a coastal salt panne and surrounding angiosperm zones in a Louisiana salt marsh. *North-east Gulf Sci.* 5:39–43.
- . 1983a. A preliminary checklist of marine planktonic diatoms of southwestern Louisiana. *Proc. La. Acad. Sci.* 46: 34–40.
- . 1983b. Community structure of diatoms epiphytic on pneumatophores of the black mangrove, *Avicennia germinans*, in a Louisiana salt marsh. *Gulf Res. Rept.* 7(3):255–259.
- Medlin, L. K. 1983. Community Analysis of Epiphytic Diatom Communities Attached to Selected Species of Macroalgal Host Plants Along the Texas Gulf Coast. Ph.D. Dissertation. Texas A&M University, College Station. 150 pp.
- Pecora, R. A. 1977. Brackish water species of *Vaucheria* (Xanthophyceae, Vaucheriales) from Louisiana and Texas. *Gulf Res. Rept.* 6(1):25–29.
- . 1980. Observations on the genus *Vaucheria* (Xanthophyceae, Vaucheriales) from the Gulf of Mexico. *Gulf Res. Rept.* 6(4):387–391.
- Simmons, E. G. and W. H. Thomas. 1962. Phytoplankton of the eastern Mississippi delta. *Contrib. Mar. Sci.* 8:269–288.
- Stowe, W. C. 1982. Diatoms epiphytic on the emergent grass, *Spartina alterniflora*, in a Louisiana salt marsh. *Trans. Am. Microsc. Soc.* 101:162–173.
- Sullivan, M. J. 1981. A preliminary checklist of marine benthic diatoms of Mississippi. *Gulf Res. Rept.* 7(1):13–18.
- Wood, E. J. F. 1963. A study of the diatom flora of fresh sediments of the South Texas bays and adjacent waters. *Publ. Inst. Mar. Sci. Univ. Tex.* 9:237–310.