

Gulf and Caribbean Research

Volume 9 | Issue 2

January 1995

A Study of Factors Influencing the Hatch Rate of *Penaeus vannamei* Eggs. I. Effects of Size, Shape and Volume of the Spawning Tank

John T. Ogle
Gulf Coast Research Laboratory

Follow this and additional works at: <https://aquila.usm.edu/gcr>



Part of the [Marine Biology Commons](#)

Recommended Citation

Ogle, J. T. 1995. A Study of Factors Influencing the Hatch Rate of *Penaeus vannamei* Eggs. I. Effects of Size, Shape and Volume of the Spawning Tank. *Gulf Research Reports* 9 (2): 123-126.
Retrieved from <https://aquila.usm.edu/gcr/vol9/iss2/7>
DOI: <https://doi.org/10.18785/grr.0902.07>

This Article is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in Gulf and Caribbean Research by an authorized editor of The Aquila Digital Community. For more information, please contact Joshua.Cromwell@usm.edu.

A STUDY OF FACTORS INFLUENCING THE HATCH RATE OF *PENAEUS VANNAMEI* EGGS. I. EFFECTS OF SIZE, SHAPE AND VOLUME OF THE SPAWNING TANK

John T. Ogle

Fisheries Section, GulfCoast Research Laboratory, P.O. Box 7000, Ocean Springs, Mississippi 39566-7000, USA

ABSTRACT The hatch rate of *Penaeus vannamei* eggs spawned by individual females in square and round tanks and in different volumes of seawater was determined. The mean hatch rates ranged from 17.3% to 64.9% and were not significantly different for volumes of 50, 100 and 200 L nor for square or round tanks of equal water volumes. Hatch rate was significantly affected by the size of the spawning tank.

INTRODUCTION

Aquaculture accounts for over 25% of the world's shrimp production (Rosenberry, 1991). The farming of shrimp requires acquisition of seed stock which is reared to marketable size. Seedstock may be harvested from the wild or produced from broodstock. Broodstock may be sourced from the wild or acquired from maturation facilities. In either case, mated oviparous shrimp are spawned under captive conditions in order to collect the larvae. A variety of systems have been used to spawn marine shrimp. Eggs or nauplii may be collected directly from the tank in which maturation and matings take place (Laubier-Bonichon and Laubier 1979; Brown et al. 1980; Lawrence et al. 1980; Simon 1982; Chen et al. 1991; Ogle 1992). However, many facilities remove the mated females to a separate tank for spawning. Wyban and Sweeney (1991) and Bray and Lawrence (1992) report the stocking of a single mated female into a tank while Lawrence et al. (1980), Kittaka (1981), Aquacop (1983), Yang (1975), Tabb et al. (1975) and Treece (1985) cite that several mated females may be placed in a common tank for spawning. In most cases, spawning and hatching are accomplished in one tank and the nauplii are collected and transferred to another tank for larval rearing. However, Salser (1978) and Bray and Lawrence (1992) recommend that after the shrimp spawn, eggs be removed to a separate tank for hatching. Tanks utilized for spawning and hatching have been as large as 145,800 L (Kittaka, 1981) and as small as 10 L (Mock and Murphy 1971). Bray and Lawrence (1992) suggest 75 L as the minimum tank size with 100 to 150 L circular tanks supplied with lids as the preferred choice for spawning and hatching. Browdy (1992) suggests using a tank size of 150-500 L and Browdy and Samocha (1985a and 1985b) report that simple flat-bottom tubs are sufficient for the spawning

and hatching of eggs. Aquacop (1983) utilizes conical bottom tanks. A comparison of the two tank types, conical and flat-bottomed, with *P. setiferus* could demonstrate no significant differences in hatch rates (Browdy, in preparation). Lotz and Ogle (1994) report an increasing hatch rate of *P. vannamei* with an increasing volume of the spawning tanks. However, the spawning tanks varied in size and shape. As these are the only comparative studies on the effect that tank shape and volume have on hatch rates, the present study was undertaken with *P. vannamei*.

MATERIAL AND METHODS

Three different tank types were used for spawning mated *P. vannamei* females. Rectangular polyethylene tubs, 0.51 m x 0.53 m x 0.32 m, with a bottom area of 0.27 m² were utilized as one of the tank types. The other two tank types were both round fiberglass tanks and are referred to as small and large. The small tanks had a diameter of 0.61 m and a depth of 0.45 m with a bottom area of 0.29 m² while the large tanks had a diameter of 1.12 m and a depth of 0.60 m with a bottom area of 0.98 m² (1 m²).

The spawning tanks were filled with a measured volume of seawater. Natural baywater pumped from Davis Bayou in the Mississippi Sound was allowed to stand for several weeks to allow suspended solids to settle. Artificial seasalt (Marine Environment, San Francisco, CA) was added to increase the ambient salinity of 25 ppt to a salinity of 30 ppt. The water was filtered through a particulate five-micron RS pleated polyester fabric media cartridge (Amtek, Flowrite, Inc., 3345 Halls Mill Rd., Mobile, AL 36606) and granular activated carbon (Amtek) before use. Sodium EDTA was added at the rate of 3 ppm.

Shrimp were matured and mated in large commercial sized maturation tanks (Ogle 1992). Mated females were sourced from the maturation tanks in the evening and placed individually into the spawning tanks. Moderate aeration was provided by a single airstone throughout spawning and hatching. The shrimp were checked for spawning after two to three hours and spent females were returned to the maturation tanks.

The number of eggs was estimated by subsampling. The water in the spawning tank was stirred and five-10 ml subsamples were collected. Subsamples were taken from the four compass directions and the center of the tank. The samples were transferred to a petri dish and the eggs counted. Data were averaged and the total number of eggs calculated. After 12 -15 hours, the number of nauplii was determined in the same fashion and the hatch rate calculated (% H = # nauplii/# eggs x 100). Only spawns that hatched were analyzed. Data were compared by ANOVA and significant (alpha = 0.05) differences noted.

The large (1m²) tanks were used to study the effect of water volume on hatch rate with 50, 100 or 200 L of seawater placed in a tank for each shrimp. Twelve shrimp were individually spawned in each of the three water volumes, for a total of 36 shrimp. Fifty L of seawater were used in all subsequent studies. The effect of tank shape on hatch rate was determined by comparing spawns in the square tubs, 0.27 m², and the small fiberglass round tanks,

0.29m². A total of 14 individual spawns were recorded for the square tanks and 15 individual spawns were recorded for the small round tanks. An additional 14 animals spawning in the small round fiberglass tanks were compared to 14 animals which spawned in the large round fiberglass tanks to determine the effect of tank size on hatch rate. In all studies, eggs were left in the spawning tanks until hatching occurred. After the eggs hatched, the tanks were drained, cleaned and filled with new seawater.

RESULTS

There were no significant differences for the effect of the water volume in the spawning tanks or the shape of the spawning tanks on hatch rates of *P. vannamei* eggs (Table 1). The average hatch rates for eggs spawned in 50, 100 and 200 L of water were 45.2, 55.2, and 42.4%, respectively. The average hatch rates for eggs spawned in the square and the small round tanks were 25.4 and 18.2%, respectively.

There was a significant effect of tank size on the hatch rates. The hatch rates for eggs spawned into a tank of 0.29 m² and 1 m² were 17.3% and 64.9%, respectively.

Overall, the minimum spawn size was 39,000 eggs and the maximum spawn size was 230,000 eggs. The minimum egg density was 780/L and the maximum was 4,600 eggs/L. This is equivalent to 10,000 to 900,000 eggs/m².

TABLE 1

The effect of tank volume, shape and size on the hatch rate of *Penaeus vannamei* eggs.

Study	Treatment	Replicates N	Hatch Rate			Significance
			Mean %	Range %	S.E.	
Volume	50 L	12	45.2	3.8 - 89.3	7.10	NS
	100 L	12	55.2	15.2 - 100.0	7.21	NS
	200 L	12	42.4	1.5 - 90.0	8.54	NS
Shape	Square	14	25.4	3.8 - 57.8	4.15	NS
	Round	15	18.2	2.4 - 51.3	5.26	NS
Size	0.29 m ²	14	17.3	2.0 - 51.3	4.15	SIG
	1.00 m ²	14	64.9	15.2 - 100.0	6.69	SIG

Volume - 1 m² round fiberglass tank. Shape - square, 0.27 m² polyethylene tanks with 50 L of water; round, 0.29 m² fiberglass tanks with 50 L of water. Size - round tanks with 50 L of water. SIG = significant. NS = non-significant.

DISCUSSION

While there was a significant effect of the tank size on the hatch rate of *P. vannamei* eggs, the reason is unclear. The volume in both the large and small tanks was 50 L which created a greater water depth in the smaller tank. The difference in depth does not appear to have an effect on hatching as demonstrated in the experiment on water volumes in this study. This suggests that as long as the female is allowed a sufficient area for unrestricted spawning, the depth of the water beneath the shrimp appears to have no effect on hatch rate. Instead, tank area may play an important role in hatch rate. The average hatch rate reported for *P. vannamei* of 30% to 47% (Wyban and Sweeney 1991) is consistent with the rates reported here of 17.3% to 64.9%.

It should be pointed out that the hatch rates reported here include three discrete events: spawning, fertilization and hatching. Spawning of *P. vannamei* occurs as the female slowly swims in circles near the surface of the water. It is possible that restricting the movement of the shrimp to a smaller area may interfere with fertilization of the eggs which would result in a lower hatch rate. It is also possible that the hatch rate may be directly influenced by the bottom area of the tank.

As the mechanisms involved in egg fertilization are not known, it is unclear how restricting the swimming activity of the shrimp during spawning would influence fertilization and hatch rates. Heldt (1938) and Hudinaga (1942) have suggested that sperm released from the spermatophore are trapped by the ventral setae of the third

and fourth pereopods which come into contact with the eggs as they are released. We have been unable to detect sperm on the pleopods or pereopods of mated *P. vannamei* by microscopic examination. Fertile spawns frequently result from matings for which the spermatophore is missing. Fertile spawns have also resulted from females which do not swim but lie perfectly still during spawning. In such incidence, the hatch rate is extremely low (Ogle 1993b).

The effect of egg density on hatching is also unknown. Primavera (1980) has recommended that the egg density in spawning tanks for *P. monodon* not exceed 2,500 to 3,000 eggs per liter. In another study, Primavera *et al.* (1977) found no effect on the hatching of *P. monodon* eggs at densities of 7000/l. However, they did not report the results in terms of area (eggs/m²). *P. monodon* eggs are larger than *P. vannamei* eggs (Ogle, in preparation) so the densities used here of 780 - 4,600 eggs/L should have no effect on the results. This is also supported by the fact that the volume of water did not significantly affect the hatch rate. Further research could be directed to determine the effect of tank area on fertilization rates and hatch rates.

ACKNOWLEDGMENTS

Appreciation is expressed to Casey Nicholson and Richard King for the daily operation of the maturation systems, Leslie Christmas for data reduction, Kathy Beaugez for aid in manuscript preparation and Dr. Terry McBee for statistical analysis. This project is funded by USDA, CSMR Grant Nos. 2-2537 and 2.2538.

LITERATURE CITED

- Aquacop. 1983. Constitution of broodstock, maturation, spawning and hatching systems for penaeid shrimps in the Centre Oceanologique du Pacifique. In: J.P. McVey (ed.), CRC Handbook of Mariculture, Vol. 1, Crustacean Aquaculture, p 105-121. CRC Press, Boca Raton, FL.
- Bray, W.A. and A.L. Lawrence. 1992. Reproduction of *Penaeus* species in captivity. In: A.W. Fast and L.J. Lester (eds.), Marine Shrimp Culture: Principles and Practices, p 105-121. Elsevier Sci Publ B.V. Amsterdam, The Netherlands. 962 p.
- Browdy, C. 1992. A review of the reproductive biology of *Penaeus* species: Perspectives on controlled shrimp maturation systems for high quality nauplii production. In: J. Wyban (ed.), 1992 Proc Spec Sess on Shrimp Farming, p 22-51. World Aquacult Soc, Baton Rouge, LA. 301 p.
- Browdy, C.L. and T.M. Samocha. 1985a. Maturation and spawning of ablated and nonablated *Penaeus semisulcatus* de Haan (1844). J World Maricult Soc 16:236-249.
- Browdy, C.L. and T.M. Samocha. 1985b. The effect of eyestalk ablation on spawning, molting and mating of *Penaeus semisulcatus* de Haan. *Aquaculture* 49(1):19-29.
- Brown Jr., A., J. McVey, B.M. Scott, T.D. Williams, B.S. Middleditch, and A.L. Lawrence. 1980. The maturation and spawning of *Penaeus stylirostris* under controlled laboratory conditions. Proc World Maricult Soc 11:488-499.
- Chen, F., B. Reid, and C.R. Arnold. 1991. Maturing, spawning and egg collecting of the white shrimp *Penaeus vannamei* Boone in a recirculating system. J World Aquacult Soc 22(3):167-172.
- Heldt, J.H. 1938. La reproduction chez les Crustacés Décapodes de la famille des Pénéides. Ann Inst Oceanogr 18(2):31-206.
- Hudinaga, M. 1942. Reproduction, development and rearing of *Penaeus japonicus* Bate. Jpn J Zool 10(2):305-393.
- Kittaka, J. 1981. Large scale production of shrimp for releasing in Japan and in the United States and the results of the releasing programme at Panama City, Florida. Kuwait Bull Mar Sci 137(2):149-163.

- Laubier-Bonichon, A. and L. Laubier. 1979. Controlled reproduction of the shrimp *Penaeus japonicus*. In: T.V.R. Pillay and Wm. A. Dill (eds.), *Advances in Aquaculture*, p 273-276. Papers presented at the FAO Tech Conf on Aquacult. Kyoto, Japan, 26 May-2 June. Fishing News Book Ltd. England. 653 pp.
- Lawrence, A.L., Y. Akamine, B.S. Middleditch, G.W. Chamberlain, and D.L. Hutchins. 1980. Maturation and reproduction of *Penaeus setiferus* in captivity. *Proc World Maricult Soc* 11:481-487.
- Lotz, J. and J.T. Ogle. 1994. Reproductive performance of the white legged shrimp *Penaeus vannamei* in recirculating seawater systems. *J World Maricult Soc* 25(3):477-482.
- Mock, C.R. and M.A. Murphy. 1971. Techniques for raising penaeid shrimp from egg to postlarvae. *Proc World Maricult Soc* 1:143-156.
- Ogle, J.T. 1992. Design and operation of the Gulf Coast Research Laboratory penaeid shrimp maturation facility I. *Penaeus vannamei*. *Gulf Coast Res Lab Tech Rep* 4, 41 p.
- _____. 1993. The state of our knowledge concerning reproduction in open thelycum penaeid shrimp with emphasis on *Penaeus vannamei*. *Invertebr Reprod Dev* 22(1-3):267-274.
- Primavera, J.H. 1980. Broodstock of sugpo (*Penaeus monodon*) and other penaeid prawns. *Aquacul Dept, SEAFDEC, Aquacul Ext Manual No. 7*. 24 p.
- Primavera, J.H., E. Borlongan and R.A. Posadas. 1977. Viability of *P. monodon* eggs after simulated transport condition. *Quarterly Res Rep SEAFDEC* 1(4):11-13.
- Rosenberry, B. 1990. Shrimp farming in the western hemisphere. In: M.B. New, H. de Saram, and T. Singh (eds.), *Technical and Economic Aspects of Shrimp Farming*, p 223-251. *Proc Aquatech 90 Conf Kuala Lumpur, Malaysia, 11-14 June 1990*. INFOFISH. 341 p.
- Salser, B. Larval penaeid shrimp culture techniques utilized by the Environmental Research Laboratory at Puerto Penasco, Sonora, Mexico. Unpublished manuscript.
- Simon, C.M. 1982. Large-scale commercial application of penaeid shrimp maturation technology. *J World Maricult Soc* 13:301-312.
- Tabb, D.C., W.T. Yang, Y. Hirono, and J. Heinen. 1972. A manual for culture of pink shrimp, *Penaeus duorarum* from eggs to post-larvae suitable for stocking. *Sea Grant Spec Bull* 7, Univ Miami, 59 p.
- Treece, G.D. 1985. Larval rearing technology. In: G.W. Chamberlain, M.G. Haby, and R.J. Miget (eds), *Texas Shrimp Farming Manual*, p III43-64. *Texas Agri. Ext. Serv. publ. invited papers presented at the Texas Shrimp Farming Workshop. 19-20 Nov. 1985, Corpus Christi, TX*.
- Wyban, J.A. and J.N. Sweeney. 1991. *Intensive shrimp production technology: The Oceanic Institute shrimp manual*. Oceanic Institute, Honolulu, HI, 158 pp.
- Yang, W.T. 1975. A manual for large-tank culture of penaeid shrimp to the postlarval stages. *Sea Grant Tech Bull* 31, Univ Miami, 93 p.