

Gulf and Caribbean Research

Volume 16 | Issue 1

January 2004

Notes on Aquarium Brood Release and Feeding of the Opossum Pipefish, *Microphis brachyurus lineatus*

Sarah Frias-Torres
Florida Institute of Technology

DOI: 10.18785/gcr.1601.11

Follow this and additional works at: <http://aquila.usm.edu/gcr>

 Part of the [Marine Biology Commons](#)

Recommended Citation

Frias-Torres, S. 2004. Notes on Aquarium Brood Release and Feeding of the Opossum Pipefish, *Microphis brachyurus lineatus*. Gulf and Caribbean Research 16 (1): 73-75.

Retrieved from <http://aquila.usm.edu/gcr/vol16/iss1/11>

This Short Communication is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in Gulf and Caribbean Research by an authorized editor of The Aquila Digital Community. For more information, please contact Joshua.Cromwell@usm.edu.

SHORT COMMUNICATION**NOTES ON AQUARIUM BROOD RELEASE AND FEEDING OF THE OPOSSUM PIPEFISH, *MICROPHIS BRACHYURUS LINEATUS*****Sarah Frias-Torres***Department of Marine and Environmental Systems, Florida Institute of Technology, 150 West University Blvd, Melbourne, FL 32901 USA**Current Address: NOAA, Southeast Fisheries Science Center, 75 Virginia Beach Drive, Miami, Florida 33149 USA, E-mail Sarah.FriasTorres@noaa.gov***INTRODUCTION**

The opossum pipefish, *Microphis brachyurus lineatus*, is a migratory syngnathid with a salmon-like life history (Frias-Torres 2002). Adults breed in freshwater and oligohaline habitats associated with emergent vegetation, mostly *Panicum* sp. and *Polygonum* sp.; breeding occurs during the wet season (May to November) (Gilmore and Hastings 1983). In southern Mexico, opossum pipefish associate with *Ruppia maritima*, and reproduction is year-round (Miranda-Marure et al. 2004). Pelagic juveniles are found in the Atlantic Ocean and are associated with *Sargassum* sp. (Böhlke and Chaplin 1968). Juveniles eventually recruit into oligohaline habitats. The species range is from Sao Paulo, Brazil, to New Jersey, USA, although the northernmost limit of permanent breeding populations occurs in the freshwater tributaries to the Indian River Lagoon in east central Florida (Gilmore and Gilbert 1992). The opossum pipefish is one of 25 species of diadromous fish at risk of extinction in North America (Musick et al. 2000) and is a candidate species to the US Threatened and Endangered Species List (Federal Register 1999).

The opossum pipefish diet was unknown until Teixeira and Perrone (1998) analyzed the gut contents of 109 opossum pipefish from the southeastern Brazil population. They found that the most abundant prey were insect larvae, followed by juvenile fish (Gerreidae) and small crustaceans. However, the diet necessary to maintain aquarium-kept opossum pipefish had not been determined prior to the present paper.

Male Syngnathidae (seahorses and pipefishes) carry a brood of fertilized eggs in a partially or completely closed pouch, depending on the species. Hatching occurs inside the pouch, followed by release of the young syngnathids.

Brood release has been documented for common pipefish and seahorse species that have been successfully kept in aquaria (Garrick-Maidment 1997). Gilmore (1977) reported opossum pipefish brood release in a freshwater aquarium, but the actual event of brood release was not observed. Long-term aquarium keeping of opossum pipefish has not been attempted before, and many fundamental questions of the biology of the species remain unresolved.

In 350 BC, Aristotle was the first scientist to describe the process of brood release in Syngnathidae, probably *Syngnathus acus*, a common pipefish with a pouch covered by a fleshy membrane found in inshore coastal areas of the Mediterranean Sea:

“When the time of parturition arrives, [the pipefish] bursts in two, and the eggs escape out (...) the fish has a diaphysis or cloven growth under the belly and abdomen, and after it has spawned by the splitting of this diaphysis, the sides of the split grow together again. (...) The young fish cluster round the parent (...) for the fish spawns onto herself; and if any one touch the young, they swim away.”

The goal of the present study was to gain insight into adult feeding and brood release of opossum pipefish in captivity as a step towards a better understanding of the biology of the species. Two primary objectives were: 1) to apply Teixeira and Perrone’s field observations to the diet supplied to captive opossum pipefish and 2) to video record an opossum pipefish brood release event.

METHODS

Male-female pairs of opossum pipefish were collected at the Fellsmere Canal, Sebastian River, east central Florida, during the wet season (May to November) of 2000. Collections were targeted to areas with underwater bank vegetation using a 30 x 40 cm square dipnet with an outer net of 4 mm and an inner net of 0.5 mm mesh size. Fish were transferred into 25 L coolers equipped with air-stones and filled with water and vegetation from the capture site. Additional *Panicum* and water were collected to establish aquaria in the laboratory. Adult male-female pairs and brooding males were transferred to freshwater 50 L glass aquaria fitted with standard undergravel filters and air stones. Several stems of *Panicum* were kept at the surface and half submerged. Illumination was set on a 13:11 light:dark cycle. Water and air temperature was kept at 25 °C.

Captive opossum pipefish were presented with a variety of dried and live food types (brine shrimp, amphipods,

freshwater shrimp and guppies) one at a time, and their reaction (consumption or rejection) over a 15 min observation period was recorded. Insect larvae were not tested as potential diet for captive opossum pipefish. Daily observation of embryo development determined when brood release was imminent. At that point, a 24 h watch was kept to allow for video recording of a brood release event.

RESULTS AND DISCUSSION

Opossum pipefish rejected brine shrimp (*Artemia salina*) and amphipods, the 2 most common types of dried and frozen foods usually sold in pet stores as ideal feeding for Syngnathidae. Opossum pipefish also rejected live amphipods but consumed all other types of live food offered: *A. salina* adults (which survive a few minutes in freshwater); juvenile freshwater glass shrimp (*Palaemonetes kadiakensis* Rathbun); fry, juveniles, and small adults of guppies (*Poecilia* sp.); and mosquitofish (*Gambusia* sp.). To ensure a constant supply of food, wild and commercially available freshwater shrimp, guppies and mosquitofish were fed to the opossum pipefish.

Brood release occurred several times in aquaria. The actual event of larval release was first observed from a 132 mm SL brooding male, 9 d after capture. Male opossum pipefish have an open brood pouch under the tail, where eggs are protected with bilateral pouch plates but are not covered by a pouch membrane (subfamily Doryrhamphinae). Therefore, direct observation of every stage of larval release was possible. On August 22, 2000 a power outage interrupted the automatic light cycle at 2:00 PM. At 3:00 PM the male started to release the brood. During brood release, the male remained close to the bottom, in a diagonal head-up orientation. The male vibrated for 2 sec from side to side, whereupon a string of 5 to 6 eggs detached from the pouch in a tail-to-head direction. Beginning with the most distal egg, the egg membrane ruptured, releasing a curled larva. Released larva then uncurled and with an undulating movement swam away from the male. In the frontal and medial sections of the pouch, direct hatching (without a preliminary release of a string of eggs) was observed (Figure 1). In this case, the larvae were violently ejected from the pouch. The observed brood release event lasted 35 min. In comparison, release of the seahorse brood (*Hippocampus* sp.) was preceded for up to 12 h of contractions, with the actual release taking just a few seconds (Garrick-Maidment, 1997).

Aquarium observations suggest that in the wild, opossum pipefish brood release occurs at dusk or at night. The power outage in the early afternoon, and the observed hatching that followed the dark period seems to confirm a




Figure 1. A 135 mm SL male opossum pipefish, *Microphis brachyurus lineatus*, releasing a 6 mm larvae. Sequence extracted from a 54 minute video recording.

dusk or nocturnal preference for larval release. Larval release was observed and video recorded for other brooding males during the wet season of 2000. All males followed a similar pattern of release at dusk, or soon after midnight, with vibrations preceding pouch release. In cases when the brood release was not observed, larvae were first seen swimming in the aquarium either late at night or just after sunrise.

ACKNOWLEDGMENTS

This work presents results of the author's doctoral dissertation submitted in 2002 to the Department of Marine and Environmental Systems, Florida Institute of Technology, Melbourne, Florida, USA. The author thanks her dissertation advisor R.G. Gilmore and co-advisor G.A. Maul for support during fieldwork and laboratory studies. I am also grateful to J. Javeck at the SEFSC in Miami for video technical support.

LITERATURE CITED

- Aristotle. 1991. *The History of the Animals*. Translated by A.L. Peck and D.M. Balme. Loeb Classical Library, Vol. 3, Book 10. Harvard University Press, Cambridge, MA, USA.
- Böhlke, J.E. and C.G.C. Chaplin. 1968. *Fishes of the Bahamas and adjacent tropical waters*. Livingston Press, Philadelphia, PA, USA, 771 p.
- Federal Register. 1999. Endangered and threatened species. Revision of candidate species list under the Endangered Species Act 64(120): 33466–33468.
- Frias-Torres, S. 2002. Oceanic transport and life history of the tropical western Atlantic opossum pipefish, *Microphis brachyurus lineatus*. Ph.D. dissertation. Florida Institute of Technology, Melbourne, FL, USA, 188 p.
- Garrick-Maidment, N. 1997. *Seahorses: Conservation and care*. T.F.H. Kingdom Books, London, England, 48 p.
- Gilmore, R.G. 1977. Notes on the opossum pipefish *Oostethus lineatus* from the Indian River Lagoon and vicinity, Florida. *Copeia* 1977:781–783.
- Gilmore, R.G. and C.R. Gilbert. 1992. Coastal habitat description (introduction) and species accounts for *Microphis brachyurus lineatus*. In: C.R. Gilbert, ed. *Rare and Endangered Biota of Florida: Fishes*, Vol. 2. University Presses of Florida, Gainesville, FL, USA, 247 p.
- Gilmore, R.G. and P.A. Hastings. 1983. Observations on the ecology and distribution of certain tropical peripheral fishes in Florida. *Florida Scientist* 46:31–51.
- Miranda-Marure, M.E., J.A. Martínez-Pérez, and N.J. Brown-Peterson. 2004. Reproductive biology of the opossum pipefish, *Microphis brachyurus lineatus*, in Tecolutla Estuary, Veracruz, Mexico. *Gulf and Caribbean Research* 16:101–108.
- Musick, J.A. et al. (17 more authors). 2000. Marine, estuarine, and diadromous fish stocks at risk of extinction in North America (exclusive of Pacific salmonids). *Fisheries* 25(11):6–30.
- Teixeira, R.L. and E.C. Perrone. 1998. Reproductive and feeding biology of freshwater and estuarine populations of the opossum pipefish, *Oostethus lineatus*, from southeastern Brazil. *Boletim do Museu de Biologia Mello Leitao (Nova Serie)* 8:3–12.