

January 2005

## Trophic Relationships of Demersal Fishes in the Shrimping Zone Off Alvarado Lagoon, Veracruz, Mexico

Edgar Pelaez-Rodriguez  
*University of Southern Mississippi*

Jonathan Franco-Lopez  
*Universidad Nacional Autonoma de Mexico*

Wilfredo A. Matamoros  
*University of Southern Mississippi*

Rafael Chavez-Lopez  
*Universidad Nacional Autonoma de Mexico*

*See next page for additional authors*

Follow this and additional works at: <https://aquila.usm.edu/gcr>



Part of the [Marine Biology Commons](#)

---

### Recommended Citation

Pelaez-Rodriguez, E., J. Franco-Lopez, W. A. Matamoros, R. Chavez-Lopez and N. J. Brown-Peterson. 2005. Trophic Relationships of Demersal Fishes in the Shrimping Zone Off Alvarado Lagoon, Veracruz, Mexico. *Gulf and Caribbean Research* 17 (1): 157-167. Retrieved from <https://aquila.usm.edu/gcr/vol17/iss1/16>  
DOI: <https://doi.org/10.18785/gcr.1701.16>

This Article is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in *Gulf and Caribbean Research* by an authorized editor of The Aquila Digital Community. For more information, please contact [Joshua.Cromwell@usm.edu](mailto:Joshua.Cromwell@usm.edu).

---

# Trophic Relationships of Demersal Fishes in the Shrimping Zone Off Alvarado Lagoon, Veracruz, Mexico

## Authors

Edgar Pelaez-Rodriguez, *University of Southern Mississippi*; Jonathan Franco-Lopez, *Universidad Nacional Autonoma de Mexico*; Wilfredo A. Matamoros, *University of Southern Mississippi*; Rafael Chavez-Lopez, *Universidad Nacional Autonoma de Mexico*; and [Nancy J. Brown-Peterson](#), *University of Southern Mississippi*

## TROPHIC RELATIONSHIPS OF DEMERSAL FISHES IN THE SHRIMPING ZONE OFF ALVARADO LAGOON, VERACRUZ, MEXICO

Edgar Peláez-Rodríguez, Jonathan Franco-López, Wilfredo A. Matamoras<sup>1\*</sup>, Rafael Chavez-López, and Nancy J. Brown-Peterson<sup>2</sup>

Laboratorio de Ecología, Facultad de Estudios Superiores Iztacala, Universidad Nacional Autónoma de México, Av. De los Barrios No 1, Los Reyes Iztacala, Tlalnepantla, México C.P. 54090 A.P. México

<sup>1\*</sup>Corresponding author: Department of Biological Sciences, The University of Southern Mississippi, Box 5018, Hattiesburg, Mississippi 39406-5018, and <sup>2</sup>Department of Coastal Sciences, The University of Southern Mississippi, 703 East Beach Drive, Ocean Springs, Mississippi 39564 USA

**ABSTRACT** The diet of demersal piscivorous fishes captured as bycatch of the commercial shrimping fleet off the Alvarado lagoonal system, Veracruz, Mexico, was studied. Nine collections distributed throughout the nortes (windy), wet, and dry seasons were made from November 1993 to January 1995. Sampling yielded a total of 646 fishes representing 10 families and 14 species, of which 44.9% had empty digestive tracts and were excluded from analysis. *Trichiurus lepturus* and *Synodus foetens* were the most abundant demersal predators in the collections. Differences in food consumption of the 7 most abundant predators were observed among the 3 seasons, with the greatest variety of prey (20 species) taken during the nortes season and the lowest variety (9 species) during the dry season. Five distinct trophic guilds were determined based on an index of relative importance of prey. Prey type and location of prey within the water column helped determine guild classification. The occurrence of different trophic guilds may allow for decreased competition for food resources on the continental shelf off Alvarado, Mexico.

**RESUMEN** Se estudio la dieta de los peces piscívoros demersales capturados como fauna acompañante del camarón en la flota de barcos camaroneros del sistema de lagunas de Alvarado, Veracruz, México. Se obtuvieron nueve colecciones que abarcaron las temporadas de nortes, lluvias y secas desde noviembre de 1993 hasta enero de 1995. Las muestras produjeron un total de 646 peces representados por 10 familias y 14 especies; 44.9% fueron encontrados con el tracto estomacal vacío y no fueron analizados. *Trichiurus lepturus* y *Synodus foetens* fueron los depredadores demersales más abundantes en nuestras muestras. Se observaron diferencias en el consumo de alimento en las tres temporadas. La temporada de nortes mostró la mayor variación de presa (20 especies), y la menor variación se observó en la temporada de secas (9 especies). Cinco distintivos gremios tróficos fueron identificados basados en el índice de importancia relativa de la presa. El tipo de presa y la localización de las presas en la columna de agua permitieron determinar la clasificación de los gremios. La existencia de diferentes gremios tróficos permite una disminución en la competencia por recursos alimenticios en la plataforma continental del Alvarado, México.

### INTRODUCTION

Shrimp trawling is one of the most important fishing industries in Mexico. In the southern Gulf of Mexico off Veracruz, a serious decline in the Mexican shrimping industry was observed from 1980 to 1991. In 1980, the shrimping industry reported a production of 5000 metric tons/year of penaeid shrimp (Grande and Díaz 1981), whereas in 1991 production using the same capture effort was only 1500 metric tons/year (SEMARNAP 1997). Currently, catches oscillate between 2000 and 3000 metric tons/year off Veracruz (Uribe-Marinez 2003). Worldwide, overfishing both by commercial and recreational fishers has reduced the abundance and biomass of apex predator species (Tegner and Dayton 1999, Jackson et al. 2001, Coleman et al. 2004) as well as non-targeted species (Burrage et al. 1993, Steele et al. 2001), leading to altered food webs in estuaries, coral reefs, and kelp forests (Jackson et al. 2001).

Data from several localities of the world show that in some types of fisheries more than 90% of the total catch (biomass) is discarded as waste bycatch (Alverson et al. 1994, Erzini et al. 2001, Kennelly and Broadhurst 2002). Studies have shown that the fish to shrimp ratio in temperate and subtropical areas of Mexico is 5:1 metric tons/yr, while the ratio in tropical areas is 10:1 metric tons/yr (Grande and Díaz 1981). Furthermore, shrimp trawling disturbs extensive areas of benthic habitat, affects the benthic macrofauna, and dramatically changes the diversity and abundance of demersal fish fauna (Alverson et al. 1994, Kaiser 1998, Rogers et al. 1999).

Little is known about the trophic structure and other ecological processes of the biotic community in the shrimping area off the Alvarado Lagoon, Veracruz, Mexico. This study was designed to examine the abundance and trophic interactions of demersal predatory fishes that are part of the bycatch in this area of high shrimp trawling effort. A common method of establishing trophic

structure is by the determination of trophic guilds (Luczkovich et al. 2002). Trophic guilds, defined as the grouping of species that share similar resources in a competitive complex (Root 1973, Blondel 2003), were determined in this study through analysis of stomach contents of trawl-caught fishes.

## METHODS

### Study Area

The study area is located immediately offshore of the Alvarado Lagoon system in the central portion of the state of Veracruz, Mexico, between 18°45'N, 95°40'W and 19°00'N, 95°42'W. Three well defined seasons characterize the region: the wet season from June through September, the nortes (windy) season from October through January, and the dry season from February through May (Contreras 1985). Highest precipitation occurs during the rainy season and oscillates between 1100–2000 mm over the year (García 1973). The Alvarado area is characterized by extensive coastal vegetation including mangroves and seagrasses and a series of lagoons and rivers that brings considerable fresh water and organic matter to the continental shelf, particularly during the rainy season.

### Sample collection and processing

We collected demersal fishes, known from the literature to be piscivores, from boats of the Alvarado shrimping fleet on 9 occasions from November 1993–January 1995, covering all 3 seasons. There were 4 collections during the nortes season, 3 collections during the wet season and 2 collections during the dry season. Boats in the fleet were equipped with a 20 m beam trawl with a 5.5 m mouth opening that was constructed with 3.85 cm mesh. Towing speed was 5–6 km/h, covering a distance of 1.8–18.5 km per sampling event. Fishing depths ranged from 30–90 m, with a mean depth of 50 m. A 30 l subsample of the bycatch (representing 25–27 kg of fish) was obtained using the methods described by Guzmán (1991) and Peláez-Rodríguez (1993) from trawls fished for 4 h between 0800–0730 local time (Central Time Zone).

Formaldehyde (10%) was injected into the oral and anal areas and then fish were immersed in the formaldehyde solution (Laevastu 1971). Fishes were labeled, bagged, and transported to the laboratory where samples were rinsed with tap water and preserved in 70% methanol within 48 to 72 hours. Species were identified with Hoese and Moore (1977), Fisher (1978), and Castro-Aguirre (1978). Fish were measured (standard length, SL, mm) and weighed to the nearest 0.1 g. Stomachs were extracted, and

their contents were identified to the lowest possible taxon using hard parts such as otoliths, scales, jaw bones and cranial bones (Windell and Stephen 1978). Prey items were blotted with desiccant paper and weighed to the nearest 0.001 g; empty stomachs were noted but not included in the analysis. Stomach contents of the 7 most abundant predators captured were used for analysis. Prey items were classified as pelagic, benthic, or benthic-pelagic according to knowledge of their general occurrence within the water column (Carpenter 2002).

### Data analysis

Abundance and biomass of the predator species were compared among seasons for each subsample with analysis of variance (ANOVA) and pairwise Sidak post-hoc tests to separate mean values if a significant F-test was determined. Species richness (S) was determined seasonally based on the abundance of the demersal, predatory fishes captured. Additionally, percent contribution of each species in terms of abundance and biomass were calculated by season.

The importance of each prey species for each of the 7 most abundant fishes was evaluated by pooling data for each season and then calculating the index of relative importance (IRI; Pinkas et al. 1971), defined as  $IRI = \%F(\%N + \%W)$ , where  $\%F$  = frequency of occurrence of a food item,  $\%N$  = numerical percentage of a food item in the stomachs, and  $\%W$  = percentage by volume of the food item in the stomachs (Pinkas et al. 1971). IRI values were standardized to  $\%IRI$  for comparison (Cortés 1997).

A Bray-Curtis dissimilarity matrix was calculated based on  $\%IRI$  values, and this matrix was used to construct a dendrogram using the unpaired grouping mean average (UPGMA) method (Field et al. 1982). ANOVA was calculated using SPSS (SPSS Inc, ver 11.5, Chicago, IL). Values were considered significantly different if  $P < 0.05$ .

## RESULTS

### Predator abundance and seasonality

Fourteen species of demersal fishes belonging to 10 families were collected during the study, yielding a total of 646 individuals with a total biomass of 54 kg (Table 1). The families Synodontidae (4 species) and Sciaenidae (2 species) contributed almost half of the total species. Of the total catch, only 362 fishes or 56.1% contained prey in their stomachs. Three species have not been previously reported for the Alvarado area; they include *Rachycentron canadum*, collected only during the nortes season, and *Synodus poeyi* and *Trachinocephalus myops*, reported for both the nortes and wet seasons (Table 1). Overall,

TABLE 1

Composition of the demersal fish fauna collected from commercial shrimp nets off the Alvarado Lagoon system during the nortes, dry and wet seasons. Abbreviations are presented for the 7 most abundant species.

Species	Nortes		Dry		Wet		Total	
	Abundance (ind)	Biomass (g)	Abundance (ind)	Biomass (g)	Abundance (ind)	Biomass (g)	Abundance (ind)	Biomass (g)
Muraenidae								
<i>Gymnothorax nigromarginatus</i>	5	685.3	7	635.9	6	709.1	18	2030.3
Ophichthidae								
<i>Myrophis punctatus</i>	2	147.0	3	192.5	6	334.7	11	674.2
Synodontidae								
<i>Synodus foetens</i> (Syfo)	67	14811.7	32	3119.3	25	1761.0	124	19692.0
<i>Synodus poeyi</i>	25	868.4			32	717.2	57	1585.6
<i>Trachinocephalus myops</i>	8	369.4			15	849.8	23	1219.2
<i>Saurida brasiliensis</i> (Sabr)	15	75.7	5	47.9	66	291.5	86	415.1
Fistulariidae								
<i>Fistularia tabacaria</i>	2	84.1			2	42.1	4	126.2
Priacanthidae								
<i>Priacanthus arenatus</i>	15	1518.6	2	373.3	6	568.7	23	2460.6
Rachycentridae								
<i>Rachycentron canadum</i>	2	1208.0					2	1208.0
Sciaenidae								
<i>Cynoscion arenarius</i> (Cyar)	5	639.3	8	560.9	6	1100.6	19	2300.8
<i>Cynoscion nothus</i> (Cyno)	25	2316.5	13	619.5	22	1853.7	60	4789.7
Sphyraenidae								
<i>Sphyraena guachancho</i> (Spgu)	5	586.0	10	445.5	13	1612.7	28	2644.2
Trichiuridae								
<i>Trichiurus lepturus</i> (Trle)	43	4154.0	19	1514.9	87	6700.5	149	12369.4
Scombridae								
<i>Scomberomorus cavalla</i> (Scca)	31	540.0	6	1449.2	5	315.2	42	2304.4
Totals	250	28004.0	174	8958.9	291	16856.8	646	53819.7
Species collected	14		10		13		14	

*Trichiurus lepturus* was the most common predator species captured during the study, with a total of 149 individuals, and was the dominant species during the wet season. *Synodus foetens* and *Saurida brasiliensis* were the second and third most abundant predatory fishes captured, while *Cynoscion nothus*, *Scomberomorus cavalla*, *Sphyraena guachancho* and *C. arenarius* rounded out the top 7 species (Table 1).

The wet season showed the highest abundance of predatory fishes in the shrimp bycatch, but it ranked second in biomass, with 291 specimens and 17 kg. The nortes season accumulated the highest biomass of bycatch predators, 28 kg, but occupied the second place in predator fish abundance with 250 specimens. The lowest values of abundance and biomass were found during the dry season with a total of 174 specimens that yielded 9 kg (Table 1).

However, there were no significant differences among seasons for either abundance (ANOVA,  $F_{2,6} = 3.46$ ,  $P = 0.100$ ) or biomass (ANOVA,  $F_{2,6} = 0.95$ ,  $P = 0.438$ ), suggesting a relatively stable and constant bycatch of predatory fishes in the shrimp trawl fishery in the area.

Seasonally, richness of predatory bycatch fishes was greater in the nortes season followed by wet and then dry seasons (Table 1); a similar pattern was seen in total abundance as well. *Synodus foetens*, *T. lepturus* and *S. brasiliensis* were important contributors numerically and/or in terms of biomass to the total species complement (Tables 1 and 2). *Synodus foetens* was first and *T. lepturus* second in the nortes and dry seasons in terms of abundance and biomass. In the wet season *T. lepturus* and *S. brasiliensis* were the first and 2nd most abundant species, whereas *T. lepturus* and *C. nothus* contributed more to biomass (Table 2).

TABLE 2

Percent contribution of abundant predatory fishes by season in terms of abundance and biomass in the shrimping zone off the Alvarado Lagoon, Veracruz, Mexico.

Species	Nortes		Dry		Wet	
	Abundance	Biomass	Abundance	Biomass	Abundance	Biomass
<i>Synodus foetens</i>	26.80	52.89	30.48	34.82	8.59	10.45
<i>Trichiurus lepturus</i>	17.20	14.83	18.09	16.91	29.90	39.75
<i>Cynoscion nothus</i>	10.00	8.27	12.38	6.91	7.56	10.99
<i>Scomberomorus cavalla</i>	12.40	1.93	5.71	16.18	1.72	1.87
<i>Saurida brasiliensis</i>	6.00	0.27	4.76	0.53	22.68	1.73
<i>Sphyaena guachancho</i>	2.00	2.09	9.52	4.97	4.47	9.57
<i>Cynoscion arenarius</i>	2.00	2.28	7.62	6.26	2.06	6.53
<i>Synodus poeyi</i>	10.00	3.10			10.99	4.25
<i>Priacanthus arenatus</i>	6.00	5.42	1.90	4.17	2.06	3.37
<i>Gymnothorax nigromarginatus</i>	2.00	2.45	6.67	7.10	2.06	4.21
<i>Trachinocephalus myops</i>	3.20	1.32			5.15	5.04
<i>Myrophis punctatus</i>	0.80	0.52	2.86	2.15	2.06	1.99
<i>Fistularia tabacaria</i>	0.80	0.30			0.69	0.25
<i>Rachycentron canadum</i>	0.80	4.31				

**Predator size and diets**

The modal SL for *S. foetens* was smaller in the nortes season than in the dry or wet seasons, whereas modal SL for *T. lepturus* was largest during the wet season (Table 3). However, the range in sizes for these 2 species overlapped for all 3 seasons. The modal SL for *S. guachancho* was larger in the nortes season compared to the dry or wet seasons, and the size range for the nortes season did not overlap with the other 2 seasons (Table 3). The remaining 4 predator size ranges and modal SL did not change much by season (Table 3). This suggests that potential ontogenetic diet shifts imbedded within seasons probably did not affect analyses of trophic spectrum.

Twenty-four prey species, including 20 fishes, three decapod crustaceans, and one cephalopod, were identified

from the stomach contents of the top 7 predators. Among the fish prey, *Bregmaceros cantori* and *Microdesmus lanceolatus* have not been previously reported from the shelf off Alvarado Lagoon (Table 4). For the 7 predator species, the lowest number of prey types consumed (9) occurred during the dry season, while the highest number of prey types (20) was found during the nortes season. Prey types varied among predators and changed seasonally (Table 4).

*Synodus foetens* was the second most abundant predatory species overall and had the largest variety of prey, with a total of 17 taxa (Table 4). This species fed on the greatest diversity of prey during the nortes season, and its prey occurred throughout the water column (Figure 1). Fifty-one percent IRI of the prey was benthic and included

TABLE 3

Summary statistics on fish standard length (range and mode, cm) by season for the 7 predators used in the diet analysis.

Species	Nortes		Dry		Wet	
	Range	Mode	Range	Mode	Range	Mode
<i>Sphyaena guachancho</i>	29.3–34.2	30.0	16.5–19.1	18.0	18.0–20.8	19.0
<i>Synodus foetens</i>	12.3–21.7	18.0	20.6–35.7	29.0	17.6–43.5	32.0
<i>Trichiurus lepturus</i>	32.0–58.6	46.0	39.4–51.8	47.0	45.6–89.7	64.0
<i>Cynoscion arenarius</i>	18.5–23.4	20.0	17.5–19.40	18.0	18.5–24.6	21.0
<i>Cynoscion nothus</i>	15.6–19.0	17.0	14.2–18.0	16.0	16.2–21.6	19.0
<i>Saurida brasiliensis</i>	7.4–8.9	8.0	5.2–7.6	6.0	8.4–11.0	9.0
<i>Scomberomorus cavalla</i>	24.5–27.3	25.0	22.6–28.4	24.0	21.5–27.6	25.0

TABLE 4

Seasonal food composition and %IRI for 7 demersal fishes off Alvarado, Veracruz.

Species	Nortes		Dry		Wet	
	Prey type	%IRI	Prey type	%IRI	Prey type	%IRI
<i>S. guachancho</i>	<i>Anchoa hepsetus</i>	68.44	<i>Bregmaceros cantori</i>	66.15	<i>Anchoa hepsetus</i>	22.02
	<i>Cynoscion nothus</i>	1.82	<i>Saurida brasiliensis</i>	8.56	<i>Saurida brasiliensis</i>	57.6
	<i>Bregmaceros cantori</i>	8.94	<i>Loligo pealei</i>	25.29	<i>Loligo pealei</i>	20.38
	<i>Saurida brasiliensis</i>	15.34				
	<i>Loligo pealei</i>	5.46				
<i>S. foetens</i>	<i>Anchoa hepsetus</i>	21.17	<i>Anchoa hepsetus</i>	51.26	<i>Saurida brasiliensis</i>	19.36
	<i>Saurida brasiliensis</i>	2.03	<i>Upeneus parvus</i>	48.74	<i>Upeneus parvus</i>	8.29
	<i>Upeneus parvus</i>	12.92			<i>Loligo pealei</i>	19.33
	<i>Loligo pealei</i>	7.44			<i>Bregmaceros cantori</i>	17.70
	<i>Harengula clupeiola</i>	14.55			<i>Pristipomoides aquilonaris</i>	1.76
	<i>Trachurus lathami</i>	2.64			<i>Diplectrum bivittatum</i>	23.17
	<i>Micropogonias furnieri</i>	2.99			<i>Syacium gunteri</i>	3.57
	<i>Pristipomoides aquilonaris</i>	9.16			<i>Trichiurus lepturus</i>	6.24
	<i>Diplectrum bivittatum</i>	8.57			<i>Engyophrys senta</i>	0.56
	<i>Symphurus plagiusa</i>	1.46				
	<i>Haemulon aurolineatum</i>	3.14				
	<i>Serranus atrobranchus</i>	13.01				
	<i>T. lepturus</i>	<i>Eucinostomus gula</i>	0.90			
<i>Anchoa hepsetus</i>		49.48	<i>Upeneus parvus</i>	33.28	<i>Anchoa hepsetus</i>	36.43
<i>Upeneus parvus</i>		15.11	<i>Harengula clupeiola</i>	24.01	<i>Upeneus parvus</i>	1.19
<i>Pristipomoides aquilonaris</i>		13.72	<i>Loligo pealei</i>	42.71	<i>Pristipomoides aquilonaris</i>	1.49
<i>Harengula jaguana</i>		12.2			<i>Diplectrum bivittatum</i>	0.55
<i>Harengula clupeiola</i>		5.85			<i>Synodus foetens</i>	0.23
<i>Loligo pealei</i>		2.43			<i>Bregmaceros cantori</i>	2.54
<i>Farfantepenaeus</i> sp.		1.21			<i>Saurida brasiliensis</i>	8.81
<i>C. arenarius</i>					<i>Cynoscion nothus</i>	0.38
	<i>Saurida brasiliensis</i>	27.05	<i>Upeneus parvus</i>	76.21	<i>Myrophis punctatus</i>	7.19
	<i>Upeneus parvus</i>	35.47	<i>Diplectrum bivittatum</i>	23.79	<i>Loligo pealei</i>	8.27
	<i>Pristipomoides aquilonaris</i>	15.21			<i>Farfantepenaeus</i> sp.	32.89
	<i>Loligo pealei</i>	0.65			<i>Saurida brasiliensis</i>	33.96
<i>C. nothus</i>	<i>Farfantepenaeus</i> sp.	21.62			<i>Upeneus parvus</i>	24.11
	<i>Pristipomoides aquilonaris</i>	37.91	<i>Bregmaceros cantori</i>	95.49	<i>Loligo pealei</i>	41.93
	<i>Bregmaceros cantori</i>	31.92	<i>Farfantepenaeus</i> sp.	4.51		
	<i>Saurida brasiliensis</i>	13.15			<i>Saurida brasiliensis</i>	20.41
	<i>Trichiurus lepturus</i>	4.16			<i>Trichiurus lepturus</i>	6.05
	<i>Microdesmus lanceolatus</i>	0.15			<i>Farfantepenaeus</i> sp.	21.94
<i>S. brasiliensis</i>	<i>Loligo pealei</i>	12.70				
	<i>Bregmaceros cantori</i>	78.42	<i>Bregmaceros cantori</i>	67.62	<i>Bregmaceros cantori</i>	75.16
	<i>Loligo pealei</i>	21.58	<i>Loligo pealei</i>	32.38	<i>Loligo pealei</i>	24.84
<i>S. cavalla</i>	<i>Anchoa hepsetus</i>	94.08	<i>Anchoa hepsetus</i>	69.67	<i>Anchoa hepsetus</i>	25.52
	<i>Bregmaceros cantori</i>	5.92	<i>Upeneus parvus</i>	30.33	<i>Diplectrum bivittatum</i>	53.52
				<i>Loligo pealei</i>	20.95	




Figure 1. Percentage of prey occurring in the pelagic, benthic-pelagic, and benthic zones of the water column for 7 demersal fishes off the Alvarado Lagoon system, Veracruz, Mexico, during the nortes, dry, and wet seasons. *Synodus foetens* (Syfo), *Saurida brasiliensis* (Sabr), *Cynoscion arenarius* (Cyar), *Cynoscion nothus* (Cyno), *Sphyraena guachancho* (Spgu), *Trichiurus lepturus* (Trle), *Scomberomorus cavalla* (Scca). Sample size for each species in the figure by season is found in Table 1.



*Upeneus parvus*, *Diplectrum bivittatum*, *Pristipomoides aquilonaris*, *Eucinostomus gula*, *Micropogonias furnieri*, *Symphurus plagiusa*, and *S. brasiliensis*. Species from the pelagic zone contributed 41.5 %IRI of stomach contents and included *Anchoa hepsetus*, *Harengula clupeiola*, and *Trachurus lathami*. A smaller percentage of the diet, 10.5 %IRI, was composed of *Loligo pealei* and *Haemulon aurolineatum* from the benthic-pelagic zone. During the dry season, *S. foetens* fed about equally on *A. hepsetus* from the pelagic zone and on *U. parvus*, a bottom dweller. During the wet season, the diet of *S. foetens* was dominated by benthic prey (80.6 %IRI) which included *S. brasiliensis*, *D. bivittatum*, *T. lepturus*. *U. parvus*, *P. aquilonaris*, *Engyophrys senta*, *Syacium gunteri*, and *B. cantori*. The benthic-pelagic zone contributed 19.3 %IRI to the diet; the only prey was *L. pealei* (Table 4).

*Saurida brasiliensis* was the smallest piscivorous predator in the study and showed no differences in seasonal prey consumption (Table 4, Figure 1). This species also had the least diverse diet, with the benthic *B. cantori* accounting for 67–78 %IRI of the diet each season. A benthic-pelagic species, *L. pealei*, made up the rest of the diet (Table 4).

Neither *Cynoscion arenarius* nor *C. nothus* consumed any pelagic prey during the course of this study (Figure 1). Both species had the greatest diversity of prey items during the nortes season. During both the nortes and dry seasons, >80 %IRI of the diet was composed of benthic species such as *U. parvus*, *Farfantepenaeus* sp., *P. aquilonaris*, *B. cantori*, and *S. brasiliensis*, whereas the remaining diet was composed of the benthic-pelagic *L. pealei* (Table 4). The diet of *C. arenarius* was dominated by benthic species during the wet season, as was the diet of the congener *C. nothus* (Figure 1). While the 2 *Cynoscion* species fed within the same areas of the water column, there were differences in the prey they captured. For instance, *B. cantori* was an important component of the diet of *C. nothus* throughout the year, yet this prey was never eaten by *C. arenarius* (Table 4). Similarly, *U. parvus* dominated the diet of *C. arenarius* but was never taken by *C. nothus* (Table 4).

*Sphyraena guachancho* consumed only 5 prey items, yet there was marked seasonal variation in the dominant prey items (Table 4). For instance, the pelagic *A. hepsetus* dominated the diet in the nortes season, while no pelagic species were consumed during the dry season when the benthic *B. cantori* dominated the diet (Table 4, Figure 1). During the wet season, benthic prey such as *S. brasiliensis* was dominant in the diet.

*Trichiurus lepturus* was the most abundant predator species captured during the study and the only species to feed throughout the water column year round (Figure 1).


During the wet season, benthic (48 %IRI) and pelagic (36.4 %IRI) species constituted the majority of the diet (13 species) of *T. lepturus*; prominent taxa included *Farfantepenaeus* sp., *S. brasiliensis*, and *A. hepsetus* (Table 4). In contrast, the pelagic species *A. hepsetus*, *Harengula jaguana*, and *H. clupeiola* dominated the diet during the nortes season (67.5 %IRI). During the dry season, *T. lepturus* fed on 3 prey species, one from each section of the water column. The benthic-pelagic *L. pealei* (42.7 %IRI) dominated the diet (Table 4, Figure 1).

*Scomberomorus cavalla* consumed only 5 prey types during the course of the study. The pelagic *A. hepsetus* dominated the diet during both the nortes (94.1 %IRI) and dry (69.7 %IRI) seasons and also accounted for 25 %IRI of the diet during the rainy season (Table 4). While benthic prey were taken throughout the year and dominated the diet in the rainy season (Figure 1), *S. cavalla* fed on different benthic species during each season (Table 4).

#### Species/season dietary patterns

Five distinct trophic guilds were delimited (Figure 2). Fishes in feeding guild A consumed mainly pelagic prey like *A. hepsetus*, *H. jaguana*, and *H. clupeiola*, whereas fish in guild B consumed not only pelagic species but transitioned to feeding on benthic-pelagic species like *L. pealei* (Figure 2). Fishes in guild C were characterized by feeding on a mixture of benthic-pelagic and benthic prey like *Farfantepenaeus* sp., *S. brasiliensis*, *U. parvus*, *Myrophis punctatus*, and *L. pealei* (Figure 2). Fishes in feeding guilds D and E tended to focus on more benthic prey like *S. brasiliensis*, *Farfantepenaeus* sp., *U. parvus*, and *B. cantori*.

In general, the species/season trophic patterns identified by guild analysis did not follow clear patterns, most likely due to body size-mouth gape differences and to seasonal prey availability. For example, *C. nothus*, *S. brasiliensis* and *C. arenarius* exhibited no seasonal differences in trophic guild, and *C. nothus* and *C. arenarius* were assigned to different guilds (Figure 2). This suggests minimal differences in prey across seasons for these species. In contrast, members of guilds A and B were comprised of different species and seasons with no clear patterns (Figure 2). Some species/season diets clustered together, and others did not. It was clear, however, that some species shifted from pelagic to benthic prey with season. For example, *S. guachancho* fed on pelagic species during the nortes and wet seasons but shifted to benthic prey during the dry season. However, the modal SL and size ranges for *S. guachancho* were virtually identical during the dry and wet seasons (Table 3), suggesting the seasonal shift in prey is not related to ontogenic feeding dif-



**Figure 2.** UPGMA cluster analysis of %IRI based on Bray Curtis dissimilarity index for 7 demersal fishes off Alvarado Lagoon Veracruz, Mexico. *Sphyraena guachancho* (Spgu), *Synodus foetens* (Syfo), *Trichiurus lepturus* (Trle), *Scomberomorus cavalla* (Scca), *Cynoscion arenarius* (Cyar), *Cynoscion nothus* (Cyno), *Saurida brasiliensis* (Sabr). N = nortes season, W = wet season, and D = dry season. Letters indicate trophic guilds identified from the cluster analysis.

ferences. Similarly, *S. foetens* and *T. lepturus* were found in 3 different guilds based on season, suggesting differences in size may not be as important as other factors determining prey selection. *Synodus foetens* fed on pelagic, benthic-pelagic, and benthic prey during all seasons, and *T. lepturus* fed on pelagic and benthic-pelagic prey. These 2 species were the most abundant species examined during this study and contributed the highest portion of biomass.

### DISCUSSION

Stomach content analysis is used widely to determine food composition, feeding strategies, trophic position, energy flow of predator and prey (Hyslop 1980), trophic structure (Luczkovich et al. 2002), and trophic partitioning (Ross 1986). Our analysis indicates the examination of stomach contents of top carnivores is an excellent way to evaluate the relationship between predators and food source in the shrimp grounds of Veracruz, Mexico.

The diets reported here for the 7 most abundant predators are generally similar to previous reports (Naughton and Saloman 1981, Mericas 1981, Divita et al. 1983,

Sheridan et al. 1984, Cruz-Escalona et al. 2005), with some notable exceptions. While fish (in particular *Anchoa*) were important in the diet of *T. lepturus* in both this study and in the northern Gulf of Mexico (GOM) (Mericas 1981, Sheridan et al. 1984), the seasonal dominance of squid in the diet (42.7 %IRI during the dry season) has not been previously reported. The diets of both *C. arenarius* and *C. nothus* captured off Veracruz differed from previous reports for the species (Sheridan et al. 1984, Sutter and McIlwain 1987) in that no pelagic prey were noted in the present study, while *Anchoa* was a major component of the diet of both species in the northern GOM (Sheridan et al. 1984). Furthermore, *Bregmoceros*, common in the diets of other predators captured in the present study, was not found in either *Cynoscion* species, although this prey species was previously reported as an important component of the diet (Sheridan et. al 1984). While the diet of *S. brasiliensis* was dominated by fish as expected, squid was a more important component of the diet of (21.5–31.3 %IRI) than the 9% frequency of occurrence previously reported by Divita et al. (1983). The predominantly piscivorous diet of *S. foetens* agrees with previous reports from the northern GOM (Divita et al. 1983) and the Veracruz,

Mexico, area (Cruz-Escalona et al. 2005), although the complete absence of penaeid shrimp in the diet is in contrast to reports from the northern GOM (Divita et al. 1983).

Our results showed patterns of resource partitioning and indicated that the 7 most abundant species examined in our study had proportioned diets based on where in the water column their prey was found. This tendency towards resource partitioning coincides with findings by Abarca-Arenas et al. (2004), who found similar evidence of resource partitioning in the Alvarado area based on the entire fish community. Macpherson (1981) and Livingston (1982) stated that in a trophic system, resource partitioning always will be observed; the pattern can be observed at the temporal level or in some cases at the diel level, even when competition among species exists

Five trophic guilds were clearly identified in our study based on the level of the water column in which the prey was obtained. Two guilds fed mainly on pelagic prey, 2 fed more on benthic prey, and one fed more on benthic-pelagic prey. Noteworthy is the large number of prey items consumed by the latter guild, demonstrating capacity to feed throughout the water column and to maintain generalist prey consumption habits.

Formation of the guilds did not appear related to body size, but rather to prey availability. Based on stomach contents, prey selection varied among the 3 seasons. Sedberry (1983) studied a community of demersal fishes on the continental shelf of the middle Atlantic Bight and also documented seasonal prey-shifting that appeared to be independent of predator size. The dry season showed the fewest taxa of prey taken, and the nortes season showed the most. With predator abundance remaining constant year-round and prey sources varying, guild structure was most likely affected. Although measurements were not made of abundance and diversity of prey beyond those obtained via stomach contents, our results suggest that prey in the nortes and wet seasons are more diverse than prey in the dry season, thus affecting the trophic guilds (sensu Darnell 1961).

Seasonal nutrient flux may influence prey availability in the study area and thus the structure of trophic guilds. Nutrients in the Alvarado Lagoon system are largely dependent upon influx from the Papaloapan River. The river deposits the largest amount of nutrients into the system during the wet and nortes seasons (Moran-Silva et al. 2005), resulting in higher productivity levels (Abarca-Arenas et al. 2004) and a general increase in the amount of exploitable resources in the system (Contreras 1985, Soberón and Yañez-Arancibia 1985). Thus, it was not a surprise that our study found the largest variety of prey and the highest abundance of predators during these 2 seasons.

Anthropogenic factors can affect the guild structure as well. Shrimp trawling is an important commercial activity off Alvarado (Grande and Díaz 1981). The effects of bycatch removal on the local demersal fish community have not been measured; however, evidence suggests that large-scale fishing affects the structure of fish communities by reducing the abundance of prey and predators and by reducing the size of predators (Pope and Knights 1982, Rice and Gislason 1996, Jennings et al. 1998, Rogers et al. 1999). In the Alvarado area, information is lacking regarding fishing activities and the life history and ecology of piscivorous fishes and their prey; thus, it is difficult to estimate the effect of the shrimp fishery and its bycatch on the trophic dynamics of the area. However, intense fishing activity in tropical waters can cause reduction in species richness and dominance of the smaller targeted and non-targeted fishes in the assemblage (Rogers et al. 1999). Our data suggest that a similar reduction in larger species may have occurred near Alvarado. For instance, large, potentially commercially important species such as *R. canadum*, *F. tabacaria*, *S. guachancho*, and *S. cavalla* composed only 11% of the total bycatch. Dominance of *S. foetens* and *T. lepturus*, 2 non-target species with the greatest variety of prey, suggests trophic adaptability and generalization may be important in this heavily fished system.

#### ACKNOWLEDGMENTS

This manuscript is based on the senior thesis of the lead author submitted in 1996 in pursuit of the biologist degree to the Facultad de Estudios Superiores at the Iztacala campus of the Universidad Nacional Autónoma de México. We thank personnel from the Cet-Mar Academy in Alvarado, Mexico, for assistance with field collections, particularly Tomas Corro-Ferreiro. In particular, we thank M.S. Peterson for assistance and advice during the preparation of this manuscript.

#### LITERATURE CITED

- Abarca-Arenas, L.G., J. Franco-Lopez, R. Chavez-Lopez, D. Arceo-Carranza, and A. Moran-Silva. 2002. Trophic analysis of the fish community taken as bycatch of the shrimp trawls off the coast of Alvarado, Mexico. *Proceedings of the Gulf and Caribbean Fisheries Institute* 55:384–394.
- Alverson, D.L., M.H. Freeberg, S.A. Murawski, and J.G. Pope. 1994. A global assessment of fisheries by-catch and discards. *FAO Fisheries Technical Paper* 339:1–233.
- Blondel, J. 2003. Guilds or functional groups: does it matter? *Oikos* 100:223–231.
- Burrage, D.D., T.J. Schultz, J.J. Ross, and P. Anglada. 1993. Bycatch reduction in the northern Gulf shrimp fishery. Final report, Gulf and South Atlantic Fisheries Development Foundation, Tampa, FL, USA, 24 p.

- Carpenter, K.E. (ed). 2002. The Living Marine Resources of the Western Central Atlantic. Vols. 2 and 3. FAO, Rome, Italy. 1500 p.
- Castro-Aguirre, J.L. 1978. Catálogo sistemático de los peces marinos que penetran a las aguas continentales de México con aspectos zoogeográficos y ecológicos. Dirección General del Instituto Nacional de Pesca. México. Serie Científica 19. 298 p.
- Coleman, F.C., W.F. Figueira, J.S. Ueland, and L.B. Crowder. 2004. The impact of United States recreational fisheries on marine fish populations. *Science* 305:1958–1960.
- Contreras, F. 1985. Las lagunas costeras mexicanas. Centro de Ecodesarrollo, Secretaría de Pesca, México, D.F., 253 p.
- Cortés, E. 1997. A critical review of methods of studying fish feeding based on analysis of stomach contents: Application to elasmobranch fishes. *Canadian Journal of Fisheries and Aquatic Sciences* 54:726–738.
- Cruz-Escalona, V.H., M.S. Peterson, L. Campos-Dávila, and M. Zetina-Rejón. 2005. Feeding habits and trophic morphology of inshore lizardfish (*Synodus foetens*) on the central continental shelf off Veracruz, Gulf of Mexico. *Journal of Applied Ichthyology* In press.
- Darnell, R.M. 1961. Trophic spectrum of an estuarine community, based on studies of Lake Pontchartrain, Louisiana. *Ecology* 42:553–568.
- Divita, R., M. Creel, and P.F. Sheridan. 1983. Foods of coastal fishes during brown shrimp, *Penaeus aztecus*, migration from Texas estuaries (June–July 1981). *Fishery Bulletin, US* 81:396–404.
- Erzini, K., M.E. Costa, L. Bentes, and T.C. Borges. 2002. A comparative study of the species composition of discards from five fisheries from the Algarve (southern Portugal). *Fisheries Management and Ecology* 9:31–40.
- Field, J.G., K.R. Clarke, and R.M. Warwick. 1982. A practical strategy for analyzing multispecies distribution patterns. *Marine Ecology Progress Series* 8:37–52.
- Fischer, W., ed. 1978. Species Identification Sheets for Fishery Purposes. Western Central Atlantic (fishing area 31). Vols. I–VII. FAO, Rome, Italy.
- García, E. 1973. Modificación del Sistema de Clasificación Climática de Koopen. Instituto de Geografía. Universidad Nacional Autónoma México, Ciudad de México, México.
- Grande, V.J. and M.L. Díaz. 1981. Situación actual y perspectivas de utilización de la fauna de acompañamiento del camarón en México. *Ciencia Pesquera* 2:43–55.
- Guzmán, P.J. 1991. Ictiofauna acompañante en zonas de pesca comercial del camarón en Alvarado, Ver. Período 1989–1991. Tesis Professional, Facultad de Estudios Superiores-Iztacala, Universidad Nacional Autónoma de México, Tlalnepantla, México, 54 p.
- Hoese H.D. and R.H. Moore. 1977. Fishes of the Gulf of Mexico, Texas, Louisiana and Adjacent Waters. Texas A&M University Press, College Station, TX, USA, 327 p.
- Hyslop, E. J. 1980. Stomach contents analysis—a review of methods and their application. *Journal of Fish Biology* 17:411–429.
- Jackson, J.B.C., M.X. Kirby, W.H. Berger, K.A. Bjorndal, L.W. Botsford, B.J. Bourque, R.H. Bradbury, R. Cooke, J. Erlandson, J.A. Estes, T.P. Hughes, S. Kidwell, C.B. Lange, H.S. Lenihan, J.M. Pandolfi, C.H. Peterson, R.S. Steneck, M.J. Tegner, and R.R. Warner. 2001. Historical overfishing and the recent collapse of coastal ecosystems. *Science* 293:629–637.
- Jennings, S., J.D. Reynolds, and S.D. Mills. 1998. Life history correlates of responses to fisheries exploitation. *Proceeding of the Royal Society of London B* 265:333–339.
- Kaiser, M.J. 1998. Significance of bottom-fishing disturbance. *Conservation Biology* 12:1230–1235.
- Kennelly, S.J. and M. Broadhurst. 2002. By-catch begone: changes in the philosophy of fishing technology. *Fish and Fisheries* 3:340–355.
- Laevastu, T. 1971. Manual de métodos de biología pesquera. Acribia, FAO, Spain.
- Livingston, R.J. 1982. Trophic organization of fishes in a coastal seagrass system. *Marine Ecology Progress Series* 7:1–12.
- Luczkovich, J.J., G. P. Ward, J.C. Johnson, R.R. Christian, D. Baird, H. Neckles, and W.M. Rizzo. 2002. Determining the trophic guilds of fishes and macroinvertebrates in a seagrass food web. *Estuaries* 25:1143–1163.
- Macpherson, E. 1981. Resource partitioning in a Mediterranean demersal fish community. *Marine Ecology Progress Series* 4:183–193.
- Mericas, D. 1981. Feeding habits of the Atlantic cutlassfish, *Trichiurus lepturus* in the Gulf of Mexico. *Northeast Gulf Science* 4:137–140.
- Moran-Silva, A., L.A. Martínez Franco, R. Chávez-López, J. Franco-López, C. M. Bedia Sánchez, F. Contreras Espinosa, F. Gutiérrez Mendieta, N.J. Brown-Peterson, and M.S. Peterson. 2005. Seasonal and spatial patterns in salinity, nutrients and chlorophyll *a* in the Alvarado lagoonal system, Veracruz, Mexico. *Gulf and Caribbean Research* 17:133–143.
- Naughton, S.P. and C.H. Salomon. 1981. Stomach contents of juveniles of king mackerel (*Scomberomorus cavalla*) and Spanish mackerel (*S. maculatus*). *Northeast Gulf Science* 5(1):71–74.
- Peláez-Rodríguez, E. 1993. Prospección ecológica de la ictiofauna de acompañamiento de camarón en Alvarado, Ver. Memorias del XVII Simposio de Biologías de Campo Universidad Nacional Autónoma de México, Iztacala, México.
- Pinkas, L., M.S. Oliphant, and I.L.K. Iverson. 1971. Food habits of albacore, bluefin tuna and bonito in California waters. California Department of Fish and Game, Fish Bulletin 152:1–105.
- Pope, J.G. and B.J. Knights. 1982. Comparison of the length distribution of combined catches of all demersal fishes in surveys in the North Sea and at Faroe Bank. *Canadian Special Publication of Fisheries and Aquatic Sciences* 59:116–118.
- Rice, J. and H. Gislason. 1996. Patterns of change in the size spectra of numbers and diversity of the North Sea assemblage, as reflected in surveys and models. *ICES Journal of Marine Science* 53:1214–1225.
- Rogers, S.I., R.K. Clarke, and J.D. Reynolds. 1999. The taxonomic distinctness of coastal bottom-dwelling fish communities of the north-east Atlantic. *Journal of Animal Ecology* 68:769–782.

- Root, R.B. 1973. Organization of a plant-arthropod association in simple and diverse habitats: The fauna of collards (*Brassica oleracea*). *Ecological Monographs* 43:95–124.
- Ross, S.T. 1986. Resource partitioning in fish assemblages: A review of field studies. *Copeia* 1986:352–388.
- Sedberry, G.R. 1983. Food and trophic relationships of a community of fishes on the outer continental shelf. National Oceanographic and Atmospheric Administration Technical Report National Marine Fishery Service SSRF-773:1–56.
- SEMARNAP. 1997. Anuario Estadístico de Pesca. Secretaría del Medio Ambiente, Recursos Naturales y Pesca (SEMARNAP), México, 212 p.
- Sheridan, P.F., D.L. Trimm, and B.M. Baker. 1984. Reproduction and food habits of seven species of northern Gulf of Mexico fishes. *Contributions in Marine Science* 27:175–204.
- Soberón, C.G. and A. Yañez-Arancibia. 1985. Control ecológico de los peces demersales: variabilidad ambiental de la zona costera y su influencia en la producción natural de los recursos pesqueros. In: Yañez-Arancibia, ed. *Recursos pesqueros potenciales de México: La pesca acompañante del camarón*. Progreso Universidad de Alimentos. Instituto de Ciencias del Mar y Limnología, Instituto Nacional de Pesca, Universidad Nacional Autónoma de México, México, 748 p.
- Steele, P., T.M. Bert, K.H. Johnston, and S. Levett. 2001. Efficiency of bycatch devices in small otter trawls used in the Florida shrimp fishery. *Fishery Bulletin*, US 100:338–350.
- Sutter, F.C. and T.D. McIlwain. 1987. Species profiles: Life histories and environmental requirements of coastal fishes and invertebrates (Gulf of Mexico). Sand seatrout and silver seatrout. US Fish and Wildlife Service Biological Reports 82(11.72). US Army Corps of Engineers, TR EL-82-4, 15 p.
- Tegner, M.J. and P. K. Dayton. 1999. Ecosystem effects of fishing. *Trends in Ecology and Evolution* 14:261–262.
- Uribe-Marinez, J.A. 2003. Condición biológica pesquera actual del camarón rosado capturado por la flota del puerto de Campeche y propuesta de veda en 2002. *Memorias del III Foro de Camarón del Golfo de México y del Mar Caribe*. In: K.A.T. Wakida, S.R. Solana and M.J. Uribe, eds. *Secretaría de Agricultura, Ganadería: Desarrollo Rural, Pesca y Alimentación (SAGARPA), Campeche, México*, p. 43–46.
- Windell, J.T. and H.B. Stephen. 1978. Methods for study of fish diets based on analysis of stomach contents. In: T.D. Bagenal, ed. *Methods for assessment of fish production in fresh water*. I.B.P. Handbook No. 3, Blackwell Scientific Publications, Oxford, England, p. 219–226.
- Yañez-Arancibia, A. 1985. Recursos demersales de alta diversidad en las costas tropicales: Perspectiva ecológica. In: A. Yañez-Arancibia, ed. *Recursos pesqueros potenciales de México: La pesca del camarón*. Progreso Universidad de Alimentos. Instituto de Ciencias del Mar y Limnología, Instituto Nacional de Pesca, Universidad Nacional Autónoma de México, México, p. 17–38.