Gulf and Caribbean Research

Volume 19 | Issue 1

January 2007

Color Variation in the Caribbean Crab *Platypodiella spectabilis* (Herbst, 1794) (Decapoda, Brachyura, Xanthidae)

Joel W. Martin

Natural History Museum of Los Angeles County

Todd L. Zimmerman

Long Island University

Follow this and additional works at: <https://aquila.usm.edu/gcr>

Part of the [Marine Biology Commons](#)

Recommended Citation

Martin, J. W. and T. L. Zimmerman. 2007. Color Variation in the Caribbean Crab *Platypodiella spectabilis* (Herbst, 1794) (Decapoda, Brachyura, Xanthidae). *Gulf and Caribbean Research* 19 (1): 59-63.

Retrieved from <https://aquila.usm.edu/gcr/vol19/iss1/8>

DOI: <https://doi.org/10.18785/gcr.1901.08>

This Short Communication is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in *Gulf and Caribbean Research* by an authorized editor of The Aquila Digital Community. For more information, please contact Joshua.Cromwell@usm.edu.

COLOR VARIATION IN THE CARIBBEAN CRAB *PLATYPODIELLA* *SPECTABILIS* (HERBST, 1794) (DECAPODA, BRACHYURA, XANTHIDAE)

Joel W. Martin¹ and Todd L. Zimmerman²

¹Natural History Museum of Los Angeles County, 900 Exposition Boulevard, Los Angeles, California 90007 USA. E-mail jmartin@nhm.org

²Long Island University, C.W. Post Campus, 720 Northern Blvd, Brookville, New York 11548-1300 USA

INTRODUCTION

Platypodiella spectabilis (Herbst, 1794) is a relatively small crab (about 10 mm carapace width) found in or near coral reefs and rocky shorelines throughout most of the Caribbean and tropical western Atlantic. Records are known from as far north as Bermuda (Chace et al. 1986, in Sterrer 1986, as *Platypodia spectabilis*) and as far south as Rio de Janeiro, Brazil, including the Fernando de Noronha Archipelago and Trinidad Island (Melo 1998: 490) and the state of São Paulo (Coelho and Ramos 1972, Fransozo et al. 2001). Thus, the range of the species is some 7,000 km from northern to southern extent. Distributional records within this range, and extending westward into the Gulf of Mexico, can be found in Rathbun (1930), Felder (1973), Powers (1977), and Abele and Kim (1986). Because of its spectacular coloration and color pattern, *P. spectabilis* is often depicted in faunistic guide books (e.g., Humann 1992, Humann and DeLoach 2002), and the species is sometimes referred to as the calico crab (e.g., Chace et al. 1986) or gaudy clown crab (Williams et al. 1989, McLaughlin et al. 2005).

The use of color in brachyuran crab systematics, and in particular the use of subtle color differences to suggest or differentiate cryptic or morphologically similar species, is now well documented (e.g., see Campbell and Mahon 1974 for species of *Leptograpsus*, Williams and Felder 1989 for species of *Menippe*, Zimmerman and Felder 1991 for species of *Sesarma*). Less clearly understood is why color patterns and intensities can sometimes vary appreciably within a species, even within narrowly restricted geographic regions. An appreciation of color patterns is critical to correctly identifying species for conservation and resource management purposes, yet often color patterns and ranges are unreported, causing confusion and sometimes misidentifications. Here we document a wide range of color patterns in a small Caribbean xanthid crab based on specimens collected in essentially the same habitat at the same time of year.

MATERIALS AND METHODS

As part of a biodiversity survey of Caribbean crypto-faunal invertebrates, we sampled several habitats from shallow waters off Guana Island, British Virgin Islands, during the summers (June–August) of 1999–2002. Specimens of *P. spectabilis* collected during that survey came almost exclusively from an area of a few square meters at our North Bay collecting site, where they were found in interstices of dead coral (mostly clumps of dead *Porites*) in shallow water (≤ 1 m) (Station 65 of the Zimmerman/Martin survey of Guana Island in 2000 and 2001). Collections were made by crushing clumps of dead coral and removing by hand the invertebrates they contained. In this manner, a large number of specimens of *P. spectabilis* were collected (including all of the photographed specimens except Figure 2d), especially in the year 2001.

Similarly-sized crabs were photographed while alive or just after immersion in ice water, a technique that rapidly kills tropical crabs while preserving their color for up to 12 hr. Photography was done by T.L. Zimmerman (Figures 1a–f; Figure 2a–c) and Leslie Harris (Figure 2d). Associated field voucher numbers are listed in the figure captions.

After being photographed, crabs were preserved in either 70% or 95% ethanol for eventual transfer to and storage in the Crustacea collections at the Natural History Museum of Los Angeles County. One ovigerous female from the North Bay site (photographic voucher number Vc0-796, Figure 1b), with a carapace width 10.5 mm and carapace length 7.1 mm, was the parental female that formed the basis of the first description of larvae in this genus (Fransozo et al. 2001).

RESULTS

As is evident from the accompanying figures (Figures 1 and 2), specimens of *Platypodiella spectabilis* collected in the British Virgin Islands (and presumably elsewhere) exhibit a wide range of color patterns. The background color varies from a light cream or beige (Figure 1a–d) to an intense yellow (Figures 1e, f; 2a–c). Regularly-spaced and nearly circular black spots ringed with white may be present on the carapace (Figure 1c, e), or the carapace can

Figure 1. Color variation in the xanthid crab *Platypodiella spectabilis* (Herbst, 1794). a: muted background yellow, no spots, no brown patches (female, Vc0-799); b: muted background yellow with many muted large brown patches (female, Vc0-796); c: muted background yellow with spots, no brown patches (male, Vc0-800); d: muted background yellow with intense large brown patches, no spots (male, Vc0-792); e: intense background yellow with few small brown patches with spots (male, Vc0-798); f: intense background with few small brown patches, no spots (male, Vc0-801). All photographs by T. Zimmerman.

lack any semblance of similar-sized spots (Figures 1a; 2b). Irregularly-shaped brown patches sometimes occur on the carapace, and are often found on the cardiac and gastric regions (Figures 1d; 2a–c); in some specimens (e.g., Figure 2c) the brown patches are coalesced to form the majority of the color on the dorsal carapace, with only limited areas displaying white or yellow. Some individuals exhibited both irregular brown patches and more regularly-spaced and evenly-sized spots (Figure 1e). Pereiopods typically bear yellow, white and brown patches so as to appear banded. Usually the order of color on a given pereopodal

article is, from the proximal to distal end, yellow followed by brown followed by white, such that the distal end of the article is white or light beige (e.g., Figures 1a, b, d; 2b). However, in some specimens the segments of the legs are almost completely brown (Figure 2c), and in others the yellow area is bordered on both sides (rather than only on the distal side) by brown (Figure 1d).

Sex of the crab does not appear to have any noticeable effect on the color pattern; figures include both males (Figures 1c–f; 2a–b) and females (all others except Figure 2d, a juvenile). The juvenile we collected was lighter

Figure 2. Color variation in the xanthid crab *Platypodiella spectabilis* (Herbst, 1794), continued. a: coalescent solid brown patches with some muted regions (and possibly spots) over diffuse intense yellow background (male, Vc0-797); b: distinct brown patches with distinct patches of intense yellow background color (male, Vc0-794); c: coalescent solid brown patches over diffuse intense yellow background (female, Vc0-793); d: coloration pattern in a juvenile (H-2037) before carapace regions are clearly demarcated. Photographs a–c by T. Zimmerman; d by Leslie Harris.

overall, with a background carapace coloration that was off-white and with a more orange (less brown) color pattern dorsally (Figure 2d).

DISCUSSION

Color plays an important role in species identification and presumably in species-species recognition among decapods. Additionally, knowledge of coloration and color variability can assist in the recognition of species by conservationists and resource managers without extensive taxonomic training. Coral-associated crabs are among the most colorful of all decapods, with trapeziids and tetraliids well known in this regard; coral-associated carpilliids and eriphiids are also often very colorful. However, few crab species exhibit coloration as striking as in *P. spectabilis*, and there are few reports of colors or color patterns that vary to the extent shown here within a known species and within a very small geographic range (in our case, within meters). A similar situation exists for a tropical hermit crab, *Calcinus gaimardi*, in the western Pacific with at least 2 distinct color morphs that occur sympatrically (C. Tudge, pers. comm., Morgan 1991, Tudge 1995).

Although it was stated earlier that the wide range of color patterns in this species might indicate a formerly unrecognized species complex (Frasozo et al. 2001), there is virtually no doubt that all specimens illustrated here belong to the same species. Even at the larval level, the widespread *P. spectabilis* does not appear to vary much across its rather large range. Frasnoso et al. (2001) documented slight differences in the morphology of reared larvae of this species from the Caribbean and from Brazil, but overall, despite the geographic distance separating the parental females, larvae from the 2 areas were found to be quite similar.

The genus *Platypodiella* was erected by Guinot (Guinot 1967: 562) to accommodate 4 species formerly treated as *Platypodia*. Two of the species, *P. gemmata* (Rathbun, 1902) and *P. rotundata* (Stimpson, 1860) are known from the eastern Pacific. Garth (1991: 131), in discussing the Galapagos crab fauna, pointed out that the 2 Pacific species are possibly the adult (described as *P. rotundata*) and the juvenile (*P. gemmata*) of the same species, with the name *P. rotundata* having precedence. The other 2 species are Atlantic, with *P. picta* (A. Milne Edwards, 1869) restricted to the eastern Atlantic and *P. spectabilis* (Herbst,

1794) known from the western Atlantic. Guinot (1967) remarked on the similarities between species of this genus and those of *Platyactaea* and, to a lesser extent, to species of *Atergatis* and *Atergatisopsis*, as well as to members of *Platypodia*. The latter 3 genera are sometimes considered members of the xanthid subfamily Zosiminae (e.g., see Serène 1984, Clark and Ng 1998). It would be interesting to examine the range of color patterns exhibited by species in these supposedly related genera to see if color variability has a phylogenetic component.

ACKNOWLEDGEMENTS

This work was funded in part by the Biotic Surveys and Inventories program of the U.S. National Science Foundation (NSF) via grant DEB 9972100 to T.L. Zimmerman and J.W. Martin, by a PEET grant (DEB 9978193) from NSF to J.W. Martin and D. Jacobs, and by the Decapod Crustacea "Assembling the Tree of Life" grant to J.W. Martin (DEB 0531616, one of 4 awards for the decapod ATOL project). We thank NSF and especially D. Causey for support and encouragement during our time on Guana Island. We also thank other members of the Guana Island field team: L. Harris, D. Cadien, R. Ware, T. Haney, K. Fitzhugh, and G. Hendler. Finally, we thank the Falconwood Corporation and especially L. Jarecki for allowing us to conduct this work on Guana Island. The manuscript was improved substantially by the constructive comments of C. Tudge and an anonymous reviewer.

LITERATURE CITED

- Abele, L.G. and W. Kim. 1986. An illustrated guide to the marine decapod crustaceans of Florida. State of Florida Department of Environmental Regulation Technical Series, Vol. 8, no. 1 (in 2 parts). 760 p.
- Campbell, N.A. and R.J. Mahon. 1974. A multivariate study of variation in two species of rock crab of the genus *Leptograpsus*. *Australian Journal of Zoology* 22:417–425.
- Chace, F.A., Jr., J.J. McDermott, P.A. McLaughlin, and R.B. Manning. 1986. Decapoda. In: W. Sterrer, ed. *Marine Fauna and Flora of Bermuda*. Wiley-Interscience, John Wiley and Sons, New York, NY, USA, p. 312–358.
- Clark, P.F. and P.K.L. Ng. 1998. The larval development of the poisonous mosaic crab, *Lophozozymus pictor* (Fabricius, 1798) (Crustacea, Decapoda, Brachyura, Xanthidae, Zosiminae), with comments on familial characters for first stage zoeas. *Zoosystema* 20:201–220.
- Coelho, P.A. and M. de Araújo Ramos. 1972. A constituição e a distribuição da fauna de decápodos do litoral leste da América do Sul entre as latitudes de 5°N e 39°S. *Trabalhos do Instituto Oceanográfico da Universidade Federal Pernambuco, Recife* 13:133–236.
- Felder, D.L. 1973. An annotated key to crabs and lobsters (Decapoda, Reptantia) from coastal waters of the north-western Gulf of Mexico. Center for Wetland Resources, Louisiana State University, Baton Rouge, LA, USA, 103 p.
- Fransozo, A., M.L. Negreiros-Fransozo, J.W. Martin, and S.E. Trautwein. 2001. Morphology of the first zoeal stage of *Platypodiella spectabilis* (Herbst, 1794) (Decapoda, Brachyura, Xanthidae) obtained in the laboratory. *Gulf and Caribbean Research* 13:79–85.
- Garth, J.S. 1991. Taxonomy, distribution, and ecology of Galápagos Brachyura. In: M.J. James, ed. *Topics in Geobiology, Volume 8: Galápagos Marine Invertebrate*. Plenum Press, New York, NY, USA, p. 123–145.
- Guinot, D. 1967. Recherches préliminaires sur les groupements naturels chez les Crustacés Décapodes Brachyours III. A propos des affinités des genres *Dairoides* Stebbing et *Daira* de Haan. *Bulletin du Museum National d'Histoire Naturelle, 2e Série*, 39:540–563.
- Humann, P. 1992. Reef Creature Identification: Florida, Caribbean, Bahamas. New World Publications, Inc., Vaughan Press, Orlando, FL, USA.
- Humann, P. and N. DeLoach. 2002. Reef Creature Identification: Florida, Caribbean, Bahamas. New World Publications Inc., Jacksonville, FL, USA.
- McLaughlin, P.A., D.K. Camp, L.G. Eldredge, D.L. Felder, J.W. Goy, H.H. Hobbs, III, B. Kensley, R. Lemaitre, and J.W. Martin. 2005. Order Decapoda. In: D. Turgeon, ed. *Common and Scientific Names of Aquatic Invertebrates of the United States and Canada. Names of Crustaceans Special Publications. Vol. 31. American Fisheries Society Special Publication*, Bethesda, MD, USA, p. 209–326.
- Melo, G.A.S. 1998. Malacostraca—Eucarida. Brachyura. Oxyrhyncha and Brachyrhyncha. In: P. Young, ed. *Catalogue of Crustacea of Brazil. Museu Nacional (Série Livros no. 6)*, Rio de Janeiro, BR, p. 455–515.
- Morgan, G.J. 1991. A review of the hermit crab genus *Calcinus* Dana (Crustacea: Decapoda: Diogenidae) from Australia, with descriptions of two new species. *Invertebrate Taxonomy* 5:869–913.
- Powers, L. 1977. A catalogue and bibliography to the crabs (Brachyura) of the Gulf of Mexico. *Contributions in Marine Science* 20 (supplement):1–190.
- Rathbun, M.J. 1930. The Cancroid crabs of America of the families Euryalidae, Portunidae, Atelecyclidae, Cancridae, and Xanthidae. *Bulletin of the US National Museum* 152:1–609.
- Serène, R. 1984. Crustacés Décapodes Brachyours de l'Océan Indien Occidental et de la Mer Rouge, Xanthoidea: Xanthidae et Trapeziidae. Avec un addendum par Crosnier, A.: Carpiliidae et Menippidae. *Faune Tropicale* 24:1–400.
- Sterrer, W. (editor). 1986. *Marine Fauna and Flora of Bermuda*. John Wiley and Sons, New York, NY, USA.
- Tudge, C.C. 1995. Hermit crabs of the Great Barrier Reef and coastal Queensland. Backhuys Publishers, Leiden, The Netherlands, 40 p.
- Williams, A.B. and D.L. Felder. 1989. Analysis of stone crabs: *Menippe mercenaria* (Say), restricted, and a previously unrecognized species described (Decapoda: Xanthidae). *Proceedings of the Biological Society of Washington* 99:517–543.

- Williams, A.B., L.G. Abele, D.L. Felder, H.H. Hobbs, Jr., R.B. Manning, P.A. McLaughlin, and I. Perez Farfante. 1989. Common and scientific names of aquatic invertebrates from the United States and Canada: Decapod Crustaceans. American Fisheries Society, Special Publication 17:1-77.
- Zimmerman, T.L. and D.L. Felder. 1991. Reproductive ecology of an intertidal brachyuran crab, *Sesarma* sp. (nr. *reticulatum*), from the Gulf of Mexico. Biological Bulletin 181:387-401.