

12-1994

# Symbiotic N Fixation by *Vigna luteloa*, A Legume of Low-Salinity Gulf Coast Marshes

C.R. Crozier  
*Louisiana State University*

J.A. Nyman  
*Louisiana State University*

R.D. DeLaune  
*Louisiana State University*

DOI: 10.18785/negs.1302.08

Follow this and additional works at: <https://aquila.usm.edu/goms>

---

### Recommended Citation

Crozier, C., J. Nyman and R. DeLaune. 1994. Symbiotic N Fixation by *Vigna luteloa*, A Legume of Low-Salinity Gulf Coast Marshes. *Northeast Gulf Science* 13 (2). Retrieved from <https://aquila.usm.edu/goms/vol13/iss2/8>

This Article is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in Gulf of Mexico Science by an authorized editor of The Aquila Digital Community. For more information, please contact [Joshua.Cromwell@usm.edu](mailto:Joshua.Cromwell@usm.edu).

## Short Papers and Notes:

### SYMBIOTIC N FIXATION BY *Vigna luteola*, A LEGUME OF LOW-SALINITY GULF COAST MARSHES

Fixation of atmospheric N may be the largest N input to low-salinity marshes of the Gulf Coast region (DeLaune and Patrick, 1990). Although several estimates of N fixation (acetylene reduction) by mats of blue-green algae or by the predominantly non-leguminous marsh vegetation are available (Whitney et al., 1975; Teal et al. 1979; Casselman et al., 1981; DeLaune and Patrick, 1990), little data exists concerning marsh legumes. *Vigna luteola* (Jacq.) Benth. (deerpea) is a perennial, viny legume occurring in fresh to brackish marshes of the Gulf Coast region. *Vigna* may account for 1.5% of total plant abundance throughout the Louisiana coastal marsh zone, and for 3.8% of total plant abundance in the intermediate salinity marshes (Chabreck, 1970).

The objectives of this study were to estimate N fixation (acetylene reduction) rates attributable to both symbiotic sources associated with *Vigna*, and to free-living N fixers in adjacent soils; and to compare landscape-level estimates of N fixation with and without inclusion of *Vigna* patches.

#### MATERIALS AND METHODS

On 17 August 1993, three replicate soil cores (14.5 cm diameter x 15 cm deep) were collected from an intermediate marsh at the eastern margin of Four League Bay in Terrebonne Parish, Louisiana. Each core was centered on the base of an individual *Vigna* plant. An abundance of root nodules was visible on the marsh soil surface and to a depth

of approximately 1-2 cm. Upon returning to the laboratory, after approximately 24 hours, nodulated root segments were separated from the soil matrix. An  $C_2H_2$  reduction assay was used to estimate N fixation rates (Turner and Gibson, 1980). Acetylene was generated from  $CaC_2$  and  $H_2O$  (Miller, 1965). Samples were placed in 140 ml glass jars, and incubated at room temperature in either air or 90% air: 10%  $C_2H_2$ . Headspace gas was sampled at 1, 4, and 16 h and concentrations of  $C_2H_4$  were determined by gas chromatography (Perkin-Elmer Model 900 GC equipped with a flame ionization detector). Production rates of  $C_2H_4$  by nodulated root segments were approximately linear over the first 4 hours of incubation. For soils, 16 hours were required for detectable  $C_2H_4$  production.

On 9 September 1993, three different intermediate salinity marsh sites near Raccourci Bay in Terrebonne Parish, Louisiana were sampled. At each site, three replicate pairs of soil cores (14.5 cm diameter x 2 cm deep) were collected from within dense *Vigna* patches and from the adjacent *Spartina patens* stand. Cores were collected from predetermined positions along a transect, with no attempt made to center on a *Vigna* plant base. Immediately after collection, each soil core was cut into four equal wedge-shaped subsamples. Two diagonally opposite subsamples were sealed in a 450 ml glass jar. An air sample was removed through a rubber sampling port, and sufficient  $C_2H_2$  was added to achieve a 10% concentration in the jar headspace. A third subsample was pooled with subsamples from the other two cores of the same vegetation type, and sealed in a glass jar to which no  $C_2H_2$  was to be added (control). Additional controls were established with  $C_2H_2$  incubated in

sample jars containing only deionized H<sub>2</sub>O and air. Samples were placed in an insulated box to reduce temperature fluctuations. Immediately after, and at 2, 4, 8 and 24 hours after injecting C<sub>2</sub>H<sub>2</sub>; 15 ml headspace gas samples were transferred to vacutainers prior to analysis. Headspace gas was maintained at atmospheric pressure by injecting 15 ml of air after each sample was collected.

N fixation rates were calculated from the moles of C<sub>2</sub>H<sub>4</sub> produced during the linear phase of incubation (0-2 hours for the field incubations), the sediment surface area represented by the field samples, and a 3:1 conversion ratio for moles of C<sub>2</sub>H<sub>2</sub> reduced:moles of N<sub>2</sub> reduced (Turner and Gibson, 1980). Treatment means were compared using Wilcoxon's signed ranks test (Siegle and Castellan, 1988).

## RESULTS AND DISCUSSION

Both nodulated root segments and nodule-free soil reduced C<sub>2</sub>H<sub>2</sub> in preliminary assays (Table 1). No detectable C<sub>2</sub>H<sub>4</sub> was produced by either nodulated roots or soils in the absence of C<sub>2</sub>H<sub>2</sub>. When samples collected on 9 September 1993 were incubated, only one of the six control microcosms produced any detectable C<sub>2</sub>H<sub>4</sub>. Even in this case, C<sub>2</sub>H<sub>4</sub> production was <1% of that in the C<sub>2</sub>H<sub>2</sub>-treated microcosms.

Rates of N fixation declined in successive field sites (Table 2). This may have been due to the cooler incubation temperatures as the afternoon progressed. Ambient temperatures on this day declined from 32°C at noon, when the first samples were collected, to 28°C at 6 pm, the time at which the samples from the third site had incubated for 2 hours.

N fixation rates across sites were much higher in soils collected from *Vigna* patches (1.90 mg N m<sup>-2</sup>d<sup>-1</sup>) than from the surrounding *Spartina* stands (0.20 mg

N m<sup>-2</sup>d<sup>-1</sup>, Table 2). Differences between vegetation types were significant at the 0.05 probability level.

Landscape-level N fixation rates for these intermediate marshes can be calculated assuming that *Vigna* occupies 3.8% of the surface area (Chabreck, 1970), and that the remaining marsh surface fixes N at the same rate as did the *Spartina* patches. Relative abundance of *S. patens* in this marsh type was reported to be five times greater than that of any other species, and no other legumes had a relative abundance even one-tenth as great as the relative abundance of *Vigna* (Chabreck, 1970). Fixation estimates are then 1.92 g N ha<sup>-1</sup> d<sup>-1</sup> without *Vigna* patches, and 2.64 g N ha<sup>-1</sup> d<sup>-1</sup> including *Vigna* patches (Table 3). Thus, failing to sample and account for N fixation rates within <5% of the land area could result in a 27% underestimation of N fixation rates.

Ecosystem N budgets for low salinity marshes of this region suggest that N fixation by free-living microbes is a major source of N inputs (60-70 kg N ha<sup>-1</sup> yr<sup>-1</sup>; DeLaune and Patrick, 1990). Accretion of excess N (20-40 kg N ha<sup>-1</sup> yr<sup>-1</sup>), was attributed to sediment deposition (DeLaune and Patrick, 1990). The expected relative errors in N fixation estimates based on this study are of similar magnitude as were the excess N accretion rates of the earlier study, and thus may lead to different conclusions

**Table 1.** Rates of C<sub>2</sub>H<sub>4</sub> production by nodulated *Vigna luteola* roots and underlying soils. Values represent means (CV) of three replicate samples.

Material	C <sub>2</sub> H <sub>4</sub> production Rate	
	C <sub>2</sub> H <sub>2</sub> †treated	Control
	— nmol g <sup>-1</sup> d <sup>-1</sup> —	
Nodulated roots †	33.9 (130)	0
Soil (2-10 cm depth)§	0.15 (63)	0

† C<sub>2</sub>H<sub>4</sub> production rates were calculated based on the 0-4 hour incubation interval.

§ C<sub>2</sub>H<sub>4</sub> production rates were calculated based on the 0-16 hour incubation interval.

**Table 2.** N fixation rates of intermediate marsh soils. Values given are the means (CV) of three replicate C<sub>2</sub>H<sub>2</sub>-treated microcosms per site.

Site	<i>Spartina patens</i>	<i>Vigna luteola</i>
	— mg N m <sup>-2</sup> d <sup>-1</sup> —	
1	0.33 (90)	4.56 (65)
2	0.15 (71)	0.83 (145)
3	0.12 (89)	0.31 (100)
Overall mean*	0.20 (55)	1.90 (122)

\*Treatment means differed significantly at the 0.05 level using Wilcoxon's signed rank test.

about sediment input rates.

This study demonstrates the potential significance of N fixation by a nodulated legume to low-salinity marsh N budgets, and how estimates of ecosystem function can be influenced by sampling schemes within a heterogeneous landscape.

### ACKNOWLEDGMENTS

This research was supported by the National Institute for Global Environmental Change (funded by the Department of Energy) Cooperative Agreement #DE-FCO3-90ER610101, and by the U.S. Geological Survey grant number 14-08-001-23320.

### LITERATURE CITED

Casselman, M. E., W. H. Patrick, Jr., and R. D. DeLaune. 1981. Nitrogen fixation in a gulf coast salt marsh. *Soil Sci. Soc. Am. J.* 45: 51-56.

Chabreck, R. H. 1970. Marsh zones and vegetative types in the Louisiana coastal marshes. Ph.D. Diss. Louisiana State University, Baton Rouge, Louisiana. 112 pp.

DeLaune, R. D., and W. H. Patrick, Jr. 1990. Nitrogen cycling in Louisiana gulf coast brackish marshes. *Hydrobiol.* 199: 73-79.

Miller, S. A. 1965. Acetylene: its properties, manufacture and uses. Volume I. Academic Press. New York. 800 pp.

Siegel, S., and N. J. Castellan, Jr. 1988. Nonparametric statistics for the behavioral sciences. Second Edition. McGraw-Hill, Inc., New York. 399 pp.

Teal, J. M., I. Valiela, and D. Berlo. 1979. Nitrogen fixation by rhizosphere and free-living bacteria in salt marsh sediments. *Limnol. Oceanogr.* 24: 126-132.

Turner, G. L., and A. H. Gibson. 1980. Measurement of nitrogen fixation by indirect means. pp. 111-138. *In* Bergersen, F. J. (ed.) *Methods for Evaluating Biological Nitrogen Fixation*. John Wiley and Sons, Ltd., New York.

Whitney, D. E., G. M. Woodwell, and R. W. Howarth. 1975. Nitrogen fixation in flax pond: a Long Island salt marsh. *Limnol. Oceanogr.* 20: 640-643.

C. R. Crozier\*, J. A. Nyman and R. D. DeLaune, Wetland Biogeochemistry Institute, Louisiana State University, Baton Rouge, Louisiana 70803-7511

\*Corresponding author  
Submitted a Note to *Northeast Gulf Science*, December 29, 1993. Revised April 29, 1994

**Table 3.** Estimates of landscape-level N fixation rates.

Vegetation type	Relative Abundance†	N Fixation Rate	Landscape-Level N Fixation Rate	
			Amount	Relative Amt.
	—%—	mg m <sup>-2</sup> d <sup>-1</sup>	g ha <sup>-1</sup> d <sup>-1</sup>	—%—
<i>Vigna</i>	3.8	1.90	0.72	27
without <i>Vigna</i> §	96.2	0.20	1.92	73
Total	100.0		2.64	100

† from Chabreck, 1970

§ *Spartina patens* patches were considered representative of all other marsh patch types.