Gulf and Caribbean Research

Volume 20 | Issue 1

January 2008

Seagrass Distribution in the Pensacola Bay System, Northwest Florida

Michael A. Lewis

U.S. Environmental Protection Agency

Richard Devereux

U.S. Environmental Protection Agency

Pete Bourgeois

National Wetlands Research Center

Follow this and additional works at: <https://aquila.usm.edu/gcr>

Part of the [Marine Biology Commons](#)

Recommended Citation

Lewis, M. A., R. Devereux and P. Bourgeois. 2008. Seagrass Distribution in the Pensacola Bay System, Northwest Florida. *Gulf and Caribbean Research* 20 (1): 21-28.

Retrieved from <https://aquila.usm.edu/gcr/vol20/iss1/4>

DOI: <https://doi.org/10.18785/gcr.2001.04>

This Article is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in *Gulf and Caribbean Research* by an authorized editor of The Aquila Digital Community. For more information, please contact Joshua.Cromwell@usm.edu.

SEAGRASS DISTRIBUTION IN THE PENSACOLA BAY SYSTEM, NORTHWEST FLORIDA

Michael A. Lewis^{1*}, Richard Devereux¹, and Pete Bourgeois²

¹U.S. Environmental Protection Agency, National Health & Environmental Effects Research Laboratory, Gulf Ecology Division, Gulf Breeze, Florida 32561 USA, *e-mail: Lewis.Michael@epa.gov

²U.S. Geological Survey, National Wetlands Research Center, Gulf Breeze Project Office, Gulf Breeze, Florida 32561 USA

ABSTRACT: Aerial surveys of seagrass coverage in the Pensacola Bay system (PBS) have been conducted during 1960, 1980, 1992 and 2003. This report summarizes the results for the 2003 survey and compares the results to those previously reported for other surveys. The estimated coverage of seagrass for the PBS during 2003 was 1,654 ha. Continuous and patchy coverages ranged from 0 to 684 ha and 11 to 543 ha, respectively, for five PBS subsystems. In 2003, the majority of seagrass coverage occurred in Santa Rosa Sound (76%). Declines in total coverage occurred for East Bay (93%) and Escambia Bay (75%) whereas increases were observed for Pensacola Bay (32%) and Santa Rosa Sound (8%). The approximate 9% decline (about 160 ha or 395 a) in total coverage since 1992 represents an estimated 7 to 8 million dollar loss in ecological services. The changes in coverage are likely due to naturally occurring and anthropogenic factors but it is not possible to differentiate the relative contributions of these factors alone and in combination on seagrass distribution. The ability of seagrasses to exist long-term in Florida's fourth largest estuarine system is uncertain due to the adverse effects of rapid urbanization in the watershed. Active resource management which includes more frequent *in-situ* monitoring and aerial assessment and the availability of relevant water and sediment quality criteria protective of submerged aquatic vegetation are needed to prevent future declines.

INTRODUCTION

Seagrasses have at least 13 ecological roles (Dawes et al. 2004) and they support a diverse biotic community that may contain as many as 113 species of epiphytes, 148 macroalgal species, 80 macrofaunal species, and 75 fish species (Humm 1964, Virnstein et al. 1983, Zieman and Zieman 1989, DeTorch et al. 1996). Seagrass meadows including those dominated by *Thalassia testudinum* Konig (turtle grass), the most abundant species in the Pensacola Bay System (PBS), supported twice the macrofauna than did unvegetated sediments (Santos and Simon 1974, Virnstein et al. 1983). The economic importance of one seagrass acre has been estimated to be between \$9,000 and \$28,000 (Texas) and \$20,500 (Florida) due to commercial, recreational and storm protection functions (Handley et al. 2007) and \$19,000 based on nutrient cycling (Costanza et al. 1997). About 1.2 million of 59 million hectares (ha) of seagrasses have been destroyed worldwide during the last decade and, similar to corals, seagrasses are considered to be in a crisis stage. Declines have occurred at 40 locations (Short and Wylie-Echeverria 1996, Hemminga and Duarte 2000) including the Gulf of Mexico (GOM), where seagrass coverage has been reduced 20 to 100% during the past 50 yrs (Dawes et al. 2004, USGS and GOMP 2004). For example, about 85% of the seagrasses along Florida's coasts have been destroyed by 1992 (USEPA 1992).

Florida's Gulf Coast contains about 680,000 ha of seagrasses of which about 2% (17,474 ha) is adjacent to the Florida panhandle region (Madley et al. 2003). The trends in coverage and condition of the seagrasses in northwest

Florida are considered to be poorly understood (Dawes et al. 2004). However, seagrass research for the PBS, the focus of this report, has been conducted intermittently during the past 40 yrs (Table 1) and includes four major aerial surveys conducted since the 1960s. The objectives of this report are to summarize the results of the 2003 aerial survey and to compare the results primarily to those for the previous survey conducted during 1992.

MATERIALS AND METHODS

Study Area

The PBS is located in northwest Florida and its watershed consists of about 18,000 km² of forests, agricultural lands, and urban and industrial areas (Figure 1). It is the fourth largest estuary in Florida and extends 32 km inland and comprises about 886 km of coastline and 435 km of inland waterways. Mean water residence time is about 25 d (Solis and Powell 1999). The PBS is comprised of five subsystems: Big Lagoon, East Bay, Escambia Bay, Pensacola Bay, and Santa Rosa Sound (Figure 1, Table 2). Santa Rosa Sound and Big Lagoon are marine lagoons that are parallel to the GOM and retain high salinities due to limited freshwater input. Both areas contain sections that are classified as Outstanding Florida Waters (FDEP 2001).

Aerial Surveys

Although aerial surveys of the PBS are available since the 1940s, only those conducted during 1960, 1980, 1992 and 2003 were specific for determining seagrass coverage. Results of the 1960, 1980, and 1992 surveys have been report-

ed previously (FDEP 2001, Schwenning et al. 2007) and the results for the October 2003 survey form the basis of this report. The experimental techniques for these surveys have differed, and these differences need to be considered in the context of the conclusions for the across-year comparisons reported here. Methodologies for the 1992 and 2003 surveys were similar but not identical. The seagrass data were derived from interpretation of 1:24,000 natural color scale photographs (1992 survey) and 1:12,000 scale color infrared photographs (2003 survey). Personnel from NASA (Stennis, MS) and Aerial Cartographics of America (Orlando, FL) performed the 1992 and 2003 flights, respectively. Black and white photographs (1:24,000 scale) and natural color photographs (1:24,000 scale) were used for the 1960 and 1980 surveys, respectively. Personnel from U.S. Geological Survey's National Wetlands Research Center (NWRC, Lafayette, LA) conducted the mapping procedure which included photo-interpretation of the aerial photographs, cartographic transfer, and digitization for all surveys. The classification scheme for all surveys was derived by USGS/NWRC based on the coastal land cover classification system of the National Oceanic and Atmospheric Administration's Coastwatch Change Analysis Project (NOAA 2003). The amount of groundtruthing or field verification of the aerial surveys has been variable, and for the 2003 survey it consisted of single visits to the five PBS subsystems to confirm seagrass presence. No *in-situ* measurements of plant condition or species identification were performed.

RESULTS AND DISCUSSION

Seagrass Coverage

An estimated 1,654 seagrass-vegetated ha (4,085 acres) were present in the PBS based on the October 2003 aerial survey (Table 3). Of this total, most coverage occurred in Santa Rosa Sound (76%). Seagrass coverage as a percent of total surface area was 18% (Santa Rosa Sound), 5% (Big Lagoon), 1% (Pensacola Bay) and < 1% (Escambia Bay, East Bay). Continuous and patchy coverages ranged from 0 to 684 ha and 11 to 543 ha, respectively in the PBS (Table 3). About 52% of the 2003 total coverage was continuous, and almost all continuous coverage (99%) occurred in Santa Rosa Sound and Big Lagoon.

The 2003 total seagrass coverage (1,654 ha) represents an almost 9% reduction relative to the estimated total for the 1992 survey (1,814 ha). Reductions in total coverage relative to 1992 occurred for East Bay (93%), and Escambia Bay (75%) while increases were observed for Pensacola Bay (32%) and Santa Rosa Sound (8%) (Figure 2). Total coverages in Big Lagoon for 1992 and 2003 were almost identical. Continuous coverage decreased in Escambia Bay (69%), Santa Rosa Sound (14%), Pensacola Bay (100%), and East Bay (100%), which contrasted an almost 64% increase in coverage for Big Lagoon. Patchy coverage decreased in Big

TABLE 1. Listing of seagrass and habitat condition research conducted for the Pensacola Bay System, Florida.

Subsystem	References
Big Lagoon	Hopkins 1973 Heck et al. 1996 Lores et al. 2000 FDEP 2001
East Bay	Van Breedveld 1966 McNulty et al. 1972 Rogers and Blisterfield 1975
Escambia Bay	Moore 1963 USDOI 1970 Livingston et al. 1971 McNulty et al. 1972 Hopkins 1973 Rogers 1974 Rogers and Blisterfield 1975 Woodward and Clyde 1997 Lewis et al. 2000 Lores et al. 2000 Lores and Sprecht 2001 Murrell et al. 2002 Murrell and Lores 2004
Pensacola Bay	Moore 1963 McNulty et al. 1972 Rogers and Blisterfield 1975 Rodriguez and Hunner 1994 Murrell et al. 2002
Santa Rosa Sound	Moore 1963 Van Breedveld 1966 McNulty et al. 1972 Hopkins 1973 Rogers and Blisterfield 1975 Winter 1978 Macauley et al. 1988 Heck et al. 1996 Lores et al. 2000 FDEP 2001 Lewis et al. 2001, 2002
Pensacola Bay System (general)	USEPA 1975 Williams 1981 Lewis 1986 Ridenauer and Shambaugh 1986 Jones et al. 1992 Collard 1991 Thorpe et al. 1997 DeBusk et al. 2002 Dawes et al. 2004 McRae et al. 2004 USEPA 2004 USEPA 2005 Schwenning et al. 2007

Figure 1.

The Pensacola Bay System in northwest Florida and its watershed (insert). Shaded areas represent continuous (green) and patchy (red) seagrass coverage estimated from a 2003 aerial survey.

TABLE 2. Geomorphological characteristics and water body classifications for Pensacola Bay subsystems. Some data adapted from USEPA (1975). N/A – not available.

Subsystem	Surface Area (km²)	Volume (10⁶ m³)	Mean Depth (m)	Water Body Classification¹	Florida Impaired Water²	Assessed Parameters
Big Lagoon	46.8	N/A	N/A	2 (OFW)	N	-
East Bay	25.9	259.3	2.4	2	Y	Fecal coliforms, Nutrients
Escambia Bay	57	225.7	2.4	3M	Y	Nutrients
Pensacola Bay	133.6	793.8	5.9	3M	Y	Bacteria
Santa Rosa Sound	68.2	N/A	N/A	2 (OFW)	Y	Fecal coliforms

¹ 2 = shellfish propagation or harvesting, 3M = Recreation, propagation and maintenance of well-balanced populations of fish and wildlife (FDEP 1996). OFW = Portion designated as Outstanding Florida Water (OFW) (FDEP 2004).

² Contains segments for TMDL development (FDEP 2006).

Lagoon (51%), East Bay (93%), and Escambia Bay (76%) and increased in Pensacola Bay (50%) and Santa Rosa Sound (58%).

Historical Perspective

The numerous seagrass investigations conducted in the PBS have been temporally and spatially inconsistent and research methodologies have varied. Seagrass mapping has been the focus of most studies, and only limited information is available describing physiological and morphological parameters of plant condition such as above- and below-ground biomass, shoot density, blade height and epiphyte/biomass ratio, which have been reported for grasses in Big Lagoon, Santa Rosa Sound and Escambia River delta region (Heck et al. 1996, Lores et al. 2000, FDEP 2001).

Historically, the largest seagrass declines within the PBS occurred between 1960 and 1980 and subsequent declines have been less severe (Table 3, Figure 2). Seagrass meadows in Santa Rosa Sound and Big Lagoon, with few exceptions, have dominated the PBS since 1960 (coverage range as a percent of total = 75 to 90%). Their combined dominance increased 12% since 1992 to 87% of total coverage in 2003. The consistent coverages in these areas are due, at least in part, to relatively less urbanization of the shorelines (sections are included in a state park and a national seashore) and to limited freshwater input which stabilizes salinity and reduces the entry of watershed contaminants. In a detailed study conducted in these areas (Heck et al. 1996), seagrasses declined during 1993 – 1995 in both areas due to a combination of propeller scarring, reduced water clarity, changes in salinity and burial due to hurricanes. Lores et al. (2000) reported seagrass coverage was also decreasing in Big Lagoon during 1997 and 1998.

Seagrass research has been conducted less frequently in East, Escambia and Pensacola Bays, where seagrasses have

been relatively uncommon since the 1960s (range of combined total coverages = 9 to 25%; Table 3). Seagrass coverage in East Bay was almost non-detectable and the least of any subsystem in 2003. Submerged aquatic vegetation in Escambia Bay is primarily limited to tidal freshwater grasses, *Vallisneria americana* Michx (tape grass) in the upper reaches and the more salinity tolerant *Ruppia maritima* L. (widgeon grass) in more seaward areas. *Vallisneria americana* is not a true seagrass, although its ecological value is similar to that of seagrasses. This species is included in this analysis, since prior assessments did not differentiate between species. Total coverage in Escambia Bay fluctuates greatly, with a

TABLE 3. Continuous (C) and patchy (P) seagrass coverage (hectares) in Pensacola Bay subsystems. Data for 1960, 1980 and 1992 aerial surveys from Schwenning et al. (2007).

Subsystem		AERIAL SURVEY			
		1960	1980	1992	2003
Big Lagoon	C	107	193	99	162
	P	164	43	118	58
East Bay	C	45	12	5	0
	P	431	87	160	11
Escambia Bay	C	4	4	36	11
	P	101	20	143	34
Pensacola Bay	C	44	10	13	0
	P	328	46	101	151
Santa Rosa Sound	C	1,247	850	796	684
	P	1,387	629	343	543
TOTAL		3,858	1,894	1,814	1,654

historical maximum occurring in 1992 followed by a 75% decline in 2003. Lores et al. (2000) reported increased coverage of *V. americana* near the Escambia River delta for 1997 and 1998 relative to 1992, but coverage decreased during 2000 due to drought and high salinity (Lores and Sprecht 2001). Although continuous seagrass coverage in Pensacola Bay was not detectable in 2003, patchy coverage increased from 101 ha to 151 ha since 1992.

Causes of Seagrass Declines

The factors responsible for declines in seagrass coverage for the PBS, other than direct effects due to mechanical and physical factors, have been more often speculative than confirmed. Supportive documentation is limited, and current understanding of the causative factors responsible for the seagrass changes alone and in combination remains elusive, which has limited effective restoration efforts and resource management. Declines have been attributed to controllable factors such as point and nonpoint source contaminants, prop scarring, dock shading, armoured shorelines, and dredging, as well as to the non-controllable effects of episodic weather events which have been increasing in recent years. A few investigators have reported the effects of nutrients (Heck et al. 1996, Lores et al. 2001) and low salinity (Lores and Specht 2001) and the potential for chemical phytotoxicity (Lewis et al. 2007) on seagrasses within the PBS.

The PBS is a contaminant-impacted estuary based on the results of many reports included in Table 1. Turbidity due to erosion and accelerated eutrophication have reduced light penetration in all subsystems. In addition, regulatory effects-based guidelines and criteria to protect marine life in water and sediment have been commonly exceeded in several subsystems for several non-nutrient contaminants (for example, DeBusk et al. 2002, USEPA 2005), suggesting a potential for toxicity. This is the case for Escambia Bay, which is the most contaminated subsystem and a priority site for conservation (Beck et al. 2000). In contrast, Santa Rosa Sound is the least chemically contaminated area within the PBS (FDEP 2001, Lewis et al. 2007).

Despite differences in environmental conditions among subsystems, it remains to be determined if the sensitivities of seagrasses and other submerged aquatic vegetation to non-nutrient contaminants is equivalent to those for zoobenthos and fish for which most regulatory numerical guidelines and criteria have been developed. In addition to this uncertainty is the continued absence of national and state numerical nutrient criteria which further hinders effective management of these marine angiosperms.

Summary and Recommendations

Despite the limitations and sources of error associated with aerial seagrass photography (see Carlson and Madley 2007), it was clear that there has been no net gain in seagrass coverage within the PBS since 1992. The few site-specific gains in coverage were overshadowed by declines in other

Figure 2.

Total seagrass coverage (hectares) for five subsystems in the Pensacola Bay System based on four aerial surveys. Data for 1960, 1980 and 1992 surveys from Schwenning et al. (2007).

subsystems which resulted in an estimated net loss of about 9% or 160 ha (~ 395 acres) system-wide. The relevance of the 2003 coverage to current coverage (2007) is unknown. Two major hurricanes, Ivan (2004) and Dennis (2005) made almost direct landfall near the PBS. Their impacts have not been reported in the scientific literature, nor have impacts associated with the ongoing urbanization of the PBS shoreline that has occurred since 2003.

The rapid urbanization of the PBS watershed is expected to continue. The populations in Escambia and Santa Rosa Counties is predicted to increase 20% and 64%, respectively, by 2020 (Zwick and Carr 2006). The magnitude of this urban transformation on near-shore seagrasses is unknown. This uncertainty will remain until an effective and long-term resource management plan is implemented, particularly for near-shore areas containing extensive seagrass coverage such as Santa Rosa Sound. Several management plans have been proposed (Rogers and Blisterfield 1975, Collard 1991, FDEP 2001), and these should include frequent aerial assessments (every 3 to 5 yrs) to determine coverage and more frequent *in-situ* evaluation to determine plant condition. These assessments are important so that in the long-term biocriteria predictive of habitat quality can be developed. Other management considerations should include promotion of shoreline configurations supportive of seagrass meadows and increased efforts to control coastal entry of non-point source contaminants. Of additional importance is the need to establish a separate designated use category to protect coastal submerged vegetation as well as the development of supportive regulatory numerical criteria for common near-shore contaminants and those of emerging concern (Daughton 2005).

Finally, assessment of the economical value of ecosys-

tem services has become increasingly important to the environmental policy and management process (Costanza et al. 1997, Carpenter and Turner 2000, USEPA 2006). The loss of ecological services associated with the estimated 395

acre decline in seagrass coverage for the PBS since 1992 represents an approximate \$8.1 million impact (\$20,500/acre; Handley et al. 2007) or, if based on nutrient cycling, a \$7.5 million loss (\$19,000/acre; Costanza et al. 1997).

ACKNOWLEDGMENTS

We thank P. Rogers (National Caucus & Center for Black Aged, Washington, DC) for manuscript preparation and P. Soderlund (Computer Services Corporation, El Segundo, CA) for graphics support. Linda Harwell (U.S. Environmental Protection Agency, Gulf Breeze, FL) assisted in data analysis. Aerial photography and analysis were conducted as part of an Intraagency Agreement with the U.S. Geological Survey's National Wetlands Research Center, Lafayette, LA.

LITERATURE CITED

- Beck, M.W., M. Odaya, J.J. Bachant, J. Bergan, B. Keller, R. Martin, R. Mathews, C. Porter, and G. Ramseur. 2000. Identification of priority sites for conservation in the northern Gulf of Mexico: an ecoregional plan. The Nature Conservancy, Arlington, VA, USA. 47 p.
- Carlson, P.R. and K. Madley. 2007. State of Florida. In: L. Handley, D. Altman, and R. DeMay, eds. Seagrass Status and Trends in the Northern Gulf of Mexico: 1940 – 2002. U.S. Geological Survey Scientific Investigations Report 2006 – 5287 and U.S. Environmental Protection Agency 855-R-04-003, U.S. Geological Survey, Reston, VA, USA, p 98-114.
- Carpenter, S.R. and M. Turner. 2000. Opening the black boxes: ecosystem science and economic evaluation. *Ecosystems* 3: 1-3.
- Collard, S. 1991. Management options for the Pensacola Bay System: the potential value of seagrass transplanting and oyster bed refurbishment programs. Water Resources Special Report 91-4. Northwest Florida Water Management District, Havana, FL, USA, 33 p.
- Costanza, R., R. d'Arge, R. de Groot, S. Farber, M. Grasso, B. Hannon, K. Limberg, S. Naeem, R.V. O'Neill, J. Parnelo, R.G. Raskin, P. Sutton, and M. van den Belt. 1997. The value of the world's ecosystem services and natural capital. *Nature* 387:253-260.
- Dawes, C.J., R.C. Phillips, and G. Morrison. 2004. Seagrass communities of the Gulf Coast of Florida: Status and Ecology. Tampa Bay Estuary Program, St. Petersburg, FL, USA, 74 p.
- Daughton, C.G. 2005. Emerging chemicals as pollutants in the environment: a 21st century perspective. *Renewable Resources Journal* 23:6-23.
- DeBusk, W.F., I. Poyer, and I. Herzfeld. 2002. Sediment quality in the Pensacola Bay System. Final Report. Northwest Florida Water Management District, Havana, FL, USA, 66 p.
- DeTroch, M., J. Mees, I. Papadopoulos, and E.O. Wakwabi. 1996. Fish communities in a tropical bay (Gazi Bay, Kenya): seagrass beds vs. unvegetated areas. *Netherlands Journal of Zoology* 46:236-252.
- Florida Department of Environmental Protection (FDEP). 1996. Chapter 62-302 of the Florida Administration Code Rule: 62-302, 530 (45), Tallahassee, FL, USA.
- Florida Department of Environmental Protection (FDEP). 2001. Seagrass management plan for Big Lagoon and Santa Rosa Sound. Florida Department of Environmental Protection, Pensacola, FL, USA, 44 p.
- Florida Department of Environmental Protection (FDEP). 2004. Integrated water quality assessment for Florida: 2004, 305 (b) report and 303 (d) list update. Florida Department of Environmental Protection, Tallahassee, FL, USA, 190 p.
- Florida Department of Environmental Protection (FDEP). 2006. Pensacola (Group 4) verified list of impaired waters. Available on-line at URL: <http://dep.state.fl.us/water/tmdl>.
- Handley, L., D. Altsman, and R. DeMay, eds. 2007. Seagrass Status and Trends in the Northern Gulf of Mexico: 1940-2002. U.S. Geological Survey Scientific Investigations Report and U.S. Environmental Protection Agency 855-R-04-003, U.S. Geological Survey, Reston, VA, USA, 267 p.
- Heck, K.L. Jr., M.J. Sullivan, J.M. Zande and C.A. Moncreiff. 1996. An ecological analysis of seagrass meadows of the Gulf Island National Seashore - years one, two, and three: seasonal assessment and inventory, interaction studies and assessment/inventory. Marine Environment Science Consortium, University of South Alabama, Dauphin Island, AL, USA, 88 p.
- Hemminga, M.A. and C.M. Duarte. 2000. Seagrass Ecology. Cambridge University Press, Cambridge, MA, USA, 326 p.
- Hopkins, T.S. 1973. Marine Ecology in Escarosa. Coastal Coordinating Council. Contract No. CCC-03-72 and CCC-14-72. Florida Department of Natural Resources, Tallahassee, FL, USA, 100 p.
- Humm, H.J. 1964. Epiphytes of the seagrass *Thalassia testudinum* in Florida. *Bulletin of Marine Science of the Gulf and Caribbean* 14:306-341.

- Jones, W.K., J.H. Cason, and R. Bjorklund. 1992. A literature-based review of the physical, sedimentary and water quality aspects of the Pensacola Bay System. Water Resources Special Report 92 - 5, Northwest Florida Water Management District, Havana, FL, USA, 58 p.
- Lewis, T. 1986. The Pensacola Bay System. An analysis of estuarine and degradations and their relationships to land management practices. Florida Department of Community Affairs, Tallahassee, FL, USA, 132 p.
- Lewis, M.A., D.E. Weber, and B. Albrecht. 2000. Quality and toxicity in Gulf of Mexico coastal areas below ten wastewater discharges. Environmental Toxicology and Chemistry 19:192-203.
- Lewis, M.A., S.S. Foss, P.S. Harris, R.S. Stanley, and J.C. Moore. 2001. Sediment chemical contamination and toxicity associated with a coastal golf course complex. Environmental Toxicology and Chemistry 20:1390-1398.
- Lewis, M.A., R. Boustany, D. Dantin, and R. Quarles. 2002. Effect of a coastal golf course complex on water quality and nutrient bioavailability to the Gulf of Mexico. Ecotoxicology and Environmental Safety 53:154-162.
- Lewis, M.A., D. Dantin, K. Abel, and C. Lewis. 2007. Florida seagrass habitat evaluation: A comparative survey for chemical quality. Environmental Pollution 146:206-218.
- Livingston, R.J. 1971. The effects of dredging and eutropication on Mulat-Mulatto Bayou (Escambia Bay, Pensacola, Florida). Report for Florida Department of Transportation. F.S.U. Research No. 111308016, Tallahassee, FL, 40 p.
- Lores, E.M., E. Pasko, J. Patrick, R.L. Quarles, J. Campbell, and J. Macauley. 2000. Mapping and monitoring of submerged aquatic vegetation in Escambia - Pensacola Bay System, Florida. Gulf of Mexico Science 18:1-14.
- Lores, E.M. and D.T. Specht. 2001. Drought-induced decline of submerged aquatic vegetation in Escambia Bay, Florida. Gulf of Mexico Science 19:161-164.
- Macauley, J.M., J.R. Clark, and W.A. Price. 1988. Seasonal changes in the standing crop and chlorophyll content of *Thalassia testudinum* Banks ex König and its epiphytes in the northern Gulf of Mexico. Aquatic Botany 31:277-287.
- Madley, K.A., J.M. Lewis, J.D. Krolick, and E.C. Santi. 2003. Florida Marine Research Institute website seagrass pages. Available online at URL: <http://floridamarine.org>.
- McNulty, J.K., W.N. Lindall, and J.E. Sykes. 1972. Cooperative Gulf of Mexico estuaries inventory and study, Florida: phase I, area description. National Oceanic and Atmospheric Administration Technical Report Circ-368, U.S. Department of Commerce, Seattle, WA, USA, 126 p.
- McRae, G., M. Julian, and H. Brown. 2004. Florida's inshore marine monitoring and assessment program (IMAP). Annual Report, Year Five. FWRI File Code: F2248-99-A4. Fish and Wildlife Research Institute, St. Petersburg, FL, USA, 93 p.
- Moore, D.R. 1963. Distribution of the seagrass, *Thalassia*, in the United States. Bulletin of Marine Science of the Gulf and Caribbean 13:329-342.
- Murrell, M.C., R.S. Stanley, E.M. Lores, G.T. DiDonato, L.M. Smith, and D.A. Flemer. 2002. Evidence that phosphorus limits phytoplankton growth in a Gulf of Mexico estuary: Pensacola Bay, FL, USA. Bulletin of Marine Science 70:155-167.
- Murrell, M.C. and E.M. Lores. 2004. Phytoplankton and zooplankton seasonal dynamics in a subtropical estuary: importance of cyanobacteria. Journal of Plankton Research 26:371-382.
- National Oceanic and Atmospheric Administration (NOAA). 2003. NOAA coastal change analysis program (C-CAP): guidance for regional implementation. Available on-line at URL: <http://csc.noaa.gov/crs/>.
- Ridenauer, J. and C. Shambaugh. 1986. An analysis of estuarine degradations within the Pensacola Bay System and their relationship to land management practices. Florida Department of Community Affairs, Tallahassee, FL, USA, 132 p.
- Rodriguez, G. and W.C. Hunner. 1994. Water quality sampling data report of Pensacola Bay for March 1993 - February 1994. Technical File Report 94-4, Northwest Florida Water Management District, Havana, FL, USA, 392 p.
- Rogers, R.G. 1974. Seagrass revegetation in Escambia Bay, Florida. In: Proceedings of the First Annual Conference on Restoration of Coastal Vegetation in Florida, Tampa, p 21-25.
- Rogers, R.G. and F.T. Blisterfield. 1975. Loss of submerged vegetation in the Pensacola Bay system 1949 - 1974. In: R.R. Lewis, ed., Proceedings of the Second Annual Conference on Restoration of Coastal Vegetation in Florida. Hillsborough Community College, Tampa, FL, USA, p 36-50.
- Santos, S.L. and J.L. Simon. 1974. Distribution and abundance of the polychaetous annelids in a south Florida estuary. Bulletin of Marine Science 24:669-689.
- Schwenning, L., T. Bruce, and L.R. Handley. 2007. Pensacola Bay System. In: L. Handley, D. Altsman, and R. DeMay eds., Gulf of Mexico Seagrass Status and Trends. U.S. Geological Survey Scientific Investigations Report, 2006 - 5287 and U.S. Environmental Protection Agency 855-R-04-003, U.S. Geological Survey, Reston, VA, USA, p 129-142.
- Short, P.T. and S. Wylie-Echeverria. 1996. Natural and human-induced disturbance of seagrasses. Environmental Conservation. 23:17-27.
- Solis, R.S. and G.L. Powell. 1999. Hydrography, mixing characteristics and residence times of Gulf of Mexico estuaries. In: T.S. Bianchi, J.R. Pennock, and R.R. Twilley, eds., Biogeochemistry of Gulf of Mexico Estuaries. John Wiley and Sons. Inc., New York, NY, USA, p. 29-62.
- Thorpe, P., R. Bartel, P. Ryan, K. Albertson, T. Pratt, and D. Cairns. 1997. The Pensacola Bay system surface water improvement and management plan, Program Development Series 97-2. Northwest Florida Water Management District, Havana, FL, USA, 146 p.
- U.S. Department of the Interior (USDOI). 1970. Effects of pollution on water quality in Escambia River and Bay, Florida. U.S. Department of Interior. Federal Water

- Pollution Control Administration, Athens, GA, USA. 63 p.
- U.S. Environmental Protection Agency (USEPA). 1975. Environmental and Recovery Studies of Escambia Bay and the Pensacola Bay System, Florida. EPA-904/9-76-016, Region IV, Atlanta, GA, USA, 322 p.
- U.S. Environmental Protection Agency (USEPA). 1992. Status and trends of emergent and submerged vegetated habitats, Gulf of Mexico, USA. EPA-800-R92-003, Office of Water, Washington, DC, USA, 161 p.
- U.S. Environmental Protection Agency (USEPA). 2004. National Coastal Condition Report II. EPA-620/R-031002, Office of Water, Office of Research and Development, Washington, DC, USA, 286 p.
- U.S. Environmental Protection Agency (USEPA). 2005. The ecological condition of the Pensacola Bay System. EPA-620/R-05/002, Gulf Ecology Division, Gulf Breeze, FL, USA, 38 p.
- U.S. Environmental Protection Agency (USEPA). 2006. Ecological benefits assessment strategic plan. EPA-240-R-06-001, Office of the Administrator, Washington, DC, USA, 32 p.
- U.S. Geological Survey (USGS) and Gulf of Mexico Program (GOMP). 2004. Seagrass habitat in the northern Gulf of Mexico: degradation, conservation and restoration of a valuable resource. EPA-855-R-04-001, Gulf of Mexico Program Office, Stennis Space Center, MS, USA, 25 p.
- Van Breedveld, J.F. 1966. Internal Report Preliminary Algal Survey. Florida Board of Conservation Marine Laboratory, Maritime Base, St. Petersburg, FL, USA, 48 p.
- Virnstein, R.W., P.W. Mikkelsen, K.D. Cairns, and M.A. Capone. 1983. Seagrass beds versus sand bottoms: The trophic importance of their associated benthic invertebrates. *Florida Scientist* 46:363-381.
- Williams, D.R. 1981. Marine grass beds inventory, North Florida coast (October 1980). Final report prepared by the U.S. Environmental Protection Agency for the U.S. Fish and Wildlife Service for the USFWS, TS-AMD-80099. U.S.EPA, Cincinnati, OH, USA, 89 p.
- Winter, P.A. 1978. Evaluation of Seagrass Resources, Santa Rosa Island, Florida. Final Report to the Army Corps of Engineers. Florida Department of Environmental Regulation, Pensacola, FL, USA, 25 p.
- Woodward and Clyde Associates. 1997. Environmental studies in Escambia Bay and Perdido Bay systems. Final Report. Prepared for Champion International Corp., Project No. 97F299, Tallahassee, FL, USA, 210 p.
- Zieman, J.C. and R.T. Zieman. 1989. The ecology of the seagrass meadows of the west coast of Florida: a community profile. Biological Report 85 (7.25), Washington, DC, USA, 155 p.
- Zwick, P.D. and M.H. Carr. 2006. Florida 2060 – a population distribution scenario for the State of Florida. Final report, Geoplan Center, University of Florida, Gainesville, FL, USA, 25 p.
-