Gulf and Caribbean Research

Volume 23 | Issue 1

2011

Odd Association and Range Extension of *Caligus rufimaculatus* Wilson, 1905; Caligidae, Siphonostomatoida, Copepoda

G.W. Benz
Middle Tennessee State University

E.C. Greiner
University of Florida

S.R. Bowen
Savannah State University

L. Goetz
National Marine Fisheries Service

See next page for additional authors

Follow this and additional works at: <https://aquila.usm.edu/gcr>

Part of the [Marine Biology Commons](#)

Recommended Citation

Benz, G., E. Greiner, S. Bowen, L. Goetz and N. Evou. 2011. Odd Association and Range Extension of *Caligus rufimaculatus* Wilson, 1905; Caligidae, Siphonostomatoida, Copepoda. *Gulf and Caribbean Research* 23 (1): 49-53.
Retrieved from <https://aquila.usm.edu/gcr/vol23/iss1/5>
DOI: <https://doi.org/10.18785/gcr.2301.05>

This Short Communication is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in *Gulf and Caribbean Research* by an authorized editor of The Aquila Digital Community. For more information, please contact Joshua.Cromwell@usm.edu.

Odd Association and Range Extension of *Caligus rufimaculatus* Wilson, 1905; Caligidae, Siphonostomatoida, Copepoda

Authors

G.W. Benz, *Middle Tennessee State University*; E.C. Greiner, *University of Florida*; S.R. Bowen, *Savannah State University*; L. Goetz, *National Marine Fisheries Service*; and N. Evou, *National Marine Fisheries Service*

SHORT COMMUNICATION

ODD ASSOCIATION AND RANGE EXTENSION OF *CALIGUS RUFIMACULATUS* WILSON, 1905; CALIGIDAE, SIPHONOSTOMATOIDA, COPEPODA

G.W. Benz^{1*}, E.C. Greiner², S.R. Bowen³, L. Goetz⁴, and N. Evou⁴

¹Department of Biology, P.O. Box 60, Middle Tennessee State University, Murfreesboro, Tennessee 37132, USA, ²Department of Infectious Diseases and Pathology, College of Veterinary Medicine, Box 110880, University of Florida, Gainesville, Florida 32611, USA, ³Marine Science Program, Box 20467, Savannah State University, Savannah, Georgia 31404, USA, ⁴Southeast Fisheries Science Center, Panama City Laboratory, National Marine Fisheries Service, 3500 Delwood Beach Road, Panama City Beach, Florida 32408, USA; * Corresponding author, e-mail: gbenz@mtsu.edu

KEY WORDS: *Caligus rufimaculatus*, marine mammal associate, range extension, parasitic copepod, *Tursiops truncatus*

Introduction

The Caligidae (Siphonostomatoida, Copepoda) comprises more than 465 parasitic species, over 250 of which belong to *Caligus* Müller, 1785 (Boxshall and Halsey 2004). *Caligus* spp. are primarily marine; but some representatives routinely inhabit brackish or fresh water (Margolis et al. 1975). Some *Caligus* spp. have been widely reported from several oceans, others have been documented from a single location, and a few have only been collected as free-swimming individuals not associated with a host (Margolis et al. 1975). *Caligus* spp. range from being stenoxenous to euryxenous, with most species infecting actinopterygians (Actinopterygii), a much smaller group exclusively or non-exclusively infecting chondrichthyans (primarily elasmobranchs; Elasmobranchii, Chondrichthyes) (Margolis et al. 1975, Tang and Newbound 2004), and one species non-exclusively infecting an invertebrate (Ruangpan and Kabata 1984). Herein, we report a modest range extension for *Caligus rufimaculatus* Wilson, 1905 based on specimens collected in an odd association with a dead dolphin in the northern Gulf of Mexico (GOM) and we contribute new details regarding the morphology of the parasite.

Materials and Methods

Copepods were collected from a bottlenose dolphin, *Tursiops truncatus* (Montagu, 1821) (Delphinidae, Odontoceti). The dolphin (juvenile male, 199.5 cm total length) was discovered dead on 26 December 2005, partially submerged under a dock in shallow water at the southern edge of Choctawhatchee Bay in Miramar Beach, Florida. Responders from the Southeast Fisheries Center (Panama City Beach, FL) collected the carcass on 30 December 2005 (the only dead animal noted in the area at that time) and moved it to a refrigerator. Copepods were collected during necropsy on 31 December 2005 from the skin of the ventral caudal region of the body and fixed in 10% formalin. It is not known

if the copepods were alive at the time the dolphin was removed from the water and no lesion was grossly apparent where the copepods were attached. This dolphin stranded during a brevetoxin-associated unusual mortality event in which 93 dolphin strandings were documented from 19 September 2005 to 30 April 2006 (Gaydos et al.¹). For taxonomic study, copepods were cleared in lactic acid into which a small amount of lignin pink had been dissolved. Fine pins mounted in the tips of thin wooden dowels were used to dissect copepods, and the wooden slide technique of Humes and Gooding (1964) facilitated the study of intact specimens and dissected appendages. Copepods were examined using bright-field microscopy. Measurements were made using a stage micrometer and illustrations were made using a camera lucida. Three specimens (2 females, 1 male) were studied using a scanning electron microscope (SEM) after routine preparation (CO₂ critical point drying, gold-palladium sputter coating, mounting on metal stubs with 2-sided sticky tape). Copepod terminology used herein conforms mostly to that of Boxshall and Halsey (2004) and host nomenclature and systematics follows Compagno (2005) for members of Chondrichthyes, FishBase (Froese and Pauly 2008) for other fishes, Reynolds III and Powell (2002) for manatees, and Jefferson et al. (1993) for other marine mammals. Copepod vouchers remain in the personal collection of the senior author.

¹Gaydos, J.K., N. Barrios, G.D. Bossart, S. Bowen, K. Evans, R. Ewing, M.L. Fleetwood, L. Flewelling, R. Hardy, C. Heil, C. Johnson, W. Jones, J. Kucklick, J. Landsberg, J. Litz, T.A. Leighfield, C. Lockyer, B. Mase, W. Noke, P. Rosel, D.S. Rotstein, T. Rowles, L. Schwacke, T. Spradlin, M. Stolen, M.J. Twiner, and F.M. Van Dolah. In review. Brevetoxin-associated bottlenose dolphin (*Tursiops truncatus*) unusual mortality events in the Florida Panhandle: 1999–2000, 2004, and 2005–2006. United States National Oceanic and Atmospheric Association Technical Memorandum, National Marine Fisheries Service—OPR.

Results and Discussion

About 30 copepods were collected from the dead dolphin and 24 individuals (6 ovigerous females, 11 non-ovigerous adult females, and 7 adult males) were each identified as *C. rufimaculatus* Wilson, 1905 (Figure 1A) based on information in Cressey (1991) and Suárez–Morales et al. (2003). In their remarks regarding *Caligus amblygenitalis* Pillai, 1961, Ho and Lin (2004) noted that few *Caligus* spp. possess an accessory process on all three outer elements of the second exopodal segment of leg 1. They included *C. rufimaculatus* as one of said species based on the illustration of leg 1 provided by Cressey (1991). However, in his text description of *C. rufimaculatus*, Cressey (1991) stated that the outermost element did not exhibit an accessory process. In his original description of *C. rufimaculatus*, Wilson (1905) did not report any of the three outermost elements (referred to by Wilson as claws) as possessing an accessory process and most recently, Suárez–Morales et al. (2003) reported that their non-type exemplars exhibited an accessory process on elements 2 and 3 (referred to by those authors as spines) among the three outer elements. Our observations corroborate the illustration of leg 1 (but not the text description) of Cressey (1991) in that we observed adult females and adult males to possess three outer elements each with an accessory process (Figure 1B). Ho and Lin (2004) also remarked that *C. amblygenitalis* can be distinguished from *C. rufimaculatus* in that the former species possesses a large patch of spinules ventrally on the basis of leg 2. Here it is noted that there seems little doubt, based on their detailed description of *C. amblygenitalis*, that Ho and Lin (2004) mistakenly remarked that the aforementioned patch of spinules is found on leg 2, as clearly their text description and illustrations establish the patch as residing on the basis of leg 1. Our observations of the adult female and male of *C. rufimaculatus* revealed tiny cuticular bumps on the ventral surface of the basis of leg 1 (Figure 1C) and no cuticular projections on the ventral surface of the basis of leg 2. These bumps (not reported by others) seem smaller than the homologous spinules illustrated by Ho and Lin (2004). Lastly, Ho and Lin (2004) remarked that another feature distinguishing *C. amblygenitalis* from *C. rufimaculatus* is an “unipectinate outer spine with strong teeth” on the second segment of leg 2. Our observations of the homologous feature of *C. rufimaculatus* revealed it to be a spine with a marginal membrane on each side, described as a “fringes” by Cressey (1991).

Few copepod species infect or otherwise associate with mammals (Boxshall and Halsey 2004) and we believe the following to be a complete accounting of said species: *Penella balaenopterae* Korea and Danielson, 1877, (Siphonostomatoida) is a mesoparasite that infects mysticets (Mysticeti) and odontocets (Odontoceti) (Hogans 1987) and the northern elephant seal, *Mirounga angustirostris*, (see Dailey et al. 2002); *Balaenophilus unisetus* Aurivillius, 1897

Figure 1. *Caligus rufimaculatus* collected from a dead bottlenose dolphin, *Tursiops truncatus*, in Choctawhatchee Bay, Florida. A. Adult female (left) and adult male (right). B. Tip of leg 1 exopod (adult female) showing three outermost elements (I, II, III) and their accessory processes (one per element; Ia, IIa, IIIa). Note: the base of seta IV (not visible in image) lies just medial to the base of element III. C. Ventral view of the basipod of leg 1 showing a patch of tiny cuticular bumps.

TABLE 1. Published host fish records for *Caligus rufimaculatus* Wilson, 1905.

Class	Order	Family	Species	Location	Attachment site on host	Reference
Chondrichthyes	Rajiformes	Rhinobatidae	<i>Rhinobatos lentiginosus</i> *	in or about Lemon Bay, FL	body surface	Bere 1936
		Myliobatidae	<i>Aetobatus narinari</i> *	in or about Lemon Bay, FL	body surface	Bere 1936
		Mobulidae	"devilfish" ¹	in or about Lemon Bay, FL	about mouth	Bere 1936
Actinopterygii	Mugiliformes	Mugilidae	<i>Mugil cephalus</i>	not reported in or about Lemon Bay, FL Biscayne Bay, FL Sapelo Island, GA	not reported mouth, body surface skin gills, body surface	Wilson 1905, 1932 Bere, 1936 Skinner 1975 Rawson 1977 ²
	Beloniformes	Belonidae	<i>Strongylura timucu</i>	in or about Lemon Bay, FL	body surface	Bere 1936
	Cyprinodontiformes	Fundulidae	<i>Fundulus heteroclitus</i>	Woods Hole, MA about Beaufort, NC	body surface body surface	Wilson 1905, 1932 Wilson 1908
			<i>F. majalis</i>	Woods Hole, MA	body surface	Wilson 1905, 1932
	Perciformes	Serranidae	<i>Centropristis striata</i> *	about Placida, FL	not reported	Cressey 1991
		Pomatomidae	<i>Pomatomus saltatrix</i>	in or about Lemon Bay, FL Maricá, Rio de Janeiro, Brazil	mouth body surface	Bere 1936 Luque et al. 1998
		Carangidae	<i>Oligoplites palometa</i>	Sepetiba Bay, Brazil	gills	Takemoto & Luque 2002
			<i>O. saliens</i>	Sepetiba Bay, Brazil	gills	Takemoto & Luque 2002
			<i>O. saurus</i>	in or about Lemon Bay, FL	mouth, body surface	Bere 1936
		Lutjanidae	<i>Lutjanus synagris</i>	about Placida, FL	not reported	Cressey 1991
		Gerreidae	<i>Eucinostomus gula</i>	about Placida, FL	not reported	Cressey 1991
			<i>E. pseudogula</i> ³	in or about Lemon Bay, FL	body surface	Bere 1936
		Haemulidae	<i>Haemulon plumieri</i> *	about Placida, FL	not reported	Cressey 1991
			<i>Orthopristis chrysoptera</i>	about Placida, FL	not reported	Cressey 1991
		Sparidae	<i>Diplodus holbrookii</i> * ⁴	about Placida, FL	not reported	Cressey 1991
			<i>Lagodon rhomboides</i>	about Placida, FL	not reported	Cressey 1991
		Scaridae	<i>Nicholsina usta</i>	about Placida, FL	not reported	Cressey 1991
		Ephippidae	<i>Chaetodipterus faber</i>	in or about Lemon Bay, FL	body surface	Bere 1936
		Trichiuridae	<i>Trichiurus lepturus</i>	on North American coast	not reported	Wilson 1905 ⁵
	Tetraodontiformes	Monacanthidae	<i>Stephanolepis hispidus</i> *	about Placida, FL	not reported	Cressey 1991
		Ostraciidae	<i>Acanthostracion quadricornis</i> *	about Placida, FL	not reported	Cressey 1991

* indicates that the host name used in that record was herein changed to conform to nomenclature in FishBase (Froese and Pauly 2008).

¹ *Mobula hypostoma* noted by author as the most probable host species.

² Results published in Rawson (1977) represent a portion of the dissertation results presented in Rawson 1973.

³ Host binomen not listed in FishBase (Froese and Pauly 2008); also referred to as "the mojarra" by author (Bere, 1936; p. 582-583).

⁴ Family entry of host in appendix of Cressey (1991; p. 18) signals an error regarding host name in text (p. 14).

⁵ Wilson's (1905; p. 573) identification reassigned as *C. rufimaculatus* by Parker (1969).

(Harpacticoida) inhabits the baleen of several rorqual species (*Balaenoptera*, Balaenopteridae) (see Bannister and Grindley 1966); *B. manatorum* (Ortiz, Lalana, and Torres—Fundora, 1992) inhabits the skin of the Antillean manatee, *Trichechus manatus manatus*, (see Suárez—Morales 2007); *Harpacticus pulex* Humes, 1964 (Harpacticoida) was discovered on sloughed skin of a captive—held bottlenose dolphin and captive—held Florida manatee, *T. m. latirostris*, (see Humes 1964); and *Caligus elongatus* Nordmann, 1832 (Siphonostomatoida) was reported attached to a dead bottlenose whale, *Hyperoodon ampullatus*, (see O'Reilly 1998). This report adds another *Caligus* sp. to the list of copepods associated with mammals; and for completeness we add that occurrences of caligids temporarily attaching to humans have been reported (Kurochkin and Kazachenko 1975, as reported by Ruangpan and Kabata 1984, O'Reilly 1998). Cases of copepods attaching to humans aside, it is notable that amongst the aforementioned copepods, only *C. elongatus* and *C. rufimaculatus* have been reported from mammals as well as representatives of other classes. Considering numbers of species and phylogenetic scope, *C. elongatus* and *C. rufimaculatus* have each been reported from a numerous and diverse variety of hosts (Margolis et al. 1975; Table 1). However, whether being euryxenous regarding fish hosts is a prerequisite for their ability to associate with marine mammals is unknown and certainly there are other euryxenous *Caligus* spp. that are not known to associate with mammals (Margolis et al. 1975). In the present case, no data support or refute the notion that the specimens of *C. rufimaculatus* were actually feeding on the bottlenose dolphin, and whether or not the copepods had attached to the dolphin prior to its death likewise remains unknown. A lack of dead fish in the immediate vicinity of the dead dolphin did not support a notion that the copepods had abandoned moribund fish hosts and settled on the dolphin; however, red tide related fish kills had occurred in Choctawhatchee Bay for months before and during the time of the dolphin stranding. This case is more curious because it involved about 30 copepods, unlike the case reported by

O'Reilly (1998) of two *C. elongatus* attached to a dead whale.

The known geographic distribution of *C. rufimaculatus* has expanded significantly since the species was first discovered (Table 1). Prior to this report the species was considered to patchily inhabit western North Atlantic coastal waters from Woods Hole, Massachusetts to Biscayne Bay in southern Florida (Wilson 1905, 1908; Rawson 1973, 1977; Skinner 1975), GOM coastal waters off southern Florida and the Yucatán Peninsula, Mexico (Bere 1936, Cressey 1991, Suárez—Morales et al. 2003), and western South Atlantic waters off Maricá and in Sepetiba Bay, Rio de Janeiro, Brazil (Luque et al. 1998, Takemoto and Luque 2002). This report extends the species range to include the northern GOM and in doing so it negates the comment of Cressey (1991) that *C. rufimaculatus* does not occur in said region. *Caligus rufimaculatus* has been collected from at least 22 species of fishes, together representing 18 families and 6 orders (Table 1). The phylogenetic scope of these hosts (elasmobranchs to members of the Tetraodontiformes) is remarkable, as is their ecological scope, ranging from small, shallow—water, and euryhaline species (e.g., *Fundulus heteroclitus*) to larger constantly swimming and wide—ranging nearshore species (e.g., *Pomatomus saltatrix*), as well as a species known to be capable of inhabiting relatively deep water (*Trichiurus lepturus*) (Froese and Pauly 2008). In addition, *C. rufimaculatus* is capable of attaching to a wide array of host surfaces, from smooth and scaleless skin (e.g., *Rhinobatos lentiginosus*, *Trichiurus lepturus*, *Tursiops truncatus*) to heavily scaled skin (*Mugil cephalus*) to hard plated skin (*Acanthostracion quadricornis*) to gills (*M. cephalus*, *Oligoplites palometa*) (Table 1). Its ability to move about freely away from a host and associate with many species would seem to predispose *C. rufimaculatus* to be a sea louse capable of negatively impacting a wide variety of aquaculture operations. Yet we are unaware of any such reports. Based on its aforementioned abilities, its known distribution within the GOM, and the general pattern of water flow within the GOM, we expect that future records of *C. rufimaculatus* will document much more of the GOM as part of its range.

Acknowledgments

We thank C. Levins (Panama City Laboratory, Southeast Fisheries Science Center, NOAA) for providing assistance with the stranding response and E. R. Salmon (Middle Tennessee State University) for leading activities associated with the SEM examination.

Literature Cited

- Bannister, J.L. and J.R. Grindley. 1966. Notes on *Balaenophilus unisetus* P. O. C. Aurivillius, 1879, and its occurrence in the southern hemisphere (Copepoda, Harpacticoida). *Crustaceana* 10:296–302.
- Bere R. 1936. Parasitic copepods from Gulf of Mexico fish. *The American Midland Naturalist* 17:577–625.
- Boxshall, G.A. and S.H. Halsey. 2004. An introduction to copepod diversity. The Ray Society, London, UK, 966 p. [in 2 parts: Part 1 pages 1–421, Part 2 pages 423–966]

- Compagno, L.J.V. 2005. Checklist of living Chondrichthyes. In: W.C. Hamlett, ed. Reproductive biology and phylogeny of Chondrichthyes. Science Publishers, Inc., Enfield, New Hampshire, USA, p. 503–549.
- Cressey, R.F. 1991. Parasitic copepods from the Gulf of Mexico and Caribbean Sea, III: *Caligus*. Smithsonian Contributions to Zoology 497:1–53.
- Dailey M.D., M. Haulena, and J. Lawrence. 2002. First report of a parasitic copepod (*Pennella balaenopterae*) infestation in a pinniped. Journal of Zoo and Wildlife Medicine 33:62–65.
- Froese, R. and D. Pauly, eds. 2008. FishBase. World Wide Web electronic publication. www.fishbase.org. (viewed on 09/01/2010).
- Ho, J.-S. and C.-L. Lin. 2004. Sea lice of Taiwan. The Sueichan Press, Keelung, Taiwan, 388 p.
- Hogans W.E. 1987. Morphological variation in *Pennella balenoptera* and *P. filosa* (Copepoda: Pennellidae) with a review of the genus *Pennella* Oken, 1816 parasitic on Cetacea. Bulletin of Marine Science 40:442–453.
- Humes, A.G. 1964. *Harpacticus pulex*, a new species of copepod from the skin of a porpoise and a manatee in Florida. Bulletin of Marine Science of the Gulf and Caribbean 14:517–528.
- Humes, A.G. and R.U. Gooding. 1964. A method for studying the external anatomy of copepods. Crustaceana 6:238–240.
- Jefferson, T.A., S. Leatherwood, and M.A. Webber. 1993. Marine mammals of the world. Food and Agriculture Organization of the United Nations, Rome, Italy, 320 p.
- Kurochkin, Y.V. and V.N. Kazachenko. 1975. On cases of attachment of parasitic caligids and argulids to human skin during submersion in water. Investiya TINRO 98:257–258. [In Russian; as cited by Ruangpan and Kabata, 1984]
- Luque, J.L., N.D. Chaves, and A.D. Cezar. 1998. Novos registros de copépodos calogóideos parasitos de peixes marinhos do Brasil. Nauplius, Rio Grande 6:9–16.
- Margolis, L., Z. Kabata, and R.R. Parker. 1975. Catalogue and synopsis of *Caligus*, a genus of Copepoda (Crustacea) parasitic on fishes. Bulletin of the Fisheries Research Board of Canada 192:1–117.
- O'Reilly, M. 1998. Whale-lice (Amphipoda: Cyamidae) and sea lice (Copepoda: Caligidae) from stranded whales in the Firth of Forth. Glasgow Naturalist 23:24–26.
- Parker, R.R. 1969. Validity of the binomen *Caligus elongatus* for a common parasitic copepod formerly misidentified with *Caligus rapax*. Journal of the Fisheries Research Board of Canada 26:1013–1035.
- Rawson Jr., M.V. 1973. The development and seasonal abundance of the parasites of striped mullet, *Mugil cephalus* L., and mummichogs, *Fundulus heteroclitus* (L.). Ph.D. thesis. University of Georgia, Athens, Georgia, USA, 100 p.
- Rawson Jr., M.V. 1977. Population biology of parasites of striped mullet, *Mugil cephalus* L. Crustacea. Journal of Fish Biology 10:441–451.
- Reynolds III, J.E. and J.A. Powell. 2002. Manatees: *Trichechus manatus*, *T. senegalensis*, and *T. inunguis*. In: W.F. Perrin, B. Würsig, and J.G.M. Thewissen, eds. Encyclopedia of marine mammals. Academic Press, San Diego, California, USA, p.709–720.
- Ruangpan, L. and Z. Kabata. 1984. An invertebrate host for *Caligus* (Copepoda: Caligidae)? Crustaceana 47:219–220.
- Skinner, R. 1975. Parasites of the striped mullet, *Mugil cephalus*, from Biscayne Bay, Florida, with descriptions of a new genus and three new species of trematodes. Bulletin of Marine Science 25:318–345.
- Suárez-Morales, E. 2007. On the taxonomic status of *Balaenophilus umimegacolus* Ogawa, Matsuzaki & Misaki, 1997 (Copepoda, Harpacticoida), a symbiont of manatees and sea turtles. Crustaceana 80:1145–1150.
- Suárez-Morales, E., I-H. Kim, and J.B. Escamilla. 2003. Illustrated record and complementary description of *Caligus rufimaculatus* Wilson (Copepoda: Siphonostomatoida) from Mexico. Caribbean Journal of Science 39:151–154.
- Takemoto, R.M. and J.L. Luque. 2002. Parasitic copepods on *Oligoplites* spp. (Osteichthyes, Carangidae) from the Brazilian coastal zone, with the redescription of *Tuxophorus caligodes* Wilson, 1908 (Siphonostomatoida, Tuxophoridae). Acta Scientiarum, Maringá 24:481–487.
- Tang, D. and D.R. Newbound. 2004. A new species of copepod (Siphonostomatoida: Caligidae) parasitic on the tiger shark *Galeocerdo cuvier* (Péron & Lesueur) from western Australian waters. Systematic Parasitology 58:69–80.
- Wilson, C.B. 1905. North American parasitic copepods belonging to the family Caligidae. Part 1. The Caliginæ. Proceedings of the United States National Museum 28:479–672.
- Wilson, C.B. 1908. North American parasitic copepods: new genera and species of Caliginæ. Proceedings of the United States National Museum 33:593–627.
- Wilson, C.B. 1932. The copepods of the Woods Hole region, Massachusetts. Bulletin of the United States National Museum 158:1–635.