Gulf and Caribbean Research

Volume 24 | Issue 1

2012

Evidence of Spawning Capable Tarpon (*Megalops atlanticus*) Off the Louisiana Coast

William Stein III
University of New Orleans

Nancy J. Brown-Peterson
University of Southern Mississippi, nancy.brown-peterson@usm.edu

James S. Franks
University of Southern Mississippi, jim.franks@usm.edu

Martin T. O'Connell
University of New Orleans

Follow this and additional works at: <https://aquila.usm.edu/gcr>

Part of the [Marine Biology Commons](#)

Recommended Citation

Stein, W. III, N. J. Brown-Peterson, J. S. Franks and M. T. O'Connell. 2012. Evidence of Spawning Capable Tarpon (*Megalops atlanticus*) Off the Louisiana Coast. *Gulf and Caribbean Research* 24 (1): 73-74.
Retrieved from <https://aquila.usm.edu/gcr/vol24/iss1/9>
DOI: <https://doi.org/10.18785/gcr.2401.09>

This Short Communication is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in *Gulf and Caribbean Research* by an authorized editor of The Aquila Digital Community. For more information, please contact Joshua.Cromwell@usm.edu.

SHORT COMMUNICATION

EVIDENCE OF SPAWNING CAPABLE TARPON (*MEGALOPS ATLANTICUS*) OFF THE LOUISIANA COAST

William Stein III^{1*}, Nancy J. Brown-Peterson², James S. Franks³, and Martin T. O'Connell¹

¹Nekton Research Laboratory, Pontchartrain Institute for Environmental Sciences, University of New Orleans, 2000 Lakeshore Dr., New Orleans, LA 70148, USA; ²Department of Coastal Sciences, The University of Southern Mississippi, 703 East Beach Dr., Ocean Springs, MS 39564, USA; and ³Gulf Coast Research Laboratory, The University of Southern Mississippi, 703 East Beach Dr., Ocean Springs, MS 39564, USA; *Corresponding author email: wstein@uno.edu

KEY WORDS: *Megalops atlanticus*, Northern Gulf of Mexico, Spawning

INTRODUCTION

Despite the fact that the tarpon (*Megalops atlanticus*) is a popular sport fish in the Gulf of Mexico (GOM; Ault et al. 2008), little is known of its spawning behavior. Spawning *M. atlanticus* have never been documented and fertilized eggs have not been observed *in situ* (Ault et al. 2008). While it has been suggested that adult *M. atlanticus* move to deep water off the southwest coast of Florida and into the Yucatan Channel to spawn, the actual locations of spawning grounds remain unknown (Smith 1980, Crabtree et al. 1995, Crabtree et al. 1997). Distribution patterns of larvae (leptocephali) have served as the basis for most of what has been inferred about the spawning areas (Smith 1980, Crabtree et al. 1997). For example, leptocephali were collected from the southwestern GOM, the Yucatan Channel, and along the west coast of Florida, and based on their size, it was assumed that *M. atlanticus* spawned in nearby areas (Smith 1980).

Histological examination of gonads has also been used to estimate the location of *M. atlanticus* spawning habitat. Females collected from the Florida Keys and Boca Grande Pass off the west coast of Florida and contained ovaries with post ovulatory follicles (POF) and advanced vitellogenic oocytes, suggesting *M. atlanticus* spawn in this region from April through July (Crabtree et al. 1997). Examination of gonads from *M. atlanticus* caught off the coast of equatorial Ceara State, Brazil suggested that spawning occurs there from October through January (de Menezes and Paiva 1966). We report here the first evidence of spawning capable *M. atlanticus* off the coast of Louisiana in the northern GOM based on histological examination of gonads.

MATERIALS AND METHODS

Two large, sexually mature *M. atlanticus* (one female, one male) were collected by anglers about 30 km south of Venice, Louisiana on 2 July 2011. A second male was captured by anglers on 28 July 2011 from the same area. Fish were weighed (kg) and measured (mm fork length, FL) and gonadal tissue from each specimen was removed, weighed, and

fixed whole in 10% neutral buffered formalin within 12 h of capture. Gonadal tissue was processed following standard histological techniques, embedded in paraffin, sectioned at 4 µm and stained with hematoxylin and eosin. Reproductive phases and gamete stages were determined following Brown-Peterson et al. (2011).

RESULTS AND DISCUSSION

The female *M. atlanticus* weighed 56.8 kg and was 1778 mm FL. The first male weighed 55.5 kg and was 1676 mm FL; the second male weighed 56.6 kg and was 1698 mm FL. These fish were within the same size ranges as reported for spawning *M. atlanticus* from Brazil and Florida (de Menezes and Paiva 1966, Crabtree et al. 1997).

The female was classified as spawning capable based on the presence of both late vitellogenic oocytes and 24 h POF in the ovaries at the time of collection (Figure 1A). The warm water at the time of specimen collection in combination with less than optimal fixation resulted in rapid degradation of the POF observed. However, these POFs appear similar to 24 h POF from other species collected at similar water temperatures (Brown-Peterson et al. 2011) and provide evidence that this female spawned 24 h or less prior to capture. While the ovary was dominated by tertiary vitellogenic oocytes (Vtg3), there was evidence of asynchronous oocyte development because secondary vitellogenic (Vtg2) were also present (Figure 1A). The two males were also both classified as spawning capable. Testes were full of spermatozoa but there was no active spermatogenesis occurring (Figure 1B), suggesting the males were at the end of the reproductive season but still capable of releasing spermatozoa. These specimens were collected near the end of the reported spawning season for *M. atlanticus* in Florida (Crabtree et al. 1997).

The collection of both male and female *M. atlanticus* in the spawning capable phase suggests that *M. atlanticus* may be spawning off Louisiana and represents the first evidence that this species appears to be reproducing in the northern

Figure 1. Histological images of gonadal tissue in the spawning capable phase from *M. atlanticus* collected off the coast of Louisiana on 2 July 2011. **A.** Female. **B.** Male. Key: PG—primary growth oocyte; POF—24 h postovulatory follicle; Vtg2—secondary vitellogenic oocyte; Vtg3—tertiary vitellogenic oocyte; GE—germinal epithelium; SZ—spermatozoa.

GOM. Juvenile *M. atlanticus* have been reported from Mississippi coastal locations by Franks (1970), Overstreet (1974) and Schofield et al. (2007). Various suggestions have been made to account for the presence of juvenile *M. atlanticus* on the coasts of Louisiana¹ and Mississippi² (Franks 1970, Overstreet 1974,

Schofield et al. 2007). It is possible that these juveniles are the product of local spawning activity based on evidence provided here. Additional collections of adult *M. atlanticus* during the spring and summer from the northern GOM would help elucidate reproductive activity of this species.

ACKNOWLEDGEMENTS

We thank M. Mullen, USM, for histological preparations. We are grateful to members of the Louisiana Tarpon Club, Louisiana Council of Underwater Dive Clubs, and Hell Divers Dive Club for providing specimens and to J. Pituch, Louisiana Department of Wildlife and Fisheries Biologist, for assistance with gonad sample collection.

LITERATURE CITED

- Ault, J.S., R. Humston, M.F. Larkin, E. Perusquia, N. . Farmer, J. Luo, N. Zurcher, S.G. Smith, L.R. Barbieri, and J.M. Posada. 2008. Population dynamics and resource ecology of Atlantic tarpon and bonefish. In: J. S. Ault, ed. *Biology and Management of the World Tarpon and Bonefish Fisheries*. CRC Press, Boca Raton, FL, USA, p. 217-258.
- Brown-Peterson, N.J., D.M. Wyanski, F. Saborido-Rey, B.J. Macewicz, and S.K. Lowerre-Barbieri. 2011. A standardized terminology for describing reproductive development in fishes. *Marine and Coastal Fisheries* 3:52-70.
- Crabtree, R.E., E.C. Cyr, and J.M. Dean. 1995. Age and growth of tarpon, *Megalops atlanticus*, from South Florida waters. *Fishery Bulletin*, U.S. 93:619-628.
- Crabtree, R.E., E.C. Cyr, D.C. Chacon-Chaverri, W.O. McLarney, and J.M. Dean. 1997. Reproduction of tarpon, *Megalops atlanticus*, from Florida and Costa Rican waters and notes on their age and growth. *Bulletin of Marine Science* 61:271-285.
- de Menezes, M.F. and M.P. Paiva. 1966. Notes on the biology of tarpon, *Megalops atlanticus* (Cuvier and Valenciennes), from coastal waters of Ceara State, Brazil. *Arquivos da Estação de Biologia Marinha da Universidade Federal do Ceará* 6:83-98.
- Franks, J.S. 1970. An investigation of the fish population within the inland waters of Horn Island, Mississippi, a barrier island in the northern Gulf of Mexico. *Gulf Research Reports* 3:3-104.
- Overstreet, R.M. 1974. An estuarine low-temperature fish-kill in Mississippi, with comments on restricted necropsies. *Gulf Research Reports* 4:328-350.
- Schofield, P.J., W.T. Slack, M.S. Peterson, and D.R. Gregoire. 2007. Assessment and control of an invasive aquaculture species: an update on Nile tilapia (*Oreochromis niloticus*) in coastal Mississippi after Hurricane Katrina. *Southeastern Fishes Council, Proceedings* 49:9-15.
- Smith, D.G. 1980. Early larvae of the tarpon, *Megalops atlanticus* Valenciennes (Pisces: Elopidae), with notes on spawning in the Gulf of Mexico and the Yucatan Channel. *Bulletin of Marine Science* 30:136-141.

¹Stein, W., J. McKenzie, and O. T. Lorenz. 2011. Potential juvenile tarpon (*Megalops atlanticus*) nursery habitats in southeastern Louisiana. The 31st Annual Meeting of the Louisiana Chapter of the American Fisheries Society, 27-28 January 2011, Lafayette, Louisiana.

²Franks, J. S., P. Grammer, J. Ballard, M. Buchanan, and G. Gray. 2009. Juvenile tarpon (*Megalops atlanticus*) in Mississippi coastal waters: short-term event or long-term trend? 35th Annual Meeting of the Mississippi Chapter of the American Fisheries Society, 11-13 February 2009, Biloxi, Mississippi.