

3-26-2015

Draft Genome Sequence of the Phenazine-Producing *Pseudomonas fluorescens* Strain 2-79

Kai Neemann
Georg-August Universität

Susanna A. Braus-Stromeyer
Georg-August Universität

Andrea Thuermer
Georg-August Universität

Rolf Daniel
Georg-August Universität

Dmitri V. Mavrodi
University of Southern Mississippi, dmitri.mavrodi@usm.edu

See next page for additional authors

Follow this and additional works at: https://aquila.usm.edu/fac_pubs



Part of the [Biology Commons](#)

Recommended Citation

Nesemann, K., Braus-Stromeyer, S. A., Thuermer, A., Daniel, R., Mavrodi, D. V., Thomashow, L. S., Weller, D. M., Braus, G. H. (2015). Draft Genome Sequence of the Phenazine-Producing *Pseudomonas fluorescens* Strain 2-79. *Genome Announcements*, 3(2), 1-2.
Available at: https://aquila.usm.edu/fac_pubs/15181

This Article is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in Faculty Publications by an authorized administrator of The Aquila Digital Community. For more information, please contact Joshua.Cromwell@usm.edu.

Authors

Kai Nesemann, Susanna A. Braus-Stromeyer, Andrea Thuermer, Rolf Daniel, Dmitri V. Mavrodi, Linda S. Thomashow, David M. Weller, and Gerard H. Braus

Draft Genome Sequence of the Phenazine-Producing *Pseudomonas fluorescens* Strain 2-79

Kai Neseemann,^a Susanna A. Braus-Stromeier,^a Andrea Thuermer,^b Rolf Daniel,^b Dmitri V. Mavrodi,^c Linda S. Thomashow,^d David M. Weller,^d Gerhard H. Braus^a

Department of Molecular Microbiology and Genetics, Institute of Microbiology and Genetics, Georg-August Universität, Göttingen, Germany^a; Department of Genomic and Applied Microbiology and Göttingen Genomics Laboratory, Institute of Microbiology and Genetics, Georg-August Universität, Göttingen, Germany^b; Department of Biological Sciences, University of Southern Mississippi, Hattiesburg, Mississippi, USA^c; USDA-ARS Root Disease and Biological Control Research Unit, Washington State University, Pullman, Washington, USA^d

***Pseudomonas fluorescens* strain 2-79, a natural isolate of the rhizosphere of wheat (*Triticum aestivum* L.), possesses antagonistic potential toward several fungal pathogens. We report the draft genome sequence of strain 2-79, which comprises 5,674 protein-coding sequences.**

Received 4 February 2015 Accepted 18 February 2015 Published 26 March 2015

Citation Neseemann K, Braus-Stromeier SA, Thuermer A, Daniel R, Mavrodi DV, Thomashow LS, Weller DM, Braus GH. 2015. Draft genome sequence of the phenazine-producing *Pseudomonas fluorescens* strain 2-79. *Genome Announc* 3(2):e00130-15. doi:10.1128/genomeA.00130-15.

Copyright © 2015 Neseemann et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Gerhard H. Braus, gbraus@gwdg.de.

The concentration and composition of antibiotic-producing, root-colonizing organisms are important factors that partially determine the suppressiveness of soils toward certain soil-borne diseases (1).

Fluorescent pseudomonads play a major role in suppressing take-all disease of wheat caused by the fungal pathogen *Gaeumannomyces graminis* var. *tritici* (Sacc.) (2). In 1979, Weller and Cook isolated bacteria from roots of wheat plants grown in take-all suppressive soils in Washington state, USA (3). *Pseudomonas fluorescens* 2-79 (NRRL B-15132) was characterized as a strong biological control agent suppressing *G. graminis* in vitro and in planta. Wheat plants infected with *G. graminis* var. *tritici* and additionally treated with *P. fluorescens* 2-79 resulted in taller plants, more heads, and fewer symptoms of root disease compared to the control plants without bacterial treatment. Bacterial treatment could increase the yield up to 147% in soils fumigated with methyl bromide and up to 27% in natural soils (3). *P. fluorescens* 2-79 produces phenazines, which represent a diverse chemical group of nitrogen-containing heterocyclic pigments possessing broadly inhibitory properties toward bacteria and fungi (4). Phenazines undergo redox reactions with NADH/NADPH, leading to an increase of toxic superoxide radicals and hydrogen peroxide in the target cells (5). Mavrodi et al. investigated the biosynthesis pathway of phenazines in *P. fluorescens* 2-79 (6).

Genomic DNA of *P. fluorescens* 2-79 was isolated by using the MasterPure Complete DNA and RNA purification kit (Epicentre, Madison, WI, USA). A shotgun sequencing library was generated employing the Nextera DNA sample preparation kit following the manufacturer's instructions. The whole genome of *P. fluorescens* 2-79 was sequenced with the Genome Analyzer IIx (Illumina, San Diego, CA, USA). In total, 8.5 million paired-end reads of 112 bp were generated. *De novo* assembly of all shotgun reads using SPAdes version 3.0.0 (7) resulted in 143 contigs >3 kb and 123-fold coverage. The draft genome sequence comprises 6.4 Mb and a GC content of 59.83%. Ge-

nome annotation was performed by using Prokka (8). The draft genome harbored 1 rRNA cluster, 47 tRNA genes, 4,286 protein-encoding genes with function prediction, and 1,388 genes coding for hypothetical proteins.

Proteins involved in secondary metabolism were analyzed. The gene *hcnA* (GenBank accession no. 15560558) involved in HCN synthesis and the phenazine operon (GenBank accession no. L48616.1) are present in *P. fluorescens* 2-79. The gene *phlD* (GenBank accession no. 15563828) necessary for the synthesis of 2,4-diacetylphloroglucinol (DAPG) is absent in 2-79.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number [JXCQ000000000](https://www.ncbi.nlm.nih.gov/nuccore/JXCQ000000000). The version described in this paper is the first version, [JXCQ010000000](https://www.ncbi.nlm.nih.gov/nuccore/JXCQ010000000).

ACKNOWLEDGMENTS

This work was supported by the Federal Ministry of Education and Research (BMBF) BioFung project and the DFG through grants awarded to G.H.B. We also are grateful for support provided by USDA-NRI grant no. 2011-67019-30212 from the USDA-NIFA Soil Processes program.

REFERENCES

- Haas D, Défago G. 2005. Biological control of soil-borne pathogens by fluorescent pseudomonads. *Nat Rev Microbiol* 3:307–319. <http://dx.doi.org/10.1038/nrmicro1129>.
- Cook RJ, Rovira AD. 1976. The role of bacteria in the biological control of *Gaeumannomyces graminis* by suppressive soils. *Soil Biol Biochem* 8:269–273. [http://dx.doi.org/10.1016/0038-0717\(76\)90056-0](http://dx.doi.org/10.1016/0038-0717(76)90056-0).
- Weller DM, Cook RJ. 1983. Suppression of take-all of wheat by seed treatments with fluorescent pseudomonads. *Phytopathology* 73:463–469. <http://dx.doi.org/10.1094/Phyto-73-463>.
- Mavrodi DV, Blankenfeldt W, Thomashow LS. 2006. Phenazine compounds in fluorescent *Pseudomonas* spp. biosynthesis and regulation. *Annu Rev Phytopathol* 44:417–445. <http://dx.doi.org/10.1146/annurev-phyto.44.013106.145710>.
- Hassett DJ, Woodruff WA, Wozniak DJ, Vasil ML, Cohen MS, Ohman DE. 1993. Cloning and characterization of the *Pseudomonas aeruginosa* *sodA* and *sodB* genes encoding manganese- and iron-cofactored superoxide

- dismutase: demonstration of increased manganese superoxide dismutase activity in alginate-producing bacteria. *J Bacteriol* 175:7658–7665.
6. Mavrodi DV, Peever TL, Mavrodi OV, Parejko JA, Raaijmakers JM, Lemanceau P, Mazurier S, Heide L, Blankenfeldt W, Weller DM, Thomashow LS. 2010. Diversity and evolution of the phenazine biosynthesis pathway. *Appl Environ Microbiol* 76:866–879. <http://dx.doi.org/10.1128/AEM.02009-09>.
 7. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <http://dx.doi.org/10.1089/cmb.2012.0021>.
 8. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068455–2069. <http://dx.doi.org/10.1093/bioinformatics/btu153>.