

12-2015

Supporting Data for the Characterization of PNA-DNA Four-Way Junctions

Douglas Van Iverson II
University of Southern Mississippi, douglas.iverson@usm.edu

Crystal C. Serrano
University of Southern Mississippi, crystal.cox@eagles.usm.edu

Ann Marie Brahan
University of Southern Mississippi

Arik Shams
University of Southern Mississippi

Filbert Totsingan
New York University, totsif@rpi.edu

See next page for additional authors

Follow this and additional works at: https://aquila.usm.edu/fac_pubs

 Part of the [Chemistry Commons](#)

Recommended Citation

Iverson, D. V., Serrano, C. C., Brahan, A. M., Shams, A., Totsingan, F., Bell, A. J. (2015). Supporting Data for the Characterization of PNA-DNA Four-Way Junctions. *Data in Brief*, 5, 756-760.
Available at: https://aquila.usm.edu/fac_pubs/15412

This Article is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in Faculty Publications by an authorized administrator of The Aquila Digital Community. For more information, please contact Joshua.Cromwell@usm.edu.

Authors

Douglas Van Iverson II, Crystal C. Serrano, Ann Marie Brahan, Arik Shams, Filbert Totsingan, and Anthony J. Bell Jr.



ELSEVIER

Contents lists available at ScienceDirect

Data in Brief

journal homepage: www.elsevier.com/locate/dib

Data Article

Supporting data for the characterization of PNA–DNA four-way junctions

Douglas Iverson^a, Crystal Serrano^a, Ann Marie Brahan^a,
Arik Shams^a, Filbert Totsingan^b, Anthony J. Bell Jr.^{a,*}^a Department of Chemistry and Biochemistry, The University of Southern Mississippi, Hattiesburg, MS, USA^b Department of Chemistry, New York University, New York, NY, USA

ARTICLE INFO

Article history:

Received 3 September 2015

Received in revised form

12 October 2015

Accepted 13 October 2015

Available online 22 October 2015

ABSTRACT

Holliday or DNA four-way junctions (4WJs) are cruciform/bent structures composed of four DNA duplexes. 4WJs are key intermediates in homologous genetic recombination and double-strand break repair. To investigate 4WJs *in vitro*, junctions are assembled using four asymmetric DNA strands. The presence of four asymmetric strands about the junction branch point eliminates branch migration, and effectively immobilizes the resulting 4WJ. The purpose of these experiments is to show that immobile 4WJs composed of DNA and peptide nucleic acids (PNAs) can be distinguished from contaminating labile nucleic acid structures. These data compare the electrophoretic mobility of hybrid PNA–DNA junctions vs. i) a classic immobile DNA 4WJ, J1 and ii) contaminating nucleic acid structures.

© 2015 Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Specifications Table

Subject area	Biochemistry and biophysics
More specific subject area	Nucleic acid biochemistry
Type of data	Nondenaturing polyacrylamide gel electrophoresis Scanned with Typhoon 9400 Phosphorimager

DOI of original article: <http://dx.doi.org/10.1016/j.dib.2015.08.017>

* Corresponding author.

E-mail address: Anthony.Bell@usm.edu (A.J. Bell Jr.).<http://dx.doi.org/10.1016/j.dib.2015.10.013>2352–3409/© 2015 Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

How data was acquired

Data format

Experimental factors

Experimental features

Data source location

Data accessibility

Scanned gel images

4WJs and DNAs are assembled by mixing, lyophilizing and suspending in annealing buffer [50 mM TrisHCl (pH 7.5), 1 mM MgCl₂].

Immobile 4WJs travel with slower mobility than mobile 4WJs and contaminating nucleic acid structures.

One DNA strand in each sample contains a flourescein label that is quantified via Phoshorimager analysis.

The University of Southern Mississippi, Hattiesburg, MS

The data are supplied with this article.

Value of the data

- The technique provides an approach to confirm the formation of previously uncharacterized 4WJs.
- The technique provides an approach to characterize the global structural features of immobile 4WJs composed of DNA and PNA.
- The technique provides an exhaustive approach to investigate different combinations of potentially contaminating DNA structures vs. immobile hybrid PNA–DNA 4WJs.




Fig. 1. DNA junction, J1, and hybrid PNA–DNA four-way junctions.

1. Data

The nucleic acid sequence of the DNA control, J1 and each hybrid PNA–DNA junction is displayed in Fig. 1 [1]. The data in Figs. 2–4 display the electrophoretic mobility patterns of six hybrid PNA–DNA 4WJs vs. i) J1 and ii) different combinations of potentially contaminating strands of DNA. The contaminating strands are composed of single strands used to form J1. The data are based on the protocol used by Kallenbach and Seaman to characterize, J1 [2,3]. In our previous study, each hybrid PNA–DNA 4WJ is evaluated vs. J1 [4].

2. Experimental design, materials and methods

2.1. Preparation of 4WJs and multi-DNAs

Each 4WJ is formed by lyophilizing a mixture of one fluorescein labeled strand (25 μ M) with 5-fold excess of three unlabeled strands (125 μ M). The multi-DNAs samples are formed by lyophilizing a mixture of a fluorescein labeled strand with 5-fold excess unlabeled stands. The flourescein strands are denoted with an asterisk (below). Each pellet is suspended in 50 mM Tris-HCl (pH 7.5) and 1.0 mM MgCl₂, incubated at 95 °C for 2 min and cooled to room temperature for 12–16 h. To determine the purity of each 4WJ, 2.5 μ M of each sample is loaded onto 15% mini-PROTEAN nondenaturing polyacrylamide gels (BioRad) and run for 1–5 h (4 °C). The gel running buffer is composed of 0.5 \times TBE · MgCl₂ buffer (45 mM Trisma, 45 mM boric acid, 1.0 mM EDTA and 1 mM MgCl₂), pH 7.6. The gels are subsequently scanned with a Typhoon 9400 Phosphorimager. A list of the corresponding oligonucleotides of each multi-DNA sample and 4WJs are shown below.

Fig. 2: Lanes 1–14: 101^{*}, 101^{*}–102, 101^{*}–103, 101^{*}–104, 102–103^{*}, 102–104^{*}, 103–104^{*}, 101^{*}–102–103, 101^{*}–102–104, 101^{*}–103–104, 102–103–104^{*}, J1(101^{*}), 4WJ:PNA₁(103^{*}) and b4WJ:PNA₁(103^{*}).

Fig. 3: Lanes 1–14: lanes 1–14: 101^{*}, 101^{*}–102, 101^{*}–103, 101^{*}–104, 102–103^{*}, 102–104^{*}, 103–104^{*}, 101^{*}–102–103, 101^{*}–102–104, 101^{*}–103–104, 102–103–104^{*}, J1(101^{*}), 4WJ:PNA₃(101^{*}) and b4WJ:PNA₃(101^{*}).

Fig. 4: Lanes 1–15: lanes 1–14: 101^{*}, 101^{*}–102, 101^{*}–103, 101^{*}–104, 102–103^{*}, 102–104^{*}, 103–104^{*}, 101^{*}–102–103, 101^{*}–102–104, 101^{*}–103–104, 102–103–104^{*}, J1 (101^{*}), 4WJ:PNA_{1,3}(104^{*}), b4WJ:PNA_{1,3}(bf104^{*}) and b4WJ:PNA_{1,3}(bf104^{*}).




Fig. 2. Comparison of multi-DNAs vs. J1, 4WJ:PNA₁ and b4WJ:PNA₁.




Fig. 3. Comparison of multi-DNAs vs. J1, 4WJ:PNA₃ and b4WJ:PNA₃.




Fig. 4. Comparison of multi-DNAs vs. J1, 4WJ:PNA_{1,3} and b4WJ:PNA_{1,3}.

3. Data interpretation

The data provide a direct method to compare the mobility of immobile hybrid PNA–DNA junctions vs. a DNA control junction (J1) and potentially contaminating multi-PNAs. In each gel (Figs. 2–4); the single strand control (101) is loaded in lane 1, the contaminating strands are loaded in lanes 2–11, the DNA control, J1 is loaded in lane 12, and the hybrid PNA–DNA junctions are loaded in lanes 13 and 14. In each case, the hybrid 4WJ with a DNA overhang(s) is loaded in lane 13 and the blunt-ended construct is loaded in lane 14.

Acknowledgments

This work was supported by the Mississippi INBRE, funded by an Institutional Development Award (IDeA) from the National Institute of General Medical Sciences of the National Institutes of Health under grant number P20GM103476.

Appendix A. Supplementary material

Supplementary data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.dib.2015.10.013>.

References

- [1] D. Iverson, C. Serrano, A.M. Brahan, A. Shams, F. Totsingan, A.J. Bell Jr., Characterization of the structural and protein recognition properties of hybrid PNA–DNA four-way junctions, *Arch. Biochem. Biophys.* 587 (2015) 1–11.
- [2] N.R. Kallenbach, R.-I. Ma, N.C. Seeman, An immobile nucleic acid junction constructed from oligonucleotides, *Nature* 305 (1983) 829–831.
- [3] N.C. Seeman, N.R. Kallenbach, Design of immobile nucleic acid junctions, *Biophys. J.* 44 (1983) 201–209.
- [4] F. Totsingan, A.J. Bell Jr, Interaction of HMG proteins and H1 with hybrid PNA–DNA junctions, *Protein Sci.* 22 (2013) 1552–1562.