

9-14-2018

## Left Ventricular Mechanical Dyssynchrony for CAD Diagnosis: Does It Have Incremental Clinical Values?

Zhixin Jiang

*The First Affiliated Hospital of Nanjing Medical University*

Weihua Zhou

*University of Southern Mississippi, Weihua.Zhou@usm.edu*

Follow this and additional works at: [https://aquila.usm.edu/fac\\_pubs](https://aquila.usm.edu/fac_pubs)



Part of the [Cardiology Commons](#), and the [Computer Sciences Commons](#)

---

### Recommended Citation

Jiang, Z., Zhou, W. (2018). Left Ventricular Mechanical Dyssynchrony for CAD Diagnosis: Does It Have Incremental Clinical Values?. *Journal of Nuclear Cardiology*.

Available at: [https://aquila.usm.edu/fac\\_pubs/15543](https://aquila.usm.edu/fac_pubs/15543)

This Editorial is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in Faculty Publications by an authorized administrator of The Aquila Digital Community. For more information, please contact [Joshua.Cromwell@usm.edu](mailto:Joshua.Cromwell@usm.edu).

**Left ventricular mechanical dyssynchrony for CAD diagnosis:****Does it have incremental clinical values?**

Zhixin Jiang, MD; Weihua Zhou, PhD

Zhixin Jiang, MD

Department of Cardiology, The First Affiliated Hospital of Nanjing Medical University,  
Nanjing, Jiangsu 210029, China

E-mail: zhixin\_jiang@njmu.edu.cn

or

Weihua Zhou, PhD

School of Computing, University of Southern Mississippi, Long Beach, MS 39560

E-Mail: weihua.zhou@usm.edu

Gated single photon emission computed tomography (SPECT) myocardial perfusion imaging (MPI) has been widely practiced to assess left ventricular (LV) function after stress and at rest, which has in turn improved the diagnosis of coronary artery disease (CAD). The visualization of wall motion and thickening has greatly improved the identification of attenuation artefacts(1, 2) and allows the recognition of stress-induced abnormalities in the case of post-ischemic stunning which helps with multi-vessel CAD detection(3).

Recently, the phase analysis technique has become a major innovation in nuclear cardiology and greatly boosted the clinical values of MPI scans. It measures the LV mechanical dyssynchrony (LVMD) and wall thickening from the regional LV count changes throughout the cardiac cycle (4, 5). Previous studies identified LVMD as an independent predictor for a favorable response to cardiac resynchronization therapy (6) and for all-cause death in patients with dilated cardiomyopathy and narrow QRS (7).

1  
2  
3 LVMD has also demonstrated its clinical values in CAD diagnosis. As reported by  
4 Uebleis et al (8), the occurrence of LVMD is mainly influenced by reduced myocardial  
5 contractility as expressed by the summed thickening score (STS). In a recent study,  
6 STS and the proportion of segments with reduced wall thickening in segments with  
7 normal perfusion were the independent risk factors of LV remodeling in myocardial  
8 infarction patients(9). Thus it is not a surprise to see that LVMD is related with stress-  
9 induced myocardial ischemia (10-13). Chen et al (11) firstly demonstrated different  
10 LVMD changes in ischemic, infarcted and normal myocardium using gated thallium-201  
11 SPECT MPI, which was acquired at 5–10 minutes after stress. Other researchers have  
12 also found the evidence of stress-induced LVMD when acquisitions were performed  
13 earlier than one hour post stress with technetium-99m sestamibi (14-16). Our research  
14 showed that, in patients with moderate to severe myocardial ischemia, LVMD changes  
15 were also related to myocardial stunning even when the acquisition was 60 minutes  
16 after stress(17).  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27

28 Furthermore, the direct comparison with coronary angiography has suggested that  
29 LVMD can help with multi-vessel CAD detection. Hida et al (14) reported that in the  
30 detection of 128 patients with multi-vessel CAD, a summed stress score (SSS)  
31 of  $\geq 9$  showed a sensitivity of 84% and a specificity of 53%, while after exercise an  
32 increase in phase standard deviation (PSD) of  $\geq 4.4^\circ$  and phase bandwidth (PBW) of  
33  $\geq 14^\circ$  had sensitivities of 74% and 68%, and specificities of 84% and 91%, respectively.  
34 Huang et al (18) demonstrated that patients with severe multi-vessel CAD had more  
35 significantly global and territorial LVMD. Gimelli et al (19) reported that in patients with  
36 suspected or known CAD, the presence and severity of LVMD was correlated with the  
37 extent of CAD, and that it was significantly more prevalent in patients with multi-vessel  
38 CAD.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48

49 In the current issue of the journal, Lin et al (20) investigated the diagnosis value of  
50 LVMD in patients with less severe CAD (coronary artery stenosis  $\geq 50\%$ ). A total of 476  
51 patients without known CAD who underwent dipyridamole gated SPECT MPI and  
52 coronary angiography within 6 months were enrolled. Depending on the duration and  
53 severity of ischemia, myocardial stunning lasts from minutes to days. Thus immediate  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 acquisition after stress is better for the detection of this phenomenon (21). In this study,  
4 with thallium-201 and cadmium-zinc-telluride detectors, the acquisition time was  
5 possible during 5 minutes after radiotracer injection. They concluded that perfusion  
6 abnormalities like SSS and summed difference score (SDS) were still the best  
7 diagnostic tools in detecting CAD, although less LVMD was observed post-stress in the  
8 CAD group compared to the non-CAD group. Regarding this study, we have following  
9 comments:

16 Firstly, it is a good attempt to use 50% as a definition for CAD, which tested the  
17 sensitivity of LVMD for CAD diagnosis. In prior studies, coronary artery stenosis of 70%  
18 - 75% was mostly used (14, 18, 19) and myocardial ischemia is usually related with  
19 more severe coronary stenosis ( $\geq 70\%$ ). The current study found that only SDS, age,  
20 hypertension and smoking were related with CAD. Though the results were negative, it  
21 still contributes to the knowledge that LVMD from gated SPECT MPI have limitations for  
22 the diagnosis of mild CAD.

29 Secondly, both global and regional LV function parameters have been used to  
30 detect myocardial stunning, such as reduced LV ejection fraction (21), enlarged end  
31 diastolic volume or end systolic volume (14), and decreased wall motion (22-24) or wall  
32 thickening (24). Whether the addition of LVMD and wall thickening can yield increased  
33 diagnostic value beyond these variables, especially STS, is unknown. Hida et al(14)  
34 found that the combination of post-stress increases in PSD, PBW, transient ischemic  
35 dilation (TID) ratio and SSS best identified multi-vessel CAD (sensitivity 77%, specificity  
36 88%), compared with TID ratio and SSS only (sensitivity 70%, specificity 76%). In  
37 another study(25), patients with systolic dysfunction were evaluated. SDS, SSS and  
38 changes in PSD and PBW with stress were greater in 26 patients with CAD than those  
39 in 15 patients without CAD. In the multivariate analysis a PSD of  $>14^\circ$  and a SSS of  $>17$   
40 best differentiated LV dysfunction of ischemic and non-ischemic etiologies, with a  
41 sensitivity of 89% and a specificity of 87%, compared with SDS only (sensitivity, 46%;  
42 specificity, 87%).

1  
2  
3 Last, it is noteworthy that a total of 476 patients and over 27 clinical variables  
4 were included in the patient baseline, univariate and multivariate, and subgroup analysis.  
5 Variable interaction may be an issue in the statistical analysis, for example, it has been  
6 demonstrated by Ludwig et al that PSD may be spuriously increased by scar(26). These  
7 may challenge the traditional linear analysis. Algorithms for the non-linear model with a  
8 big number of inter-correlated variables may further improve the accuracy of the  
9 prediction. Machine learning may be a good choice. In a recent multi-center study with  
10 10,030 patients with suspected CAD, Motwani et al used machine learning to predict 5-  
11 year all-cause mortality and achieved an AUC of 0.79(27). In their method, the  
12 information gain algorithm was employed to select important variables from 25 clinical  
13 and 44 coronary CTA parameters and the LogitBoost algorithm was employed to build  
14 the prediction model.  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26

## 27 References

- 28 (1) Chua T, Kiat H, Germano G, Maurer G, van Train K, Friedman J et al. Gated technetium-99m  
29 sestamibi for simultaneous assessment of stress myocardial perfusion, postexercise regional  
30 ventricular function and myocardial viability. Correlation with echocardiography and rest  
31 thallium-201 scintigraphy. *J Am Coll Cardiol* 1994;23:1107-14.
- 32 (2) Nichols K, DePuey EG, Krasnow N, Lefkowitz D, Rozanski A. Reliability of enhanced gated SPECT  
33 in assessing wall motion of severely hypoperfused myocardium: echocardiographic validation. *J*  
34 *Nucl Cardiol* 1998;5:387-94.
- 35 (3) Sciagra R. The expanding role of left ventricular functional assessment using gated myocardial  
36 perfusion SPECT: the supporting actor is stealing the scene. *Eur J Nucl Med Mol Imaging*  
37 2007;34:1107-22.
- 38 (4) Chen J, Garcia EV, Folks RD, Cooke CD, Faber TL, Tauxe EL et al. Onset of left ventricular  
39 mechanical contraction as determined by phase analysis of ECG-gated myocardial perfusion  
40 SPECT imaging: development of a diagnostic tool for assessment of cardiac mechanical  
41 dyssynchrony. *J Nucl Cardiol* 2005;12:687-95.
- 42 (5) Zhou W, Garcia EV. Nuclear Image-Guided Approaches for Cardiac Resynchronization Therapy  
43 (CRT). *Curr Cardiol Rep* 2016;18:7.
- 44 (6) Henneman MM, Chen J, Dibbets-Schneider P, Stokkel MP, Bleeker GB, Ypenburg C et al. Can LV  
45 dyssynchrony as assessed with phase analysis on gated myocardial perfusion SPECT predict  
46 response to CRT? *J Nucl Med* 2007;48:1104-11.
- 47 (7) Kano N, Okumura T, Isobe S, Sawamura A, Watanabe N, Fukaya K et al. Left ventricular phase  
48 entropy: Novel prognostic predictor in patients with dilated cardiomyopathy and narrow QRS. *J*  
49 *Nucl Cardiol* 2017.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1
- 2
- 3 (8) Uebleis C, Hoyer X, Van Krieking SD, Schuessler F, Laubender RP, Becker A et al. Association
- 4 between left ventricular mechanical dyssynchrony with myocardial perfusion and functional
- 5 parameters in patients with left bundle branch block. *J Nucl Cardiol* 2013;20:253-61.
- 6 (9) Yang W, Zhang F, Tang H, Shao X, Wang J, Wang X et al. Summed thickening score by myocardial
- 7 perfusion imaging: A risk factor of left ventricular remodeling in patients with myocardial
- 8 infarction. *J Nucl Cardiol* 2018;25:742-53.
- 9
- 10 (10) Aljaroudi W, Koneru J, Heo J, Iskandrian AE. Impact of ischemia on left ventricular dyssynchrony
- 11 by phase analysis of gated single photon emission computed tomography myocardial perfusion
- 12 imaging. *J Nucl Cardiol* 2011;18:36-42.
- 13 (11) Chen CC, Shen TY, Chang MC, Hung GU, Chen WC, Kao CH et al. Stress-induced myocardial
- 14 ischemia is associated with early post-stress left ventricular mechanical dyssynchrony as
- 15 assessed by phase analysis of 201TI gated SPECT myocardial perfusion imaging. *Eur J Nucl Med*
- 16 *Mol Imaging* 2012;39:1904-9.
- 17 (12) Singh H, Patel CD, Sharma P, Naik N, Singh S, Narang R. Does perfusion pattern influence stress-
- 18 induced changes in left ventricular mechanical dyssynchrony on thallium-201-gated SPECT
- 19 myocardial perfusion imaging? *J Nucl Cardiol* 2015;22:36-43.
- 20 (13) Zhou Y, Li D, Feng J, Yuan D, Patel Z, Cao K et al. Left Ventricular Dyssynchrony Parameters
- 21 Measured by Phase Analysis of Post-stress and Resting Gated SPECT Myocardial Perfusion
- 22 Imaging. *World journal of nuclear medicine* 2013;12:3-7.
- 23 (14) Hida S, Chikamori T, Tanaka H, Igarashi Y, Shiba C, Usui Y et al. Diagnostic value of left ventricular
- 24 dyssynchrony after exercise and at rest in the detection of multivessel coronary artery disease
- 25 on single-photon emission computed tomography. *Circ J* 2012;76:1942-52.
- 26 (15) Karacalioglu AO, Balta S, Emer O, Demirkol S, Celik T, Ozguven M. Phase analysis in patients with
- 27 reversible perfusion defects and normal coronary arteries at angiography. *Ann Nucl Med*
- 28 *2013;27:416-22.*
- 29 (16) Peix A, Cabrera LO, Padron K, Rodriguez L, Fernandez J, Lopez G et al. Association between non-
- 30 perfusion parameters and presence of ischemia in gated-SPECT myocardial perfusion imaging
- 31 studies. *Journal of nuclear cardiology : official publication of the American Society of Nuclear*
- 32 *Cardiology* 2016.
- 33 (17) Jiang Z, Tang H, Shi J, Zhou Y, Wang C, Li D et al. Myocardial stunning-induced left ventricular
- 34 dyssynchrony on gated single-photon emission computed tomography myocardial perfusion
- 35 imaging. *Nucl Med Commun* 2018;39:725-31.
- 36 (18) Huang WS, Huang CH, Lee CL, Chen CP, Hung GU, Chen J. Relation of early post-stress left
- 37 ventricular dyssynchrony and the extent of angiographic coronary artery disease. *J Nucl Cardiol*
- 38 *2014;21:1048-56.*
- 39 (19) Gimelli A, Liga R, Giorgetti A, Favilli B, Pasanisi EM, Marzullo P. Determinants of left ventricular
- 40 mechanical dyssynchrony in patients submitted to myocardial perfusion imaging: A cardiac CZT
- 41 study. *J Nucl Cardiol* 2016;23:728-36.
- 42 (20) Lin WL, Wang SY, Shiao YC, Wu YW. The Clinical Usefulness of Phase Analysis in Detecting
- 43 Coronary Artery Disease Using Dipyridamole Thallium-201-Gated Myocardial Perfusion Imaging
- 44 with aCadmium-Zinc-Telluride Camera. *J Nucl Cardiol* 2018. (accepted)
- 45 (21) Mut F, Giubbini R, Vitola J, Lusa L, Sobic-Saranovic D, Peix A et al. Detection of post-exercise
- 46 stunning by early gated SPECT myocardial perfusion imaging: results from the IAEA multi-center
- 47 study. *J Nucl Cardiol* 2014;21:1168-76.
- 48 (22) Ward RP, Gundeck EL, Lang RM, Spencer KT, Williams KA. Overestimation of postischemic
- 49 myocardial stunning on gated SPECT imaging: correlation with echocardiography. *Journal of*
- 50 *nuclear cardiology : official publication of the American Society of Nuclear Cardiology*
- 51 *2006;13:514-20.*
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

- 1  
2  
3 (23) Hacker M, Rieber J, Schmid R, Lafougere C, Tausig A, Theisen K et al. Comparison of Tc-99m  
4 sestamibi SPECT with fractional flow reserve in patients with intermediate coronary artery  
5 stenoses. *Journal of nuclear cardiology : official publication of the American Society of Nuclear*  
6 *Cardiology* 2005;12:645-54.  
7  
8 (24) Santiago JF, Heiba SI, Jana S, Mirzaitehrane M, Dede F, Abdel-Dayem HM. Transient ischemic  
9 stunning of the myocardium in stress thallium-201 gated SPET myocardial perfusion imaging:  
10 segmental analysis of myocardial perfusion, wall motion and wall thickening changes. *European*  
11 *journal of nuclear medicine and molecular imaging* 2002;29:979-83.  
12 (25) Igarashi Y, Chikamori T, Hida S, Tanaka H, Shiba C, Usui Y et al. Usefulness of phase analysis to  
13 differentiate ischemic and non-ischemic etiologies of left ventricular systolic dysfunction in  
14 patients with heart failure. *Circ J* 2014;78:141-50.  
15 (26) Ludwig DR, Friehling M, Schelbert EB, Schwartzman D. Impact of scar on SPECT assay of left  
16 ventricular contraction dyssynchrony. *Eur J Nucl Med Mol Imaging* 2014;41:529-35.  
17 (27) Motwani M, Dey D, Berman DS, Germano G, Achenbach S, Al-Mallah MH et al. Machine learning  
18 for prediction of all-cause mortality in patients with suspected coronary artery disease: a 5-year  
19 multicentre prospective registry analysis. *Eur Heart J* 2017;38:500-7.  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

For Peer Review



# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

For Peer Review