

2-1-2005

A Regional Food-Frequency Questionnaire for the US Mississippi Delta

Katherine L. Tucker
U.S. Dept. of Agriculture

Janice Maras
U.S. Dept. of Agriculture

Catherine Champagne
Pennington Biomedical Research Center

Carol L. Connell
University of Southern Mississippi, carol.connell@usm.edu

Susan Goolsby
Arkansas Children's Hospital Research Institute

See next page for additional authors

Follow this and additional works at: https://aquila.usm.edu/fac_pubs



Part of the [Nutrition Commons](#)

Recommended Citation

Tucker, K. L., Maras, J., Champagne, C., Connell, C. L., Goolsby, S., Weber, J., Zaghoul, S., Carithers, T., Bogle, M. L. (2005). A Regional Food-Frequency Questionnaire for the US Mississippi Delta. *Public Health Nutrition*, 8(1), 87-96.
Available at: https://aquila.usm.edu/fac_pubs/2881

Authors

Katherine L. Tucker, Janice Maras, Catherine Champagne, Carol L. Connell, Susan Goolsby, Judith Weber, Sahar Zaghoul, Teresa Carithers, and Margaret L. Bogle

A regional food-frequency questionnaire for the US Mississippi Delta

Katherine L Tucker^{1,*}, Janice Maras¹, Catherine Champagne², Carol Connell³, Susan Goolsby⁴, Judith Weber⁴, Sahar Zaghoul^{5,†}, Teresa Carithers⁶ and Margaret L Bogle⁷

¹Jean Mayer US Department of Agriculture (USDA) Human Nutrition Research Center on Aging at Tufts University, 711 Washington Street, Boston, MA 02111, USA: ²Pennington Biomedical Research Center, Baton Rouge, LA, USA: ³University of Southern Mississippi, Hattiesburg, MS, USA: ⁴Arkansas Children's Hospital Research Institute, Little Rock, AR, USA: ⁵University of Arkansas at Pine Bluff, Pine Bluff, AR, USA: ⁶The Jackson Heart Study, Jackson, MS, USA: ⁷USDA Agricultural Research Service, Little Rock, AR, USA

Submitted 16 April 2004: Accepted 13 July 2004

Abstract

Objective: To describe food sources of nutrient intake for white and African American adults in the Lower Mississippi Delta (LMD), and their use in the development of a regional food-frequency questionnaire (FFQ) based on an earlier version of the National Cancer Institute's Health Habits and History Questionnaire.

Design: We ranked food sources of energy, macronutrients, vitamins and minerals, and examined portion size distributions for 842 white and 857 African American residents aged 19 years and older, using 24-hour dietary intake recall data from a telephone survey of 36 LMD counties. These values were used to develop a regional FFQ, which was then field-tested with 100 subjects and revised to improve interpretability.

Setting: The LMD region of the USA.

Subjects: White and African American adult residents of the LMD.

Results: LMD African Americans obtained more of their energy and nutrient intakes from poultry, processed meat, salty snacks, fruit drinks, pork and cornbread; and less from milk, alcohol, legumes, salad dressing, butter/margarine and sweetened tea than did white residents. Regional foods not on nationally used FFQs included grits, turnip greens, okra, ham hocks, chitterlings, crawfish, catfish, cracklings, jambalaya, potato logs, chicken and dumplings, and sweet potato pie. Based on responses during field-testing, the questionnaire was also designed to add four portion sizes for each food item, presented as questions, rather than in grid format.

Conclusions: Regional food use patterns differ from national patterns and furthermore differ between African American and white adults in the LMD. The resulting Delta NIRI FFQ for Adults should contribute to improved assessment of usual intake for use in studies of diet and health in this region.

Keywords

Food-frequency questionnaire
Lower Mississippi Delta
Dietary methodology
Dietary intake
African Americans

Despite limitations in the quantitative assessment of macronutrients¹, the food-frequency questionnaire (FFQ) has proved to be a useful tool for ranking individuals' usual nutrient intakes for assessing associations between dietary components and health outcomes². The most widely used FFQs have been designed to capture the foods most commonly consumed in the USA^{3,4}. However, the intakes of individuals who follow very different dietary patterns from the national norm may be underestimated due to the absence of specific foods used from the food list, or may be incorrectly estimated due to differing

recipes for commonly used foods. When a different dietary pattern is prevalent in the population under study, the error may lead to extensive misclassification. In addition to attenuating the ability to relate dietary components to specific outcome measures, this may result in biased assessments of a group or subgroup. It is therefore important to test questionnaires and, if necessary, adapt them to each population under study.

The Lower Mississippi Delta (LMD) region of the USA is an important area in which to conduct nutrition research. One of the most impoverished rural regions in the country, it has higher than national average mortality rates from nutrition-related chronic diseases including heart disease and cancer⁵. Despite the key role that diet is known to play

†Present address: University of Hawaii at Manoa, Honolulu, HI, USA.

*Corresponding author: Email katherine.tucker@tufts.edu

in the aetiology of these diseases, few studies have examined the potential role of diet in these excessive health risks. None of the major national nutrition surveys has collected dietary data in this region. One goal of the Lower Mississippi Delta Nutrition Intervention Research Initiative (Delta NIRI), funded by the US Department of Agriculture's (USDA) Agricultural Research Service, is to obtain data that will help clarify these associations and to identify effective nutrition interventions that may assist the population in reducing these risks. A telephone survey in the Delta region collected representative dietary data using 24-hour dietary recalls. A previous report of these data showed that intakes of most micronutrients were significantly lower in the LMD, relative to national data, and that this was particularly true among the African American subset⁶.

We used these data to develop a new FFQ (the Delta NIRI Food Frequency Questionnaire for Adults) designed for use in the LMD region. In this report, we present the methodology used for this development and highlight the uniqueness of foods, preparations and portion sizes used by African American and white adults in this region.

Materials and methods

Data on dietary intakes in this region were available from 24-hour dietary recalls collected in the Delta NIRI Foods of our Delta Survey 2000 (FOODS 2000), between January and June 2000. FOODS 2000 was a cross-sectional telephone survey of a representative sample of the population 3 years of age and older in 36 counties and parishes in the LMD of Arkansas, Louisiana and Mississippi. Each sample person provided a single 24-hour recall. A previous validation study conducted by Delta NIRI had confirmed the feasibility of using a telephone survey to collect dietary data within this region⁷. For FFQ development, we selected data from adults aged 19 years and older ($n = 842$ white and 857 African American) to describe the major food contributors to energy and nutrient intake, to identify commonly used portion sizes and to obtain recipes for unique food items. This information was then used to develop a regionally appropriate food list and portion size options for the FFQ; and to provide weighting and recipe information for the nutrient database used to analyse questionnaire responses.

FOODS 2000 methodology

Westat (Rockville, MD, USA) conducted participant sampling, interviewer training and telephone interviews for FOODS 2000. Sampling was done using a two-stage stratified cluster design, as previously described⁶. Briefly, 36 Delta counties were assigned to nine strata according to population size, racial composition and income; and two sampling units (based on census definition) per

strata were selected randomly. List-assisted methodology was used to select a sample of 4377 working residential telephone numbers in the 18 sample units. Response to the screener interview was 58.6%, and a total of 2162 households agreed to participate. One adult per household was selected randomly to complete the 24-hour recall.

Thirty-three telephone interviewers completed a 4-day training session on administration of the dietary questionnaires, using food-measuring guides. The 24-hour dietary recall was conducted using the USDA multiple-pass procedure⁸. Prior to completing the recall, the food measurement guide was mailed to participants for use during the interview. The response rate was 80.5%. Data were coded, entered and analysed for nutrient content at the Pennington Biomedical Research Center using the USDA SurveyNet nutrient composition tables⁹.

Ranking of food contributions to nutrient intake

The resulting food and nutrient variables from the survey were transferred to the Dietary Assessment and Epidemiology Research Program at the USDA Jean Mayer Human Nutrition Research Center on Aging at Tufts University. These were analysed for food contributions to energy and nutrient intakes, portion size distributions and recipe detail. Foods reported on the 24-hour dietary recalls were grouped into 140 food groups, based on similarity of nutrient composition and common usage. These food groupings are similar, though not identical, to the 112 groups used by Subar *et al.* in their description of the dietary sources of nutrients in the diets of US adults¹⁰. These food groups were then ranked by their percentage contribution to individual nutrients, using the RANK procedure in SAS¹¹. Foods that contributed 0.5% or more to the intake of energy or any single nutrient were included in the list of foods on the questionnaire. Local names of foods were used, as appropriate, based on pre-testing with adult groups in the LMD. Foods that did not appear on existing commonly used FFQs, but were relevant contributors to intake, were identified from these lists for inclusion on the new FFQ.

FFQ form development, field trial and revision

Beginning with an adapted version of the 1989 National Cancer Institute (NCI)–Block FFQ¹², and following the methodology used in the development of that questionnaire, we revised the food list to include all foods that contributed at least 0.5% to any of the nutrients examined. Because of the extended distribution of portion sizes reported in the FOODS 2000 data, we replaced the small, medium and large designations with four target sizes, based on the distribution of portion sizes observed for each food.

Questions on the use of common vitamin and mineral supplement formulations, as well as separate use of 12 individual vitamin and mineral supplements, were also

included so that total nutrient intakes could be calculated. Another modification, borrowed from the Harvard FFQ⁴, was the inclusion of open-ended responses for types of cereal most frequently consumed and for additional foods consumed at least once per week. These were coded directly onto the form prior to scanning, based on a code book of reported foods that can be adapted to include new responses linked to parallel additions to the database.

The draft form was field-tested at several community sites in the LMD and in Jackson, Mississippi. Using a variant of cognitive interviewing methodology, more than 100 questionnaires were interviewer-administered to individuals in the region. Reactions, difficulties or questions by respondents were noted systematically and discussed in depth at investigator meetings.

Database development

Nutrient data from the USDA SurveyNet database⁹ were used for the development of a nutrient database for the FFQ. Each food line represented several specific food items. Within each line item, the frequency of foods reported from the 24-hour dietary recall was ranked. For example, in the line 'Strawberries, other berries', all reported berries were ranked according to frequency of appearance in the dataset. Those contributing at least 5% of reported usage were assigned a proportional share so that the sum of those included totalled 100%. The nutrient value for each food was then weighted by this proportion to represent a composite food for the line.

In addition to weighting of specific foods, recipes for combination food items were added. For example, gumbo and jambalaya are local preparations that may vary from household to household. Specific recipe information collected during the FOODS 2000 24-hour dietary intake recall was examined to determine whether a standard recipe could be used. When multiple recipes were identified, they were ranked, as described above for foods, and the nutrient values of each were included in relative proportion to their reported frequency.

Results

Food contributors to energy intake

For this sample of both African American ($n = 857$) and white ($n = 842$) LMD adults, approximately 7% of energy came from soft drinks, followed by 6% from white bread, 4–5% from burgers or meatloaf and 3–4% from salty snacks. After that, however, food contributions differed considerably between the two racial groups. African Americans reported greater relative consumption than white LMD adults of fried chicken, sausage and fried fish; and lower relative consumption of salad dressing. Other top energy contributors for African Americans included cornbread and luncheon meat; while for white adults, these included beefsteaks and roasts, sweetened tea and milk (Table 1).

Food contributors to macronutrient intake

The top five food contributors of carbohydrate, protein, fat and dietary fibre are presented in Table 2 for LMD residents, by race. For both groups, the top sources of carbohydrate were soft drinks (15%) and bread (about 9%). Salty snacks were also important contributors to both groups. Cakes, muffins and sweetened tea contributed more carbohydrate intake for whites, and fruit drinks and candy for African Americans. The top protein source for African American adults was fried chicken, followed by burgers and meatloaf, poultry (not fried) and fried fish; while for white adults they included luncheon meats, and beefsteaks and roasts, but not fish. Fat sources also differed by race; white adults consumed more fat from salad dressings and cheese, and African Americans consumed more from fried chicken, sausage, salty snacks and fried fish. Both groups consumed the greatest proportion of fibre from white bread, followed by beans.

Food contributors to vitamin intake

Fortified white bread was the major source of folate for both groups (Table 3), followed by breakfast cereal, tea and orange juice. Breakfast cereals were also important contributors of vitamin B₆ in this population along with bananas. African Americans consumed most of their vitamin B₆ from fried chicken, while white adults consumed more vitamin B₆ from beefsteaks and roasts. Although both groups consumed most of their vitamin B₁₂ from burgers and meatloaf, white adults consumed relatively more from beefsteaks and roasts and milk, and less from fried fish, liver and sausage, than the African American adults.

While orange juice was the major source of vitamin C for both groups, African Americans consumed relatively more vitamin C from fruit drinks, while white adults consumed more from potatoes (Table 4). Vitamin A sources differed considerably by race. White adults consumed 10% of their vitamin A from carrots, but carrots were not within the top five sources for African Americans. Rather, the top source of vitamin A for African Americans was sweet potatoes, followed by chicken liver, greens and fried beef liver. The major source of vitamin E for African Americans was salty snacks (12%), followed by fried fish, while for white adults the major source was mayonnaise and salad dressing.

Food contributors to mineral intake

White adults obtained 26% of their calcium from milk, relative to less than 15% for African Americans (Table 5). African Americans consumed almost 5% of their calcium from cornbread. The largest source of iron for both racial groups was white bread, followed by breakfast cereal and beefsteaks and roasts for white adults; and ground beef, fried chicken and grits for African American adults. Beef was the major source of zinc for both groups. Other top sources included gumbo and luncheon meat for white adults, and poultry and sausage for African American

Table 1 Top 20 contributors to energy intake in adults of the Lower Mississippi Delta, by race

Food	White (n = 842)		African American (n = 857)	
	%	Rank	%	Rank
Soft drinks	7.0	1	7.1	1
White bread, rolls	5.8	2	6.0	2
Burgers, meatloaf	3.8	3	4.6	4
Salty snacks	2.8	4	4.2	5
Cakes, muffins	2.6	5	2.0	12
Mayonnaise, salad dressing	2.4	6	1.4	20
French fries	2.3	7	2.1	11
Fried chicken	2.3	8	4.6	3
Beefsteaks, roasts	2.2	9		
Fried fish	2.1	10	3.0	7
Cookies	1.9	11	2.3	8
Tea, sweetened	1.9	12		
2% Milk	1.8	13		
Cheese	1.7	14	1.5	19
Whole milk	1.6	15		
Other potatoes	1.6	16		
Biscuits	1.6	17		
Candy (not chocolate)	1.5	18	1.9	13
Sausage	1.4	19	3.1	6
Ice cream	1.4	20	1.6	15
Cornbread			2.1	10
Luncheon meat			1.6	18
Spaghetti and sauce, lasagne			1.6	16
Orange juice			1.6	14
Fruit drinks			2.2	9
Poultry, broiled			1.6	17
Intake, mean \pm standard deviation	Men	Women	Men	Women
Energy (kcal)	2450 \pm 1107	1791 \pm 828	2327 \pm 1215	1672 \pm 925

adults. Among white adults, the major magnesium sources were coffee and milk. In contrast, African Americans consumed the largest proportion (5%) of their magnesium from salty snacks.

While these results describe differences in sources of these nutrients, and the comparison of nutrient intakes is the subject of another publication⁶, it is important to note that the African American men and women in this sample reported, on average, lower intakes of each of these minerals than did the white men and women. This was reflected in apparently unusual sources for some nutrients. For example, the average magnesium intakes were 248 mg for African American men and 182 mg for women (relative to current Recommended Dietary Allowances¹³ of 420 and 320 mg, respectively), and this is reflected in the fact that foods commonly consumed, like salty snacks, make larger contributions to intake than foods that are considerably better sources of the nutrient per serving.

Foods added to the questionnaire

Although only the top five food contributors of each nutrient are presented in the tables here for space considerations, regional foods identified from the 24-hour recalls that were seen to contribute more than 0.5% of any of the nutrients examined were included on the questionnaire (Table 6). These include grits, turnip greens, field peas, okra, ham hocks, chitterlings, squirrel, crawfish, catfish and cracklings. Regionally common preparations and recipes added to the database included

broccoli and rice casserole, jambalaya, dirty rice, potato logs, chicken and dumplings, and sweet potato pie. In addition, local names for foods were added, such as 'chicken fried steak', a commonly used beef preparation. Because of the large contribution of drinks such as lemonade and powdered drink mixes for the African American group and sweetened iced tea for the white adults to the energy, carbohydrate and/or vitamin C intakes of this population, more specification of these categories was added to the FFQ. Incorporation of all foods reported on the 24-hour recalls that contributed at least 0.5% to the intake of energy or tested nutrients, separately for white and African American adults, captured more than 98% of reported energy and of each of the examined macronutrient, vitamin and mineral intakes for both race groups.

Portion size

The distributions of portion sizes of common foods on the FFQ were examined in the FOODS 2000 data. We present the 15th, 50th and 85th percentiles in Table 7. Examination of these distributions suggested that the inclusion of four portion size options on the questionnaire was necessary to capture the variation in reported portion sizes. For many foods, the 85th percentile exceeded two times the median. On most existing FFQs, a small portion is considered half of a medium portion, and a large portion is 1.5 times a medium portion. Using these designations would clearly underestimate some of the large portion sizes observed

Table 2 Top food contributors to macronutrient intakes in adults of the Lower Mississippi Delta, by race

Food	White (n = 842)		African American (n = 857)	
	%	Rank	%	Rank
Carbohydrate				
Soft drinks	14.9	1	15.0	1
White bread, rolls	8.5	2	8.7	2
Tea, sweetened	4.1	3		
Cakes, muffins	3.5	4		
Salty snacks (chips)	2.7	5	3.9	4
Candy (not chocolate)			3.3	5
Fruit drinks			6.2	3
Protein				
Burgers, meatloaf	7.1	1	8.4	2
Beefsteaks, roast	7.1	2		
Fried chicken	5.4	3	11.8	1
White bread, rolls	4.5	4	4.7	5
Luncheon meat	3.9	5		
Fried fish			5.0	4
Poultry, not fried			5.6	3
Fat				
Mayonnaise, salad dressing	6.2	1		
Burgers, meatloaf	5.9	2	6.9	1
Salty snacks (chips)	3.9	3	5.9	4
Fried fish	3.3	4	4.8	5
Cheese	3.3	5		
Fried chicken			6.4	3
Sausage			6.4	2
Fibre				
White bread, rolls	7.9	1	9.1	1
Beans	6.6	2	7.7	2
High-fibre cereal	5.1	3		
Salty snacks (chips)	4.8	4	7.3	3
French fries	3.8	5	3.6	4
Cornbread			3.2	5
Intake, mean \pm standard deviation				
	Men	Women	Men	Women
Carbohydrate (g)	288 \pm 143	219 \pm 106	273 \pm 151	205 \pm 116
Protein (g)	94 \pm 50	66 \pm 33	89 \pm 50	63 \pm 38
Fat (g)	96 \pm 54	71 \pm 44	94 \pm 61	67 \pm 47
Fibre (g)	15.3 \pm 10.5	12.7 \pm 8.2	13.5 \pm 9.8	10.7 \pm 8.5

here. Based on the pattern of these distributions for most foods, and for ease of later computation, portion sizes were designed with the second portion choices representing a standard reference amount (usually the median and most common size). The first choices were half of the reference portion, the third 1.5 times the reference portion, and the fourth twice the reference portion in order to capture the extra large servings that subsets of the population reported consuming.

FFQ format

Results of the field trials revealed that the grid format for combined frequency and portion sizes was confusing for the respondents. Participants responded favourably to the grid format for frequencies, but preferred portion sizes to be asked as separate questions following each frequency grid. For example, portion size choices were presented for cooked vegetables, rather than repeatedly for broccoli, carrots, corn and so on. In many cases, this also allowed for the collapsing of portion sizes of similar foods, previously found to be repetitive. Despite evidence of very large portions in the recall data, participants seemed reluctant to select the extra large category, suggesting that

this label may hold negative connotations and could lead to underreporting if presented in this way. We therefore changed the portion sizes to standard reference amounts, rather than labelling them as small, medium, large and extra large. Additionally, we found that participants responded more rapidly when identifying the target sizes their portion was closest to, rather than ranges.

This new format (Appendix) also allowed for the inclusion of adjustment questions within each food group section throughout the questionnaire. Rather than including items such as milk in coffee on another line on the grid, or having a separate section for fat adjustment questions like removing skin on chicken, it was preferable to respondents to answer these questions at the end of each food group section. This, along with the inclusion of open-ended questions on specific cereal brands and additional foods not on the list, allowed for some simplification of foods listed in the grid.

The average time of completion for the FFQ was 45 min before modification. Modification did not lead to major changes in the list of food line items, with 142 in the final version. Rather, it led to changes in the ordering of items, changes in the names of some items and in the addition of

Table 3 Top food contributors to B vitamin intakes in adults of the Lower Mississippi Delta, by race

Food	White (<i>n</i> = 842)		African American (<i>n</i> = 857)	
	%	Rank	%	Rank
Folate				
White bread, rolls	11.9	1	13.2	1
Regular breakfast cereals	5.4	2	5.7	2
Tea	4.9	3	4.6	4
Orange juice	3.9	4	5.6	3
High-fibre cereals	3.7	5		
Rice			3.5	5
Vitamin B₆				
Regular breakfast cereals	5.5	1	5.7	2
Bananas	5.2	2	5.4	3
Beefsteaks, roast, ribs	4.6	3		
Other potatoes	4.4	4		
Fried chicken	4.2	5	9.1	1
Burgers, meatloaf			4.6	4
Salty snacks (chips)			4.5	5
Vitamin B₁₂				
Burgers, meatloaf	11.5	1	14.3	1
Beefsteaks, roast, ribs	10.3	2		
Fried fish	5.8	3	8.9	2
2% Milk	5.7	4		
Whole milk	4.0	5		
Fried liver			7.2	3
Chicken liver			6.9	4
Sausage			5.7	5
Intake, mean ± standard deviation				
	Men	Women	Men	Women
Folate (µg)	398 ± 236	297 ± 177	338 ± 209	262 ± 177
Vitamin B ₆ (mg)	2.0 ± 1.3	1.4 ± 0.9	1.8 ± 1.3	1.3 ± 0.8
Vitamin B ₁₂ (µg)	5.8 ± 5.3	3.9 ± 4.1	4.5 ± 5.1	3.8 ± 7.7

Table 4 Top food contributors to intakes of vitamins C, A and E from the Lower Mississippi Delta Nutrition Intervention Research Initiative Foods of our Delta Survey 2000

Food	White (<i>n</i> = 842)		African American (<i>n</i> = 857)	
	%	Rank	%	Rank
Vitamin C				
Orange juice	23.8	1	27.7	1
Fruit drinks	6.6	2	21.2	2
Oranges	6.0	3	5.5	3
Tomatoes	4.6	4	3.2	4
Other potatoes	3.7	5		
Grapefruit juice			3.2	5
Vitamin A				
Carrots	10.2	1		
Sweet potatoes	5.4	2	9.8	1
2% Milk	5.3	3		
Regular breakfast cereals	4.1	4	5.0	5
High-fibre cereals	3.6	5		
Chicken liver			6.4	2
Greens			6.3	3
Fried liver			5.5	4
Vitamin E				
Mayonnaise, salad dressing	9.1	1	5.8	3
Salty snacks (chips)	4.5	2	12.2	1
White bread, rolls	4.3	3	4.5	4
Fried fish	4.3	4	6.7	2
Highly fortified cereals	4.3	5		
Fried chicken			3.9	5
Intake, mean ± standard deviation				
	Men	Women	Men	Women
Vitamin C (mg)	80 ± 91	73 ± 85	102 ± 112	86 ± 90
Vitamin A (RE)	873 ± 999	760 ± 833	756 ± 1250	692 ± 1247
Vitamin E (TE)	10.3 ± 11.6	8.2 ± 7.0	10.1 ± 25.5	6.6 ± 5.3

RE – retinol equivalents; TE – tocopherol equivalents.

Table 5 Top food contributors to mineral intakes from the Lower Mississippi Delta Nutrition Intervention Research Initiative Foods of our Delta Survey 2000

Food	White (<i>n</i> = 842)		African American (<i>n</i> = 857)	
	%	Rank	%	Rank
Calcium				
2% Milk	12.3	1	6.5	4
Cheese	9.1	2	10.2	1
Whole milk	8.7	3	7.9	2
White bread, rolls	6.0	4	7.8	3
Skimmed or 1% milk	5.2	5		
Cornbread			4.6	5
Iron				
White bread, rolls	8.5	1	9.9	1
Regular breakfast cereals	7.5	2	5.9	3
High-fibre cereals	5.5	3		
Burgers, meatloaf	4.6	4	6.1	2
Beefsteaks, roast	3.6	5		
Fried chicken			3.7	4
Grits			3.0	5
Zinc				
Beefsteaks, roast	9.5	1	4.6	3
Burgers, meatloaf	9.4	2	12.5	1
Gumbo	4.0	3		
High-fibre cereals	3.2	4		
Luncheon meats	2.7	5		
Fried chicken			6.4	2
Poultry, not fried			4.4	4
Sausage			3.9	5
Magnesium				
Coffee	6.7	1		
2% Milk	3.8	2		
White bread, rolls	3.7	3	5.0	2
Salty snacks (chips)	3.2	4	5.4	1
High-fibre cereals	2.9	5		
Orange juice			3.9	3
Fried chicken			3.8	4
Burgers, meatloaf			3.4	5
Intake, mean ± standard deviation				
	Men	Women	Men	Women
Calcium (mg)	841 ± 633	662 ± 420	643 ± 443	506 ± 407
Iron (mg)	16.9 ± 11.9	12.8 ± 7.9	14.7 ± 8.5	10.9 ± 6.9
Zinc (mg)	14.2 ± 12.0	10.0 ± 7.9	11.7 ± 9.2	8.3 ± 6.1
Magnesium (mg)	319 ± 192	234 ± 115	248 ± 147	182 ± 112

recipe questions to clarify the preparation methods. Although separation of the portion sizes from the grid format and the addition of preparation questions led to increased physical length, it did not increase the time of completion, which remained at approximately 45 min with no apparent differences in time of completion by race or sex.

Discussion

The accurate assessment of dietary intake is critical to the success of population-based studies that link dietary behaviour with health outcomes. Although a single 24-hour dietary recall per person has been shown to represent mean intake for a large group, it is well known that this method is less successful in ranking individual food and nutrient intakes¹⁴. Use of a single 24-hour dietary intake recall results in misclassification of individual usual intake due to normal day-to-day variation in intake, and makes it more difficult to see existing associations

between food or nutrient intake and disease outcomes for individuals². For some nutrients, many days of dietary intake are needed to average this variation to be representative of individual nutrient intake¹⁵. Both the collection and processing of dietary recall or food intake record data are also demanding and expensive. For these reasons, the FFQ is used extensively in population studies².

The most commonly used FFQs were designed for use in the general US population. The foods listed capture either the greatest proportion of total nutrient intake in the population³ or the greatest variance in intake⁴ and contain assumptions about average portion size. In either case, subgroups of the population with diets that vary from the norm are unlikely to be represented in these decisions. The importance of a culturally and regionally specific questionnaire was demonstrated previously among the Puerto Rican population in the north-eastern USA, where it was shown that use of a nationally used questionnaire would not only underestimate intakes of energy and

Table 6 Regional foods added to the Lower Mississippi Delta Nutrition Intervention Research Initiative Food Frequency Questionnaire for Adults*

Food	Note
Biscuits	Made with white flour, frequently served with gravy
Broccoli and rice casserole	Popular casserole that also contains processed cheese
Cheese dip	Commercially processed
Chicken and dumplings, pot pie	Chicken in gravy with flour dumplings or pie crust
Cracklings	Deep fried pork skin
Fried beef (chicken fried steak)	Beef loin, flattened, battered and fried
Fried catfish, fish sandwich	Usually deep fried
Fried potatoes: potato logs	Large pieces of potato, deep fried
Fruit drinks: orangeade, lemonade	Usually commercially prepared, condensed or powdered
Game: venison, squirrel	From local hunters
Gravy (on meat or biscuits)	Usually prepared from a commercial mix
Greens: mustard, turnip, collards, poke salat	Locally popular green vegetables
Grits	Corn-based cereal, cooked like rice
Home-made soup: gumbo	Usually contains okra, rice, sausage and special seasonings
Jambalaya, dirty rice	Rice recipe with meat or fish and special seasonings
Neck bones, ham hock, pig's feet	May be smoked or pickled
Okra	Locally popular vegetable
Organ meats: chitterlings	Fried intestines
Peas: field, black-eyed, purple hull	Locally popular peas
Powdered drink mixes	Mostly sugar with artificial flavour
Root crops: turnips, rutabaga	Locally popular root vegetables
Shellfish: crawfish	River crustacean
Sweet potato pie	Sweet pie made with sweet potatoes, milk and sugar
Sweet tea	Iced tea prepared with sugar

* Foods identified on the 24-hour dietary recalls that contributed at least 0.5% of energy or of any other nutrient examined.

nutrients for this group but could also lead to biased comparisons with non-Hispanic whites¹⁶. In addition to consumption of foods not on the food list, portion sizes of ethnic subgroups can vary widely from that assumed for the general population. Furthermore, unique preparations of commonly used foods by a subgroup may have a major impact on the nutrient content of the diet.

This analysis indicates that the population of the LMD region of the USA does consume many foods not on existing FFQs. Furthermore, within the Delta region, differences in dietary patterns were evident between white and African American residents. For example, white Delta adults consumed considerably more 'sweet tea' (sugar added to hot tea and then iced), while the African American adults were much more likely to consume fruit drinks. The importance of these beverages in this population required reconsideration of their placement on the questionnaire and of the detail required for measuring them correctly. Additionally, cornbread, grits, greens, sausage and liver appeared as important contributors of specific nutrients for the African American, and gumbo for the white adults in the LMD. Consideration of these groups separately ensures that the most important foods for each group are represented in a single FFQ that can accurately capture the diet of both. We found that a questionnaire for the LMD required numerous differences from those nationally used, including several new foods, food names as used in the local context, specific food preparation details, increased portion sizes and locally defined recipes.

One limitation of the data used to develop the food list was the collection of recall information only between January and June. It is therefore possible that some

seasonal foods were underrepresented in the food intake rankings. However, we tested the FFQs during the summer months, and specifically requested whether there were any additional foods that were consumed frequently. While this request led to changes in recipes, it did not lead to the need to add additional foods to the questionnaire.

The concerns with questionnaire completion that we noted in our field tests were similar to those identified by others using cognitive interviewing techniques. For example, based on questionnaire testing, the NCI abandoned the grid format¹⁷. The Delta NRI experience differed, however, as we found that the grid was well accepted for frequency but not for portion size. Our final form, therefore, includes alternating sections of frequency grids, expanded sections for portion size and adjustment questions, allowing for a more natural flow of related information, while obtaining more precise detail on individual food preparations. The inclusion of four rather than three portion size options allowed individuals in the upper tail of the distribution to report their portion sizes more accurately, thereby reducing underestimation of intake of specific foods consumed in large quantity by some individuals. The use of specific numbers for each portion size allowed the respondent to think of their own portion in relation to the closest portion. This was found to be easier and faster than either attempting to calculate from information only for a medium serving or from considering the information included in ranges.

In addition to providing valuable data from which to design the FFQ, the ranking of food items to nutrient intake provides an important look at the diets consumed by population subgroups. The patterns described here

Table 7 Portion size distributions for selected foods (g), as consumed by adults of the Lower Mississippi Delta

Food item	15th percentile	Median	85th percentile
Apple	69	138	212
Bacon	8	16	32
Beef	51	124	276
Beer	360	720	1440
Biscuit	29	51	90
Butter	5	8	16
Cake	18	40	87
Candy	4	13	40
Carrot	8	32	78
Cheese	14	26	49
Coffee	179	253	592
Corn	41	85	164
Cornbread	21	63	112
Eggs	40	80	118
Fruit drink	188	358	579
Fried chicken	45	86	153
Fried fish	75	150	298
French fries	35	70	125
Gravy	14	31	120
Greens	36	88	164
Grits	98	139	242
Ham	21	42	85
Hamburger	68	113	227
Hot dog	27	57	118
Ice cream	59	116	241
Liquor	42	111	348
Orange	83	131	184
Peas	33	88	170
Pie	69	114	198
Pizza	63	163	284
Pork	44	85	186
Potato chips	14	28	74
Rice	44	119	171
Salad dressing	5	15	39
Soda	248	369	553
Soup	112	241	494
Spaghetti	36	107	210
Sweet potato	45	98	150
Sweet tea	237	474	858
Tomato	15	59	184
Tuna	9	80	170
Whole milk	61	203	366
White bread	23	40	60

show that both racial groups in the LMD tended to have higher intakes of soft drinks, other sweetened drinks, fried chicken, fried fish and pork than the general US population¹⁰. Together, soft drinks, white bread, ground beef and salty snacks contribute more than 20% of energy in the LMD population, while few fruits or vegetables appear. The ranking of food sources reveals that when intake is low, as in the example of magnesium among African Americans, the primary source may appear unusual, such as salty snacks. The prominence of white bread and salty snacks in the lists of contributors to several nutrients illustrates that good sources of these nutrients are not included as part of the dietary pattern.

The LMD is one of the poorest regions of the country and the states therein have high prevalences of obesity, hypertension, diabetes, heart disease, cancer and other diet-related conditions^{5,18–20}. An earlier report documented lower intakes of protein and most vitamins

and minerals in this sample relative to a national survey⁶. The results presented here clarify the food choices in these populations and suggest areas for further research and nutrition intervention.

The experience described here illustrates the complexity involved in the development or modification of an FFQ for populations with diets that differ from those already in use. In each case, not only the food items but also the preparation methods, portion sizes and internal variation within the population must be fully considered. Other important considerations include the educational level of the population, their previous exposure (or lack thereof) to research questionnaires, and the clarity and interpretability of the tool, within these contexts.

The development of this regionally specific FFQ for adults will allow future research on diet and health outcomes and will be of value to scientists studying food and nutrient intakes of African Americans and white adults in the Southern United States. Calibration of this instrument, in both a full and abbreviated form, is in progress with a subset of participants in the Jackson Heart Study, where each version will be compared with multiple 24-hour recalls and serum nutrient biomarkers. Once fully calibrated, the use of this FFQ will also facilitate studies of dietary change through interventions, thereby contributing to the increasing evidence that dietary improvements can have a powerful effect on nutritional status and health.

Acknowledgements

This work was supported in part by USDA, Agricultural Research Service Co-operative Agreements 58-6251-2-0008 and 58-1950-9-001.

References

- 1 Kipnis V, Subar AF, Midthune D, Freedman LS, Ballard-Barbash R, Troiano RP, *et al.* Structure of dietary measurement error: results of the OPEN biomarker study. *American Journal of Epidemiology* 2003; **158**: 14–21, discussion 22–6.
- 2 Willett W. *Nutritional Epidemiology*, 2nd ed. New York: Oxford University Press, 1998.
- 3 Block G, Hartman AM, Dresser CM, Carroll MD, Gannon J, Gardner L. A data-based approach to diet questionnaire design and testing. *American Journal of Epidemiology* 1986; **124**: 453–69.
- 4 Willett WC, Sampson L, Stampfer MJ, Rosner B, Bain C, Witschi J, *et al.* Reproducibility and validity of a semiquantitative food frequency questionnaire. *American Journal of Epidemiology* 1985; **122**: 51–65.
- 5 Pickle L, Mungiole M, Jones G, White A. *Atlas of United States Mortality*. Hyattsville, MD: National Center for Health Statistics, 1997.
- 6 Champagne CM, Bogle ML, McGee BB, Yadrick K, Allen HR, Kramer TR, *et al.* Dietary intake in the lower Mississippi delta region: results from the Foods of our Delta Study. *Journal of the American Dietetic Association* 2004; **104**: 199–207.
- 7 Bogle M, Stuff J, Davis L, Forrester I, Strickland E, Casey PH, *et al.* Validity of a telephone-administered 24-hour dietary recall in telephone and non-telephone households in the

