

12-13-2006

Amyloid- β -Induced Pathological Behaviors are Suppressed by *Ginkgo biloba* extract EGb 761 and Ginkgolides in Transgenic *Caenorhabditis elegans*

Yanjue Wu
University of Maryland

Zhixin Wu
University of Southern Mississippi

Peter Butko
University of Southern Mississippi, pbutko@chem.nagoya-u.ac.jp

Yves Christen
Ipsen

Mary P. Lambert
Northwestern University

See next page for additional authors

Follow this and additional works at: https://aquila.usm.edu/fac_pubs

 Part of the [Biology Commons](#)

Recommended Citation

Wu, Y., Wu, Z., Butko, P., Christen, Y., Lambert, M. P., Klein, W. L., Link, C. D., Luo, Y. (2006). Amyloid- β -Induced Pathological Behaviors are Suppressed by *Ginkgo biloba* extract EGb 761 and Ginkgolides in Transgenic *Caenorhabditis elegans*. *Journal of Neuroscience*, 26(50), 13102-13113.

Available at: https://aquila.usm.edu/fac_pubs/8519

Authors

Yanjue Wu, Zhixin Wu, Peter Butko, Yves Christen, Mary P. Lambert, William L. Klein, Christopher D. Link, and Yuan Luo

Amyloid- β -Induced Pathological Behaviors Are Suppressed by *Ginkgo biloba* Extract EGb 761 and Ginkgolides in Transgenic *Caenorhabditis elegans*

Yanjue Wu,¹ Zhixin Wu,² Peter Butko,³ Yves Christen,⁴ Mary P. Lambert,⁵ William L. Klein,⁵ Christopher D. Link,⁶ and Yuan Luo¹

¹Department of Pharmaceutical Sciences, School of Pharmacy, University of Maryland, Baltimore, Maryland 21201, Departments of ²Biological Sciences and ³Chemistry and Biochemistry, University of Southern Mississippi, Hattiesburg, Mississippi 39406, ⁴Ipsen, 75016 Paris, France, ⁵Department of Neurobiology and Physiology, Northwestern University, Evanston, Illinois 60208, and ⁶Institute for Behavioral Genetics, University of Colorado, Boulder, Colorado 80309

Amyloid- β ($A\beta$) toxicity has been postulated to initiate synaptic loss and subsequent neuronal degeneration seen in Alzheimer's disease (AD). We previously demonstrated that the standardized *Ginkgo biloba* extract EGb 761, commonly used to enhance memory and by AD patients for dementia, inhibits $A\beta$ -induced apoptosis in neuroblastoma cells. In this study, we use EGb 761 and its single constituents to associate $A\beta$ species with $A\beta$ -induced pathological behaviors in a model organism, *Caenorhabditis elegans*. We report that EGb 761 and one of its components, ginkgolide A, alleviates $A\beta$ -induced pathological behaviors, including paralysis, and reduces chemotaxis behavior and 5-HT hypersensitivity in a transgenic *C. elegans*. We also show that EGb 761 inhibits $A\beta$ oligomerization and $A\beta$ deposits in the worms. Moreover, reducing oxidative stress is not the mechanism by which EGb 761 and ginkgolide A suppress $A\beta$ -induced paralysis because the antioxidant L-ascorbic acid reduced intracellular levels of hydrogen peroxide to the same extent as EGb 761, but was not nearly as effective in suppressing paralysis in the transgenic *C. elegans*. These findings suggest that (1) EGb 761 suppresses $A\beta$ -related pathological behaviors, (2) the protection against $A\beta$ toxicity by EGb 761 is mediated primarily by modulating $A\beta$ oligomeric species, and (3) ginkgolide A has therapeutic potential for prevention and treatment of AD.

Key words: $A\beta$ peptide; Alzheimer's disease; behavior; mutant; phenotype; serotonin

Introduction

Despite a widely accepted "amyloid cascade hypothesis" for Alzheimer's disease (AD) (Hardy and Selkoe, 2002), the current explanation for amyloid- β ($A\beta$) toxicity in AD still remains controversial in view of the fact that the relationship between $A\beta$ species and $A\beta$ -specific behavior has not been defined *in vivo* (Lorenzo and Yankner, 1994; Pike et al., 1995; Lambert et al., 1998). A leading theory is that $A\beta$ oligomers are responsible for initiation of synaptic dysfunction, an early event that leads to neurodegeneration observed in AD (Walsh and Selkoe, 2004; Roselli et al., 2005; Lesne et al., 2006; Oddo et al., 2006). The evidence for or against these hypotheses is critical for specific therapeutic strategies. It has previously been demonstrated that small molecules inhibiting $A\beta$ oligomers also reduced its toxicity (Walsh et al., 2005; Yang et al., 2005; Maezawa et al., 2006).

However, most of these studies were conducted *in vitro*, and the use of a transgenic mice model of AD for pharmacological evaluation and mechanistic studies is time-consuming. Simple invertebrate models of neurodegenerative diseases offer experimental advantages for addressing basic cellular processes that are conserved among all animals (Link, 2005; Wu and Luo, 2005).

The standard *Ginkgo biloba* leaf extract EGb 761 has been given routinely as a prescription drug in many countries, and as a dietary supplement in the United States, for Alzheimer's dementia (Christen and Maixent, 2002). Several clinical trials have provided evidence of efficacy (Le Bars et al., 1997; Birks et al., 2002; Mix and Crews, 2002; Le Bars, 2003), comparable with Donepezil (Mazza et al., 2006), as a symptomatic treatment of mild to moderate AD, and suggestive for AD prevention (Andrieu et al., 2003). Currently, a National Institutes of Health-supported GEM (Ginkgo Evaluation of Memory) study in the United States and a GuidAge study in Europe are underway to evaluate EGb 761 as a preventive drug (DeKosky et al., 2006). Substantial experimental data indicate that EGb 761 has neuroprotective and neuromodulatory effects (Luo, 2001; Watanabe et al., 2001; Yao et al., 2001; DeFeudis, 2002). Two of its major constituents, flavonoids (24%) and terpenoids (6%), have been actively investigated for their neuroprotective properties (DeFeudis, 1998; Smith and Luo, 2003). The ginkgolides, known as potent antagonists of platelet-

Received Aug. 9, 2006; revised Oct. 12, 2006; accepted Nov. 9, 2006.

This work was supported by National Institutes of Health Grant R01AT001928-03A1 (Y.L.) from the National Center for Complementary and Alternative Medicine and by Ipsen (Paris, France). We thank Dr. Dominic Walsh of University College Dublin (Dublin, Ireland) for helpful discussion and Zhiming Cao for technical assistance.

Correspondence should be addressed to Yuan Luo, Department of Pharmaceutical Sciences, School of Pharmacy, University of Maryland, Baltimore, MD 21201. E-mail: yluo@rx.umaryland.edu.

DOI:10.1523/JNEUROSCI.3448-06.2006

Copyright © 2006 Society for Neuroscience 0270-6474/06/2613102-12\$15.00/0




Figure 1. Chemical structures of the ginkgolides (A), the bilobalide (B), and three flavonol glycosides (C). The ginkgolides and bilobalide are specific to the *G. biloba* tree. Ginkgolides A, B, C, J, and bilobalide, but not ginkgolide M, are constituents of EGb 761. The flavonol glycosides quercetin, kaempferol, and isorhamnetin are present in trace amounts in EGb 761 (DeFuidis, 1998).

-activating factor receptor (PAFR) (Smith et al., 1996) are unique to the *G. biloba* tree (Nakanishi, 1967; Ivic et al., 2003; Jaracz et al., 2004). The flavonoids are involved in antioxidative properties of EGb 761.

We and others previously described an inhibitory effect of EGb 761 on A β oligomerization in solution (Luo et al., 2002; Chromy et al., 2003) providing evidence that EGb 761 can alter the biochemical and biophysical properties of A β that may underpin its toxicity. In the present study, we use the transgenic *Caenorhabditis elegans* model (Link, 1995) that exhibits several pathological behaviors to associate A β species with A β -induced abnormal behaviors in the organism. The current results support the hypothesis that A β oligomers are the toxic species and suggest that beneficial effects of EGb 761 on the cognitive function of AD patients is, at least in part, attributable to its modulation of A β oligomerization.

Materials and Methods

G. biloba extract EGb 761 was provided by Schwabe Pharmaceuticals (Karlsruhe, Germany); the extract is well characterized and is the one used in the ongoing clinic trials. The flavonoid fraction (Fig. 1C) was from Ipsen (Paris France). The individual constituents of EGb 761 ginkgolide A (GA), ginkgolide B (GB), ginkgolide C (GC), ginkgolide J (GJ) (Fig. 1A), and bilobalide (BB) (Fig. 1B) were obtained from Dr. I. A. Khan at the National Center for Natural Products Research (Oxford, MS). Congo red (CR), L-ascorbic acid [vitamin C (VC)] as well as other chemicals were from Sigma (St. Louis, MO). Stock solutions of all chemicals were made either with 100% ethanol or with distilled water only. The final concentration of ethanol did not exceed 0.01% in the food (*Escherichia coli* strain OP50). All chemicals for treatment of experimental animals were added directly to the OP50 food source and the working concentration was equal to 100 μ g/ml for EGb 761, or 10 μ g/ml for single constituents of EGb 761.

C. elegans strains. (For a summary, see Table 1.) The wild-type strain

Table 1. Description of the well characterized transgenic *C. elegans* used in the study

Strains	Transgene	Expression	Phenotype
N2		Wild type	Wt movement
CL4176	<i>myo-3/Aβ_{1–42}</i>	Inducible muscle A β _{1–42}	Rapid paralysis
CL1175	<i>myo-3</i>	Control for CL4176	Normal
CL2006	<i>unc-54/Aβ_{1–42}</i>	Constitutive muscle A β _{1–42}	Progressive paralysis
CL2355	<i>snb-1/Aβ_{1–42}</i>	Inducible neuronal A β _{1–42}	Reduced chemotaxis Hypersensitive to 5-HT
CL2122	<i>mtl-2/GFP</i>	Control for CL2355	Wt movement
CL2179	<i>myo-3/GFP</i>	GFP control for CL4176	Wt movement

Wt, Wild type.

N2 (Bristol) was from *Caenorhabditis* Genetics Center (University of Minnesota, Minneapolis, MN). The construction and characterization of the transgenic nematode strains CL2006 (*unc-54/A β _{1–42}*) and CL4176 (*smg-1^{ts} [myo-3/A β _{1–42} long 3′-untranslated region (UTR)]*) and its control strain CL1175 (*smg-1^{ts}*) have been described previously (Link, 1995; Link et al., 2001). The CL2006 strain constitutively produces a body-wall muscle-specific A β _{1–42}, whereas the expression of muscle-specific A β _{1–42} in CL4176 depends on upshifting temperature from 16 to 23°C. The transgenic arrays in the CL2006, CL1175, and CL4176 strains all contain the dominant *rol-6(su1066)* morphological marker. The CL2355 strain [*smg-1^{ts} (snb-1/A β _{1–42}/long 3′-UTR)*] employs the promoter of the *C. elegans* synaptobrevin ortholog (*snb-1*) to drive pan-neuronal expression of A β _{1–42}, which is inducible by temperature upshift. The transgenic arrays in CL2355 and the control strain CL2122 are marked with a *mtl-2/green fluorescent protein (GFP)* transgene that causes constitutive intestinal GFP expression. An integrated *myo-3/GFP/long 3′-UTR* strain (CL2179) is used as a control for CL4176. This strain has a GFP reporter tagged to the same promoter as the transgenic strains and has detectable GFP at 16°C, which is significantly stronger at higher temperatures.

C. elegans maintenance and treatment. The wild-type (N2) and the transgenic CL2006 were propagated at 20°C, CL4176 and CL2355 and their controls at 16°C, on solid nematode growth medium (NGM) seeded with 100 μ l spots of *E. coli* (OP50) for food. To prepare age-synchronized animals, nematodes were transferred to fresh NGM plates on reaching reproductive maturity at 3 d of age and allowed to lay eggs for 4–6 h (overnight for Western blotting of CL4176 and CL2355). Isolated hatchlings from the synchronized eggs (day 1) were cultured on fresh NGM plates in either a 20°C or a 16°C (for CL4176, CL2355, and their control strains) temperature-controlled incubator (model 2005; Sheldon Manufacturing, Cornelius, OR). The worms were fed with the drugs either from stage L1 (1 d of age) or starting from the egg.

Paralysis assays. The strain CL4176 (Drake et al., 2003; Link et al., 2003) maintained at 16°C was egg-synchronized onto the 35 × 10 mm culture plates containing either a vehicle or drug. Transgene expression was induced by upshifting the temperature from 16 to 23°C, started at the 36th hour after egg laying and lasted until the end of the paralysis assay. Paralysis was scored at 1 h intervals until the last worm became paralyzed.

Chemotaxis assays. Chemotaxis assays (see Fig. 3A) were performed as described by Bargmann et al. (1993). Synchronized transgenic *C. elegans* CL2355 and its control strain CL2122 were treated with or without 100 μ g/ml EGb 761, or 10 μ g/ml single components, starting from the egg. They were cultured in 16°C for 36 h, and then in 23°C for another 36 h. The worms were then collected, washed with M9 (Wood, 1988) three times, and assayed in 100 mm plates containing 1.9% agar, 1 mM CaCl₂, 1 mM MgSO₄, and 25 mM phosphate buffer, pH 6.0. Twenty worms were placed to the center of the plate. After all animals were transferred to the center of the assay plate, 1 μ l of odorant (0.1% benzaldehyde in 100% ethanol) (Sigma) along with 1 μ l of 1 M sodium azide were added to the original spot. On the opposite side of the attractant, 1 μ l drop of sodium azide and 1 μ l of control odorant (100% ethanol) were added. Assay plates were incubated at 23°C for 1 h and chemotaxis index (CI) was scored. Chemotaxis index is defined as follows: (number of worms at the attractant location – number of worms at the control location)/total number of worms on the plate.

5-HT sensitivity assay. Synchronized transgenic worms (CL2355) and the control strain (CL2122) fed with or without the drugs were collected at 36 h after temperature upshift. Serotonin (creatinine sulfate salt; Sigma) was dissolved in M9 buffer to 1 mM. Twenty worms in each group were washed with M9 buffer for three times and transferred into 200 μ l serotonin solution in a 96-well assay plate. Worms were scored after 5 min as active or paralyzed (nonmotile for 5 s).

Western blotting of A β species. The A β species in the transgenic *C. elegans* strains was identified by immunoblotting using a Tris-Tricine gel and the standard Western blotting protocol except that the polyvinylidene difluoride membranes were boiled for 5 min after the transfer. After the experimental treatments, the worms were collected by washing with distilled water, quickly frozen in liquid nitrogen, sonicated in the cell lysis buffer (50 mM HEPES, pH 7.5, 6 mM MgCl₂, 1 mM EDTA, 75 mM sucrose, 25 mM benzamide, 1 mM DTT, 1% Triton X-100), and heated with sample buffer containing 5% β -mercaptoethanol (1:1; Bio-Rad, Hercules, CA). To detect A β oligomers, samples were extracted in PBS containing proteinase inhibitor mixture (Sigma) with (for antibody A11) or without 2% SDS (for antibody NU-4). After mixing with the sample buffer, proteins were unheated and loaded on the gel. Equal amounts of the total protein (80 μ g) were loaded in each lane. Antibody to A β _{1–17} (6E10; at 1:500 dilutions) was from Signet (Dedham, MA). Antibody selective against oligomers (A-11; at 1:1000 dilution) was from Biosource (Camarillo, CA) (Kayed et al., 2003). Antibody specific to A β oligomers (NU4; 1:3000) was generated in W. L. Klein's laboratory (Lambert, 2006). Anti-mouse IgG or anti-rabbit IgG (1:5000; Signet) was used as the secondary antibody. The mean densities of the A β bands were analyzed by a gel documentation system (Alpha Innotech 801054; Imgen Technologies, Alexandria, VA).

Fluorescence staining of A β deposits. Individual CL2006 transgenic nematodes were fixed in 4% paraformaldehyde/PBS, pH 7.4, for 24 h at 4°C, and permeabilized in 5% fresh β -mercaptoethanol, 1% Triton X-100, 125 mM Tris, pH 7.4, in a 37°C incubator for 24 h. The nematodes were stained with 0.125% thioflavin S (Sigma) in 50% ethanol for 2 min, destained, and mounted on slides for microscopy. Fluorescence images were acquired at the same exposure parameters using a 40 \times objective of the microscope (BX 60; Olympus, Tokyo, Japan) equipped with a digital camera (Micropublisher 5.0; QImaging, Burnaby, British Columbia, Canada). The number of thioflavin S-reactive deposits in the area anterior of the pharyngeal bulb in individual animals was scored.

H₂O₂ assay in *C. elegans*. Intracellular ROS were measured in *C. elegans* using 2,7-dichlorofluorescein diacetate (DCF-DA) (Invitrogen, Eugene, OR) as described previously (Smith and Luo, 2003). Age-synchronized *C. elegans* were collected at 36 h after temperature upshift (CL4176/1175) into 100 μ l of PBS with 1% Tween 20 (PBST) in Eppendorf tubes in groups of 40 worms. The worms were then subjected to equally timed homogenization (Pellet Pestle Motor; MG Scientific, Pleasant Prairie, WI) and sonication (Branson Sonifier 250; VWR Scientific, Suwanee, GA) to break up the outer cuticle. Samples were collected




Figure 2. Effect of EGb 761 and its constituents on A β -induced paralysis in muscle A β strain CL4176. **A**, Representative images of A β -induced paralysis in the transgenic CL4176 strain without transgene expression (no A β), untreated with EGb 761 (Ctrl; left panel), and in the transgenic CL4176 strain (muscle A β strain) fed with (EGb; right) or without EGb 761 (Ctrl; middle) at 36 h after temperature upshift. **B**, Time course of paralysis assays in CL4176 fed with different drugs. Synchronized eggs of CL4176 *C. elegans* were maintained at 16°C, on the 35 \times 10 mm culture plates (~35 eggs/plate) containing vehicle (Ctrl), EGb 761 (100 μ g/ml), or CR (139 μ g/ml). The hatched worms were grown for 38 h at 16°C followed by upshifting the temperature to 23°C to induce the transgene expression. The paralysis was scored at 60 min intervals. Data are expressed as percentage of nonparalyzed worms from at least three independent assays of 100 worms in each experiment. Paralysis in the transgenic strain CL4176 is attributable to A β expression compared with the control strain CL1175, which does not express A β transgene (filled triangles). **C**, Time course of paralysis assay in CL4176 worms fed with ginkgolide A or J (10 μ g/ml for each). Paralysis was scored at 30 min intervals. Data were obtained from three different experiments with 100 worms in each group. Error bars indicate SEM. **D**, An illustrative diagram indicating the time duration of temperature upshift for expressing A β transgene, onset of paralysis, detectable A β , and EGb protective effect at different treatment regimens.

into wells of 96-well plates. DCF-DA (final concentration, 50 μ M in PBS) was added to each well at 37°C for quantification of fluorescence in an FLx800 Microplate Fluorescence Reader (Bio-Tek Instruments, Winooski, VT) with the excitation at 485 nm and emission at 530 nm. Samples were read every 20 min for 2.5 h.

Statistical analyses. Differences between untreated and drug-treated groups were analyzed for statistical significance by independent Student's *t* test of two groups using Origin 6.0 software (Microcal Software, Northampton, MA). A value of *p* < 0.05 is considered statistically significant. Correlation analysis was performed with the GraphPad (San Diego, CA) Prism 4.0a, using a one-tailed Pearson test.

Table 2. Quantitative analysis of paralysis

Experiments	PT ₅₀	<i>p</i> value
Control	5.7 ± 0.8	
Egb 761	9.4 ± 1.3*	0.03
Ascorbic acid	3.4 ± 0.3	0.17
Congo red	7.5 ± 0.8	0.15
Control	3.5 ± 0.3	
Ginkgolide A	7.2 ± 0.3**	0.001
Ginkgolide B	2.8 ± 0.4	0.18
Ginkgolide C	3.5 ± 0.3	1.00
Ginkgolide J	6.8 ± 0.6*	0.01
Bilobalide B	2.8 ± 0.4	0.18
Flavonoids	2.7 ± 0.6	0.24

The paralysis assays were quantified for mean time duration at which 50% worms were paralyzed (PT₅₀) from the transgenic worms fed with or without the drugs. Values of *p* were obtained from nine independent assays for the worms fed with EGb 761 or Congo red, each paired with untreated controls, and from three assays for L-ascorbic acid, and single components paired with the control.

Results

Egb 761, ginkgolide A and J alleviate A β -induced paralysis in the transgenic *C. elegans*

To determine whether EGb 761 specifically protects against A β -induced toxicity *in vivo*, we first fed EGb 761 to a transgenic *C. elegans* line, in which expression of human A β peptide in the muscle cells induces an A β -dependent paralysis phenotype in the worms (muscle A β strain, CL4176) (Link et al., 2003). Synchronized eggs from transgenic worms, or control worms were placed on OP50 food containing vehicle or EGb 761 (100 μ g/ml) for 36 h at 16°C, followed by temperature upshift from 16 to 23°C to induce transgene expression. Figure 2A is a set of photographs representing the following: the transgenic muscle A β expression strain (CL4176) without the temperature upshift and untreated with EGb 761 [control (Ctrl), no A β] (Fig. 2A, left panel), the temperature upshifted (for 36 h) CL4176, untreated with EGb 761 (Ctrl, muscle A β) (Fig. 2A, middle panel), or treated with EGb 761 (EGb, muscle A β) (Fig. 2A, right panel). Worms that did not move or only moved the head, under a gentle touch with a platinum loop were scored as paralyzed. In Figure 2A, the shape of movement and tracks left on the food (indicated with black lines) in nonparalyzed worms are obvious. The arrowheads indicate paralyzed worms (straight line shape), and the arrows indicate the moving worms (the “C” or “S” shape). At 36 h, ~100% worms without the transgene expression were nonparalyzed (Fig. 2A, left). At this time, only ~20% A β transgene expressing worms without EGb feeding were nonparalyzed (Fig. 2A, middle), and ~60% transgene expressing worms fed with EGb were nonparalyzed (Fig. 2A, right). Note the tracks left behind on the plate in the EGb 761-fed *C. elegans* (Fig. 2A, right panel).

Figure 2B is a time course of paralysis in the *C. elegans* CL4176 fed with a vehicle (0.01% ethanol; open circles), EGb 761 (EGb, 100 μ g/ml; filled circles), or CR (139 μ g/ml; filled squares) from day 1 till after the temperature upshift for 36 h. CL1175 (filled triangles) is a transgenic control strain that does not express A β . A notable delay of paralysis was observed in the transgenic worms fed with EGb 761 compared with the untreated controls. Congo red also moderately delayed paralysis. Interestingly, the worms fed with 100 μ g/ml L-ascorbic acid, a known antioxidant, did not show any delay in paralysis (for clarity of the graph, data are presented only in Table 2). To further determine which constituent(s) of EGb 761 contributed to reducing A β -induced paralysis, transgenic *C. elegans* CL4176 were fed with GA, GB, GC, GJ, BB, and a flavonoid (Flav) fraction (10 μ g/ml each). Among individual components tested, only GA and GJ exhibited signifi-

cant delay of A β -induced paralysis (for clarity, the paralysis time courses for other components are not shown in Fig. 2C) (for statistical analysis, see Table 1). Reproducibility of paralysis at a given temperature in individual trials was consistent. Because the transgene product is temperature sensitive, the onset of paralysis is related to the specific value of room temperature, at which the paralysis assays were conducted. Thus, the variation in the onset of paralysis between Figure 2, B and C, is attributed to the slight difference in room temperature on the days when the assays were performed (22 vs 23°C). Nevertheless, the difference between the worms fed with and without EGb 761 was consistently significant at both 22 and 23°C.

For quantitative analysis, we define PT₅₀ as the time interval, from the onset of paralysis, at which 50% of the worms were paralyzed (i.e., PT₅₀ of 5.7 h for control was obtained by subtracting the onset time of paralysis 27 h from the time when 50% of the worms were paralyzed at 32.7 h) (Table 1). A statistically significant delay of A β -induced paralysis was observed in the worms fed with EGb 761 (Table 2) (control vs EGb, *p* = 0.03; *n* = 9 assays; 40 worms in each assay group), but not in Congo red-treated animals (control vs CR, *p* = 0.15; *n* = 9 assays; 30 worms in each assay group). Among six single components of EGb 761, only ginkgolides A and J exhibited a statistically significant delay of paralysis (control vs GA, *p* = 0.001; control vs GJ, *p* = 0.01; *n* = 3 assays; 30 worms in each assay group). Moreover, EGb 761 feeding was found to delay paralysis through a wide range of concentrations (10–500 μ g/ml) (data not shown). Paralysis was not delayed in the worms fed with EGb 761 before A β induction, at the time of A β induction, or after A β induction by temperature upshift (data not shown) (see diagram in Fig. 2D), suggesting that short duration of feeding may not be sufficient to alleviate the A β toxicity.

Egb 761 suppresses neuronal A β expression-induced defect in chemotaxis behavior and 5-HT sensitivity

The paralysis phenotype in muscle of the A β expression strain CL4176, used above, has been established and used to illustrate several important molecular processes related to A β toxicity (Drake et al., 2003; Link et al., 2003). To relate A β toxicity with neurological functions, we characterized a behavior phenotype of the transgenic strain in which A β was expressed in the neuronal cells (CL2355). Two characteristic neuronal controlled behaviors, chemotaxis and 5-HT sensitivity, were assayed in these worms. The chemotaxis response in *C. elegans* is mediated by activation of several sensory neurons and interneurons to stimulate the motor neurons (Hobert, 2003). The CI is a measure of the fraction of worms that are able to arrive at the location of the attractants (Matsuura et al., 2005). To investigate the effect of EGb 761 on the performance of chemotaxis behavior, we applied the chemical benzaldehyde as an attractant and ethanol as a control, both containing sodium azide, which paralyzes the worms on contact (Fig. 3A). The chemotaxis index was scored in the transgenic strain and the control strain at day 4 of age. Figure 3B shows that the transgenic mutant CL2355 exhibits a significant reduction of CI compared with the control strain CL2122, or no A β strain untreated (Ctrl CI_{no A β} , 0.36 ± 0.02 vs Ctrl CI_{A β} , 0.27 ± 0.02; *n* = 4; *p* = 0.01). EGb 761 feeding of the control strain had no effect on their CI (Ctrl CI_{no A β} , 0.36 ± 0.02, vs EGb CI_{no A β} , 0.35 ± 0.04; *n* = 4; *p* = 0.31), but it significantly normalizes the reduced CI in the neuronal A β transgenic strain (Ctrl CI_{A β} , 0.27 ± 0.02, vs EGb CI_{A β} , 0.37 ± 0.03; *n* = 4; *p* = 0.01; a total of 240 worms was used in each assay). A similar effect was observed

in the transgenic worms fed with GA (Ctrl CL_{A β} , 0.27 ± 0.02 , vs GA CL_{A β} , 0.34 ± 0.01 ; $n = 4$; $p = 0.03$).

Taking advantage of the ability of *C. elegans* to take up exogenous 5-HT (Sawin et al., 2000), we conducted a serotonin (5-HT) hypersensitivity assay to further determine whether 5-HT-mediated neurotransmission is affected by expressing A β transgene in the neurons. 5-HT is a key neurotransmitter that modulates several behaviors of *C. elegans*, including egg laying, locomotion, and olfactory learning (Schafer and Kenyon, 1995; Sawin et al., 2000; Nuttley et al., 2002; Zhang et al., 2005). When exogenous 5-HT is applied to the nematodes, they become paralyzed as a result of the sensitivity to excessive 5-HT. This 5-HT sensitivity assay has previously been used to identify 5-HT hypersensitive mutants, which revealed the functional relationship of the genes involved in 5-HT signaling (Schafer et al., 1996; Ranganathan et al., 2001). We used this assay to test the response to 5-HT in the transgenic mutant CL2355 in comparison with the control strain CL2122. Twenty animals from each strain were placed in 200 μ l of 5-HT solution (1 mM) in 96-well microtiter wells. Paralysis was scored at 5 min after exposure to 5-HT. Figure 3C demonstrates that the transgenic neuronal line CL2355 untreated (Ctrl) was particularly sensitive to 5-HT, compared with the transgenic control CL2122 (no A β strain) untreated (percentage of Ctrl_{CL2355}, $35.9 \pm 7.8\%$, vs percentage of Ctrl_{CL2122}, 73.2 ± 3.3 ; $n = 3$; $p < 0.001$). The neuronal A β worms (CL2355) then were fed with EGb 761 or its single components. Only feeding with EGb 761 (EGb vs Ctrl, $69.5 \pm 4.9\%$, vs $35.9 \pm 7.8\%$; $n = 3$; $p = 0.003$) and ginkgolide A (GA vs Ctrl, 65.5 ± 4.0 , vs $35.9 \pm 7.8\%$; $n = 3$; $p = 0.005$) significantly normalized the defective response to 5-HT in the transgenic strain. Neither the antioxidant L-ascorbic acid (VC), kaempferol, the flavonoid fraction of EGb 761 (KA), or CR normalized the paralysis in the neuronal A β strain CL2355.

Among behavior assays available to examine neuronal toxicity (including egg laying, body bend, pharyngeal pumping, locomotion/swimming, and paralysis), the serotonin sensitivity assay (swimming) was performed to determine a behavior phenotype in neuronal A β -expressing strain for the following reasons: (1) early work by Horvitz et al. (1982) demonstrated that exogenous 5-HT affected the wild-type *C. elegans* locomotion, or swimming behavior. Subsequently, it was reported that the effect was blocked in the MOD-1 mutant (a 5-HT-gated chloride channel) (Ranganathan et al., 2000), suggesting that this assay measures a 5-HT-mediated response; (2) in *C. elegans*, the experience-dependent behavior (an enhanced slowing of the response) is modulated by 5-HT (Sawin et al., 2000), which provides a link between the behavior plasticity and the neuronal plasticity associated with the impairment of cognitive function observed in AD; (3) among others, this phenotype exhibited the most significant difference between the control strain (CL2122) and the transgenic strain (CL2355). Thus, the A β -dependent behavior may provide the opportunity to elucidate 5-HT-mediated neuroprotection against A β neurotoxicity.

EGb 761 modulates A β oligomers in the transgenic *C. elegans*

To determine whether inhibition of A β oligomerization by EGb 761 underpins its mechanism for suppressing pathological behaviors in *C. elegans*, we analyzed A β species from the transgenic *C. elegans* fed with or without EGb 761 by Western blotting using antibodies against A β (6E10) or specific against A β oligomers (A-11 and NU-4). Figure 4A shows A β -immunoreactive (6E10) bands (M_r , ~ 7.2 – 28 kDa) detected in the tissues from the transgenic worms (CL4176) fed with or without EGb 761 and the




Figure 3. Assays for chemotaxis behavior and serotonin sensitivity in neuronal A β -expressing strain CL2355. **A**, Schematic diagram of the chemotaxis assay. Synchronized L4 worms treated with or without drugs (20 worms for each assay) were placed in the center of an assay plate (100 \times 15 mm) containing 1 μ l of attractant (0.1% benzaldehyde) plus 1 μ l of 1 M sodium azide at the edge of the plate on one side (black dot), and 1 μ l of control odorant (100% ethanol, *) plus 1 μ l of 1 M sodium azide on the opposite edge of the plate (star). After 1 h incubation at room temperature, CI was scored as follows: CI = (number of worms at attractant – number of worms at control)/total number of worms. In each experiment, at least 60 worms from each group were analyzed. **B**, Chemotaxis behavior in neuronal strain CL2355 (neuronal A β strain) was reduced compared with the transgenic control strain CL2122 (no A β strain). Feeding with 100 μ g/ml EGb 761 for 4 d alleviated this descent in the transgenic strain, but not in the control strains ($n = 4$; * $p < 0.05$). Error bars indicate SEM. **C**, Serotonin hypersensitivity in neuronal-expressing A β strain (CL2355) is normalized by EGb 761 and its constituents. Strain CL2355 were fed with EGb 761 (100 μ g/ml), vehicle (Ctrl; 100 μ g/ml), ginkgolides (GA, GB, GC, GJ) (10 μ g/ml each), BB (10 μ g/ml), quercetin (QC) (10 μ g/ml), kaempferol (KA; 10 μ g/ml) L-ascorbic acid (VC) (100 μ g/ml), and CR (139 μ g/ml) for 4 d. The worms were collected at 36 h after temperature upshift to 23°C. 5-HT sensitivity assay was conducted in 96-well plate containing 200 μ l of 1 mM 5-HT solution. Percentage of active worms is calculated as follows: number of the active worms (after placed into serotonin solution for 5 min and still kept moving for 5 s)/total worms. A total of 60 worms of each group was used for each time ($n = 3$; ** $p < 0.01$).




Figure 4. A β species in the transgenic *C. elegans*. **A**, Representative Western blot (BL) of A β species in the transgenic *C. elegans* CL4176 fed with or without EGb 761 (EGb), L-ascorbic acid (VC), and CR. The CL4176 worms maintained at 16°C were fed with a vehicle (Ctrl), EGb 761 (100 μ g/ml), or L-ascorbic acid (100 μ g/ml), Congo red (139 μ g/ml) at day 1 of age for 72 h. The worms were collected, and equal amounts of protein were loaded on each gel lane and immunoblotted with an anti-A β antibody (6E10). The arrow indicates changes of A β oligomers and the monomers in the worms after the treatment. **B**, Quantification of immunoreactive A β oligomers (the bend at 20 kDa; filled bars) and A β monomers (dashed bars) by Gel Documentary System Fluochem SP (Alpha Innotech, San Leandro, CA). Data are expressed as mean density of an indicated band from three independent experiments. Error bars represent SEM. * p < 0.05. **C**, A β species in transgenic *C. elegans* CL4176 fed with or without EGb 761 (100 μ g/ml), GA, GB, GC, GJ, BB, and a flavonol diglycosides quercetin (QC) (10 μ g/ml for each component). A β oligomers (top) and the protein loading control actin (bottom) are indicated. Note that immunoreactive A β monomers (arrow on the right) are enhanced in worms fed with GA. **D**, Quantification of immunoreactive A β oligomers (solid bars) and A β monomers (dashed bars) from three independent experiments. Error bars represent SEM. * p < 0.05; ** p < 0.01; # p < 0.05 (4 kDa monomer). **E**, Immunoblot using antibody A11 (anti-oligomers) in the *C. elegans* strains CL4176 fed with EGb 761 (100 μ g/ml) or vehicle (Ctrl) for 4 d. The blot represents three independent experiments. **F**, Immunoblot using NU-4 (anti-A β oligomers) antibody in *C. elegans* strains CL4176 and CL2355 (-Tg) fed with EGb 761 (100 μ g/ml) or vehicle (Ctrl). The blot represents two independent experiments. **G**, An integrated *myo-3*/GFP strain (CL2179), as a control for CL4176, was fed with (bottom panel) or without (top) EGb 761 in the same way as CL4176 in **A**. GFP fluorescence was examined at 36 h after temperature upshift, under a 20 \times objective on the fluorescence microscope (E800; Nikon). The images were processed by IPLab software 3.7 (Scanalytics, Fairfax, VA). No difference in GFP fluorescence density and intensity were observed in the two groups.

comparison chemicals. Two immunoreactive A β species with molecular weights at \sim 20 and 28 kDa were significantly decreased in the worms fed with EGb 761 (100 μ g/ml for 72 h) (Fig. 4A, EGb, lane 3). Interestingly, a simultaneous increase of the A β monomers was observed in those worms. Consistent with the

observation in CL4176, a similar inhibitory effect by EGb 761 was observed in the transgenic strain CL2006. In this strain, EGb 761 inhibited the same A β species and, at the same time, increased A β monomers in the same tissue extract (data not shown). In contrast, L-ascorbic acid (VC) (100 μ g/ml for 72 h) did not inhibit the A β oligomerization or increase the monomer content as EGb 761 did, in the transgenic worm CL4176 (Fig. 4A, lane 4). CR (139 μ g/ml) inhibited the oligomerization to a lesser degree than EGb 761 but also failed to increase the A β monomer content in the same samples in the transgenic worms (Fig. 4A, CR, lane 5). The mean densities of the A β oligomer band at \sim 20 kDa as well as the A β monomer bands were analyzed. Statistically (Fig. 4B), EGb 761 and Congo red significantly reduced the oligomers (Fig. 4B, solid bars) (n = 3; control vs EGb, p = 0.04; Ctrl vs CR, p = 0.02). Most importantly, only EGb 761 significantly increased the A β monomers from the same *C. elegans* samples (Fig. 4B, dashed bars) (n = 3; control vs EGb, p = 0.01).

Neuroprotection by EGb 761 is known as a “multipotent” action (i.e., it is achieved by modulating a variety of biological pathways simultaneously because of its complex nature) (DeFeudis, 2002; Smith and Luo, 2003). We next asked whether the inhibitory effect on oligomerization by an individual constituent is sufficient to correlate with its alleviative effect on A β -induced paralysis. The transgenic *C. elegans* was fed with different single constituents and A β species were analyzed by Western blotting using antibody 6E10. Figure 4C represents a noticeable shift of A β oligomer species to monomers by EGb 761 and ginkgolide A. Blots from three independent experiments were quantified in Figure 4D. It shows that EGb 761 (EGb), GA, and GJ remarkably reduced the A β oligomers band (filled bars) (Ctrl vs EGb, ** p = 0.006; Ctrl vs GA, ** p = 0.006; Ctrl vs GJ, ** p = 0.007). GB, GC, and GJ also significantly reduced this A β species, to a somehow lesser degree (Ctrl vs GB, * p = 0.04; Ctrl vs GC, * p = 0.04), compared with untreated transgenic worms (Ctrl). Only EGb 761 and GA significantly enhanced the A β monomers (dashed bars) (Ctrl vs EGb, # p = 0.02; Ctrl vs GA, # p = 0.03).

To confirm the properties of the A β species modulated by EGb 761, we used two independent antibodies A11 (Fig. 4E) and NU4 (Fig. 4F). Antibody A11 is selective for high molecular weight A β oligomers as well as other oligomeric proteins (Kayed et al., 2003). This antibody was used to reveal a temporal profile of A β oligomerization in mice model of

AD (Oddo et al., 2006). Antibody NU-4 is a mice monoclonal antibody generated in the W. L. Klein laboratory (Lambert, 2006), which is an improved version of the previously generated low molecular weight oligomer [A β -derived diffusible ligand (ADDL)]-selective polyclonal rabbit antibody (M93/M94) (Lambert et al., 2001). Levels of oligomers reactive to this antibody have been reported to be 70-fold higher in the AD patient's brain compared with the control brains (Gong et al., 2003). Prevention by EGb 761 of the oligomer formation *in vitro* was demonstrated using this antibody (Chromy et al., 2003).

Multiple immunoreactive bands were recognized by A11 in the transgenic *C. elegans* CL4176 (Fig. 4E, lane 2). Among these, a major band at ~50 kDa disappeared in the CL4176 strain fed with EGb 761 (Fig. 4E, lane 3). The blot is representative of three independent experiments. Because A11 also recognizes oligomeric structures other than A β , we further performed the immunoblotting using an antibody specific to the toxic A β oligomers referred to as ADDL (NU-4). Figure 4F is representative of two independent blots using antibody NU-4. An immunoreactive band at 50 kDa, presumably a 12-mer species previously found to be abundant in AD brain (Gong et al., 2003), was detected only in the transgenic strain CL4176, but not in the transgenic control strain. EGb 761 inhibited formation of this species completely (Fig. 4F, lane 3). EGb 761 also inhibited formation of higher-order A β species (~70–80 kDa, probably 16- to 18-mers) in the neuronal strain CL2355. In addition, the A β species detected by 6E10 and A11/NU4 do not overlap, in terms of the size, which suggests that multiple A β oligomers are attenuated by EGb 761. The shift from larger A β oligomers to monomers caused by EGb 761 is striking, but it only can be detected by 6E10, because A11 and Nu-4 do not recognize monomeric A β (Chromy et al., 2003; Lambert, 2006).

To determine whether EGb 761 affects the expression levels of the A β transgene, rather than interacting with the A β peptide species, an integrated *myo-3*/GFP strain (CL2179) with the same promoter tagged with a GFP reporter was used as a control for CL4176. To determine whether EGb 761 affects transcription of the transgenes, the strain was handled the same way as CL4176, in terms of temperature upshift regimen and EGb 761 feeding. Figure 4G shows that comparing the worms (CL2179) fed with or without EGb, there is no visible difference in GFP fluorescence density and intensity in the two groups, suggesting that EGb 761 feeding does not affect the levels of A β transgene expression in the worms. Data from a microarray of the transgenic worms fed with or without EGb 761 further supported this observation (data not shown).

EGb 761 inhibits amyloid deposits in transgenic *C. elegans*

To decide whether the inhibitory effect of EGb 761 on A β oligomerization would affect amyloid formation, the number of amyloid deposits was scored in the worm head region, which is separated from the rest of the body by the pharyngeal bulb (black arrows). Figure 5A shows A β deposits (black arrowheads) detected in the transgenic *C. elegans* (CL2006) (Fig. 5Ac) but not the wild type (N2) (Fig. 5Ab), as previously observed (Link, 1995). The number of A β deposits per nematode was reduced in the transgenic *C. elegans* CL2006 fed with EGb 761 (Fig. 5Ad). The worms exhibit background fluorescence without thioflavin S staining (Fig. 5Aa), which provides a guide to define the head area in the animals. Figure 5B shows that the mean number of A β deposits, from 24 worms in each group, was significantly reduced in worms fed with 100 μ g/ml EGb 761 (Ctrl, 7.7 ± 0.5 , vs EGb, 4.9 ± 0.6 ; $n = 24$; $p = 0.002$) or with 200 μ M Congo red (Ctrl,




Figure 5. A β deposits in transgenic *C. elegans* CL2006 fed with or without drugs. **A**, Representative images of background fluorescence in a *C. elegans* without staining (**a**), with thioflavin S staining in the wild type (**b**), or in the transgenic strain CL2006 fed with (**d**) or without EGb 761 (**c**). Synchronized CL2006 worms were maintained at 20°C on NGM agar plates seeded with *E. coli* as the food source. The worms were fed with EGb 761 or Congo red for 2 d starting at day 4 of age. At the end of the treatment, the nematodes were fixed in 4% paraformaldehyde, permeabilized, and then stained with 0.125% thioflavin S. β -Amyloid deposits were examined using a fluorescence microscope attached to a digital camera. The numbers of deposits (arrowheads) were scored in the worm head, which is separated from the body by pharyngeal bulb (arrows). **B**, Quantitative analysis of β -amyloid deposits in the transgenic *C. elegans* CL2006 fed with different chemicals for 48 h (EGb 761, 100 μ g/ml; the flavonoid fraction of EGb 761, 100 μ g/ml; Congo red, 139 μ g/ml). The quantity is expressed as mean number of β -amyloid deposits/anterior area of the worm ($n = 24$ for each analysis). Error bars indicate SEM. * $p < 0.05$; ** $p < 0.01$.

6.2 ± 0.8 , vs CR, 5.5 ± 1.1 ; $n = 24$; $p = 0.025$), but not with the flavonoid fraction of EGb 761 (Ctrl, 7.1 ± 0.9 , vs Flav, 6.9 ± 1.0 ; $n = 24$; $p > 0.05$). These results suggest that EGb 761 inhibits A β oligomerization, which leads to an increase in the nontoxic A β monomers and the reduced amyloid deposits. Congo red, which binds to A β fibrils, also reduced the A β deposits in *C. elegans*.

Levels of H₂O₂ in the transgenic *C. elegans* fed with EGb 761 and other chemicals

Given that numerous lines of evidence have associated oxidative stress with AD and A β toxicity, we hypothesized that the antioxidative properties also contribute to the protective effects of EGb 761 against A β toxicity. We first tested the levels of H₂O₂ in the transgenic control strain (CL1175) and the transgenic muscle A β strain (CL4176). Figure 6A demonstrates that the untreated transgenic strain CL4176 (Ctrl) exhibits increased levels of H₂O₂




Figure 6. Levels of ROS in transgenic *C. elegans*. **A**, Levels of H₂O₂ in the *C. elegans* fed with different concentration of EGb 761 (0, 25, 50, 100 μ g/ml). Age-synchronized groups of transgenic *C. elegans* strain CL4176 and the control strain CL1175 maintained at 16°C for 38 h were temperature upshifted to 23°C for 38 h followed by the DCF assay for H₂O₂ described in Materials and Methods. Statistical significance is observed in the worms fed with 100 μ g/ml EGb 761 (25 μ g/ml, $p = 0.13$; 50 μ g/ml, $p = 0.09$; 100 μ g/ml, $p = 0.03$; $n = 3$; total of 120 worms in each group). **B**, Levels of H₂O₂ in CL4176 worms fed with vehicle (Ctrl), EGb 761 (EGb), L-ascorbic acid (VC), or CR from 1 d of age until 3 d of age. **C**, Levels of H₂O₂ in the *C. elegans* fed with different constituents of EGb 761 (ginkgolides GA, GB, GC, GJ). At least 60 animals from each group were analyzed for the levels of H₂O₂. Results are expressed as percentage of fluorescence (%DCF) relative to vehicle-treated controls, which is set as 100%. Statistical significance is represented as follows: * $p < 0.05$; ** $p < 0.01$. Error bars indicate SEM.

compared with the control strain (CL1175) that does not express A β (no A β strain, $p = 0.001$). Next, we asked whether feeding EGb 761 to the worms before the induction of A β expression would prevent the elevation of H₂O₂ levels in the strain CL4176.

Three concentrations of EGb 761 (25, 50, and 100 μ g/ml) were added to the transgenic *C. elegans* diet from day 1 of age until the end of the temperature upshift. Feeding EGb 761 attenuated the intracellular levels of H₂O₂ in a dose-dependent manner (Fig. 6A). Feeding 100 μ g/ml EGb 761 exhibited most significant reduction (Ctrl, 100 \pm 23%; EGb, 42 \pm 7%; $n = 6$; $p = 0.04$). Interestingly, no significant reduction was observed in the control worms fed with EGb 761 (Fig. 6A), suggesting that the effect of EGb 761 on attenuating H₂O₂ may be specific to A β expression. In comparison, 50 μ g/ml L-ascorbic acid (VC) (Ctrl, 100 \pm 3%; VC, 31 \pm 3%; $n = 3$, $p = 0.04$), but not CR (139 μ g/ml) (Ctrl, 100 \pm 3%; CR, 101 \pm 18%; $n = 3$; $p = 0.11$; total of 300 worms in each group), also showed significant attenuation of the A β -induced elevation of H₂O₂. Among single components of EGb 761 tested (10 μ g/ml for each), only ginkgolide A significantly attenuated the levels of H₂O₂ ($p = 0.047$) (Fig. 6C).

Paralysis is associated with A β oligomers but not the level of H₂O₂


To determine whether there is a correlative association between the behavioral rescue and the decrease in A β oligomers by EGb 761 and its constituents, correlation analysis was performed. Figure 7A shows a clear correlation between the amount of A β oligomers (mean density) and paralysis (PT₅₀), with a value of the Pearson correlation coefficient r of 0.566 ($p = 0.044$). Among all of the compounds, GJ, GA, and EGb 761 appear to contribute the most to the correlation, in contrast with GC, GB, and BB. The latter three compounds decrease the A β oligomers density without improving the worms' paralysis. Figure 7B is a plot of paralysis (PT₅₀) versus levels of H₂O₂ measured by DCF fluorescence (see Materials and Methods). Paralysis is dissociated from the level of reactive oxygen species (ROS), because no correlation of the paralysis with the levels of H₂O₂ was found. The value of the Pearson correlation coefficient r was 0.27 ($p = 0.07$).

As summarized in Table 3, chemotaxis behavior does not correlate with the mean density of the A β oligomers. It is possible that the chemosensory neural circuits are less sensitive to the A β toxicity, and thus the chemotaxis behavior may not be the best assay for the latter. However, the serotonin hypersensitivity and paralysis correlate well with the A β oligomers density, supporting our hypothesis that EGb 761 and its constituents alleviate the behavioral abnormalities by decreasing the levels of the toxic A β oligomers.

Discussion

In the present experiments, we sought to associate A β species with A β -specific pathological behaviors using transgenic *C. elegans* as a model and EGb 761 as a pharmacological modulator. We found that EGb 761 alleviated A β -induced paralysis, chemotaxis dysfunction, and 5-HT hypersensitivity in the transgenic *C. elegans* expressing A β . EGb 761 also modulated A β oligomers and attenuated levels of H₂O₂ in the transgenic *C. elegans* (CL4176). Interestingly, suppression of paralysis is associated with inhibition of A β oligomerization but is disassociated with the antioxidative effect (Fig. 7A,B), suggesting that protective effects of EGb 761 against A β toxicity is mediated primarily by inhibition of A β oligomerization. If a similar mechanism is shared with other species, it may represent a rationale for the beneficial effects of EGb 761 in humans with AD-related dementia (Le Bars et al., 2000; Mazza et al., 2006) and for the enhanced

A Correlation analysis between paralysis and A β oligomers



B Correlation analysis between paralysis and levels of H₂O₂




Figure 7. Correlation analysis between paralysis (PT₅₀) and 20 kDa A β oligomers (A) or with levels of H₂O₂ (B). Correlation analysis was performed with the GraphPad Prism 4.0a, using a one-tailed Pearson test. The linear regression line along with the 95% confidence interval is shown only for illustrative purposes.

Table 3. Summary of correlation analysis

Behavior	Pearson <i>r</i> value	<i>p</i> value	Correlation with A β oligomers
Paralysis	−0.566	0.044	Yes
5-HT sensitivity	−0.644	0.043	Yes
Chemotaxis	−0.451	0.131	No

The data are from 10 different drugs tested in three behavior assays. Correlation of paralysis (PT₅₀), chemotaxis (CI), and 5-HT sensitivity with the 20 kDa A β oligomers mean density was tested by Pearson's test (see Materials and Methods).

cognitive function by EGb 761 in the transgenic mice model of AD (Tg2576) (Stackman et al., 2003).

EGb 761 has been known as an antioxidant that protects neuronal cells against amyloid toxicity in the cells (Bastianetto et al., 2000; Yao et al., 2001). We demonstrate here that EGb 761 and ginkgolide A suppress both A β oligomerization and A β -induced paralysis in a model organism, in a manner that does not parallel the effects of known antioxidants. We believe that the effect of

EGb 761 and ginkgolide A on A β toxicity is specific for the following reasons.

EGb 761 modulates A β specific pathological behaviors (Figs. 2, 3)

C. elegans is an ideal model organism for functional analysis of the age-associated neurodegeneration because of its available genetic information as well as the simple structure of its nervous system, which consist of only 302 neurons in an adult nematode. We took advantage of an established relationship between onset of A β expression and paralysis phenotype in a transgenic *C. elegans* model (Link, 1995). The absence of endogenous A β production in the worms offers an opportunity to find a direct role of the A β involvement in pathological behaviors (Wu and Luo, 2005). In addition, predominantly intracellular expression of A β provides another tool to address specific role of intracellular A β in relation to its toxicity (Gutierrez-Zepeda and Luo, 2004). Substantial evidence implicates intracellular A β oligomers in early events related to AD (Kienlen-Campard et al., 2002).

For the first time, two neuronal behavior phenotypes were characterized in a neuronal A β -expressing strain CL2355. Both chemotaxis and 5-HT signaling phenotypes are biologically relevant to A β -induced toxicity. *C. elegans* uses six primary sensory neurons to respond to >40 different attractants and repellents (Bargmann et al., 1993). The reinforcement is mediated by 5-HT signaling in the worms (Zhang et al., 2005). Several possible mechanisms could explain the 5-HT hypersensitive phenotype of the A β transgene, including the following: (1) response to 5-HT, which is modulated by calcium channel-dependent calcium influx (Schafer and Kenyon, 1995), was affected by the transgene A β expression (Ingram, 2005); (2) reduced acetylcholine (ACh) by A β accumulation in the worms; ACh is a negative regulator of 5-HT sensitivity in *C. elegans* (Schafer et al., 1996); and (3) A β may directly or indirectly block 5-HT reuptake in the worms. The pathological behavior assays used here are simple and relatively reproducible compared with other behavior assays in the worms. Whether this form of pathological behavior uses molecular mechanisms common to higher animals remains to be determined. A similar approach has recently revealed polyglutamine threshold toxicity in a transgenic *C. elegans* model of Huntington's disease (Brignull et al., 2006).

EGb 761 and ginkgolide A inhibit A β oligomerization (Fig. 4A–D)

Accumulation of A β oligomers seems to be one of the earliest events in the transgenic mice of AD (Oddo et al., 2006), which impairs long-term potentiation (LTP) and memory (Walsh et al., 2005; Lesne et al., 2006) and which correlates better with severity of dementia in AD patients than the density of amyloid plaques (Gong et al., 2003). Some oligomers (~20 kDa) inhibited by EGb 761 and ginkgolide A, as observed in this study (Fig. 4A–D), might be similar to, or identical with the neurotoxic ADDLs, or A β -derived diffusible ligands (Lambert et al., 1998). Previous studies showed that EGb 761 could inhibit formation of these species in solution (Luo et al., 2002; Chromy et al., 2003). The higher-order A β oligomers (~50 kDa) inhibited by EGb 761 in the worms (Fig. 4E,F) might be relevant to the species previously reported to be abundant in the AD brain (Gong et al., 2003). Our results suggest, but do not prove, that all of the oligomeric A β species inhibited by EGb 761 and ginkgolide A are toxic.

Based on *in vitro* studies, a linear pathway leading from A β monomers via paranuclei, oligomers, to protofibrils, and then

to fibrils was proposed (Bitan et al., 2003; Urbanc et al., 2004). This pathway may provide an explanation for our observation that Congo red, although it reduced A β deposits (Fig. 5B), did not significantly delay A β -induced paralysis (Fig. 2B). We speculate that EGb 761 and Congo red may bind to A β oligomers differently. For example, EGb 761 may have a higher affinity for certain oligomeric species, whereas Congo red favors the fibril form of A β . Thus, Congo red probably enters the linear process of fibrillogenesis at later stages than EGb 761, which would still lead to reduced overall A β oligomers (Fig. 4A) and decreased A β deposits (Fig. 5B), but not the appearance of A β monomers or significant suppression of paralysis (Fig. 2B) in the transgenic *C. elegans*. This theory agrees with the previous observation that paralysis occurs before detectable β -amyloid deposition in *C. elegans* (Drake et al., 2003). Therefore, the paralysis suppression by EGb 761 and ginkgolide A might be a consequence of the shift from the toxic A β oligomer to the nontoxic A β monomers (Fig. 4A). Given its “multipotent” nature, it is also possible that EGb 761 differentially modulates different processes of oligomerization.

The unique structure of ginkgolide A provides a rationale for its specific effect

It is believed that the unique biological properties of ginkgolides arise from their unique “cage skeleton” structure. This structure may share a common motif with Congo red and/or curcumin (Yang et al., 2005), which all display affinity for amyloidogenic conformations. It is not surprising that, among all ginkgolides, only GA exhibited a correlation between reducing paralysis (Fig. 2C) and inhibiting A β oligomers (Fig. 4C,D). Biological effects of ginkgolides can markedly vary with minor differences in their chemical structure. For example, the presence of one extra hydroxyl group at position 7 converts GB, a potent PAFR antagonist ($K_i = 0.88 \mu\text{M}$) into GC, a constituent with much lower activity ($K_i = 12.6 \mu\text{M}$) (Vogensen et al., 2003). Indeed, GA and GJ are the components of the extract that inhibit or even eliminate the deadly effects of A β on LTP (Nakanishi, 2005). GB affects hippocampal LTP by inhibiting the PAFR (Kondratskaya et al., 2004). Given that very little is known about which components of EGb 761 are the most efficacious, additional analyses of specific molecular structures of ginkgolides and of their interactions with A β may reveal novel compound(s) to protect against AD.

The fact that the antioxidant L-ascorbic acid lowered the levels of H₂O₂ (Fig. 6B), but did not inhibit A β oligomerization (Fig. 4A–C) nor suppressed paralysis (Table 2), suggests that a mechanism(s) other than the relief from antioxidative stress is operative in the ginkgolide-mediated protection against A β toxicity in *C. elegans*. This finding is consistent with the *in vitro* observation that no correlation exists between relative potencies of polyphenols in inhibiting A β fibril formation and in their oxidative features (Ono and Yamada, 2006). However, the dissociation of antioxidative effect from delaying paralysis does not rule out the contribution of oxidative damage in A β -dependent abnormal behaviors. Recent reports demonstrate that A β -induced, and a pro-aggregation-protein-induced paralysis in transgenic *C. elegans* were reduced by slowing aging (Cohen et al., 2006), or by coexpression of a stress response protein, the small heat shock protein 16 (sHSP-16) (Link et al., 2006) in the worms, respectively. EGb 761 attenuated the expression of sHSP-16 protein in the wild-type worms exposed to oxidative stress (Strayer et al.,

2003). A possible explanation is that EGb 761 might act in a manner similar to that of sHSP-16 in protecting against the deleterious process, which would lead to a decreased need for sHSP-16 expression. The antioxidative action of EGb 761 (Fig. 6B) may contribute to the beneficial effect in AD (Zandi et al., 2004), without being manifested in *C. elegans*.

Together, the neuroprotective effects of EGb 761 and ginkgolides on transgenic *C. elegans* behaviors provide strong evidence that formation of intracellular A β _{1–42} oligomers is directly associated with A β toxicity, supporting the view that the A β oligomers are the toxic species in AD (Kayed et al., 2003; Walsh and Selkoe, 2004). It is likely that the sequence of events manifested in the behavior of the transgenic worms, as well as the pharmacological efficacy, share the similar mechanisms with the cognitive impairment in mammals.

References

- Andrieu S, Gillette S, Amouyal K, Nourhashemi F, Reynish E, Ousset PJ, Albarede JL, Vellas B, Grandjean H (2003) Association of Alzheimer's disease onset with *Ginkgo biloba* and other symptomatic cognitive treatments in a population of women aged 75 years and older from the EPIDOS study. *J Gerontol A Biol Sci Med Sci* 58:372–377.
- Bargmann CI, Hartwig E, Horvitz HR (1993) Odorant-selective genes and neurons mediate olfaction in *C. elegans*. *Cell* 74:515–527.
- Bastianetto S, Ramassamy C, Dore S, Christen Y, Poirier J, Quirion R (2000) The *Ginkgo biloba* extract (EGb 761) protects hippocampal neurons against cell death induced by beta-amyloid. *Eur J Neurosci* 12:1882–1890.
- Birks J, Grimley EV, Van Dongen M (2002) *Ginkgo biloba* for cognitive impairment and dementia. *Cochrane Database Syst Rev* 4:CD003120.
- Bitan G, Kirkitadze MD, Lomakin A, Vollers SS, Benedek GB, Teplow DB (2003) Amyloid beta-protein (A β) assembly: A β 40 and A β 42 oligomerize through distinct pathways. *Proc Natl Acad Sci USA* 100:330–335.
- Brignull HR, Moore FE, Tang SJ, Morimoto RI (2006) Polyglutamine proteins at the pathogenic threshold display neuron-specific aggregation in a pan-neuronal *Caenorhabditis elegans* model. *J Neurosci* 26:7597–7606.
- Christen Y, Maixent JM (2002) What is *Ginkgo biloba* extract EGb 761? An overview—from molecular biology to clinical medicine. *Cell and Mol Biol* 48:601–611.
- Chromy BA, Nowak RJ, Lambert MP, Viola KL, Chang L, Velasco PT, Jones BW, Fernandez SJ, Lacor PN, Horowitz P, Finch CE, Krafft GA, Klein WL (2003) Self-assembly of A β (1–42) into globular neurotoxins. *Biochemistry* 42:12749–12760.
- Cohen E, Bieschke J, Perciavalle RM, Kelly JW, Dillin A (2006) Opposing activities protect against age-onset proteotoxicity. *Science* 313:1604–1610.
- DeFeudis FV (1998) *Ginkgo biloba* extract (EGb 761): from chemistry to clinic. Weisbaden, Germany: Ullstein Medical.
- DeFeudis FV (2002) Effects of *Ginkgo biloba* extract (EGb761) on gene expression: possible relevance to neurological disorders and age-associated cognitive impairment. *Drug Dev Res* 57:214–235.
- DeKosky ST, Fitzpatrick A, Ives DG, Saxton J, Williamson J, Lopez OL, Burke G, Fried L, Kuller LH, Robbins J, Tracy R, Woolard N, Dunn L, Kronmal R, Nahin R, Furberg C (2006) The Ginkgo Evaluation of Memory (GEM) study: design and baseline data of a randomized trial of *Ginkgo biloba* extract in prevention of dementia. *Contemp Clin Trials* 27:238–253.
- Drake J, Link CD, Butterfield DA (2003) Oxidative stress precedes fibrillar deposition of Alzheimer's disease amyloid beta-peptide (1–42) in a transgenic *Caenorhabditis elegans* model. *Neurobiol Aging* 24:415–420.
- Gong Y, Chang L, Viola KL, Lacor PN, Lambert MP, Finch CE, Krafft GA, Klein WL (2003) Alzheimer's disease-affected brain: presence of oligomeric A β ligands (ADDLs) suggests a molecular basis for reversible memory loss. *Proc Natl Acad Sci USA* 100:10417–10422.
- Gutierrez-Zepeda A, Luo Y (2004) Testing the amyloid toxicity hypothesis of Alzheimer's disease in transgenic *Caenorhabditis elegans* model. *Front Biosci* 9:3333–3338.

- Hardy J, Selkoe DJ (2002) The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science* 297:353–356.
- Hoertel O (2003) Behavioral plasticity in *C. elegans*: paradigms, circuits, genes. *J Neurobiol* 54:203–223.
- Horvitz HR, Chalfie M, Trent C, Sulston JE, Evans PD (1982) Serotonin and octopamine in the nematode *Caenorhabditis elegans*. *Science* 216:1012–1014.
- Ingram VM (2005) The role of Alzheimer Abeta peptides in ion transport across cell membranes. *Subcell Biochem* 38:339–349.
- Ivic L, Sands TT, Fishkin N, Nakanishi K, Kriegstein AR, Stromgaard K (2003) Terpene trilactones from *Ginkgo biloba* are antagonists of cortical glycine and GABA(A) receptors. *J Biol Chem* 278:49279–49285.
- Jaracz S, Malik S, Nakanishi K (2004) Isolation of ginkgolides A, B, C, J and bilobalide from *G. biloba* extracts. *Phytochemistry* 65:2897–2902.
- Kayed R, Head E, Thompson JL, McIntire TM, Milton SC, Cotman CW, Glabe CG (2003) Common structure of soluble amyloid oligomers implies common mechanism of pathogenesis. *Science* 300:486–489.
- Kienlen-Campard P, Miolet S, Tasiaux B, Octave JN (2002) Intracellular amyloid-beta 1–42, but not extracellular soluble amyloid-beta peptides, induces neuronal apoptosis. *J Biol Chem* 277:15666–15670.
- Kondratskaya EL, Pankratov YV, Lalo UV, Chatterjee SS, Krishtal OA (2004) Inhibition of hippocampal LTP by ginkgolide B is mediated by its blocking action on PAF rather than glycine receptors. *Neurochem Int* 44:171–177.
- Lambert MP (2006) Monoclonal antibodies that target pathological assemblies of A β . *J Neurochem*, in press.
- Lambert MP, Barlow AK, Chromy BA, Edwards C, Freed R, Liosatos M, Morgan TE, Rozovsky I, Trommer B, Viola KL, Wals P, Zhang C, Finch CE, Krafft GA, Klein WL (1998) Diffusible, nonfibrillar ligands derived from Abeta-42 are potent central nervous system neurotoxins. *Proc Natl Acad Sci USA* 95:6448–6453.
- Lambert MP, Viola KL, Chromy BA, Chang L, Morgan TE, Yu J, Venton DL, Krafft GA, Finch CE, Klein WL (2001) Vaccination with soluble Abeta oligomers generates toxicity-neutralizing antibodies. *J Neurochem* 79:595–605.
- Le Bars PL (2003) Magnitude of effect and special approach to *Ginkgo biloba* extract EGb 761 in cognitive disorders. *Pharmacopsychiatry* 36 [Suppl 1]:S44–S49.
- Le Bars PL, Katz MM, Berman N, Itil TM, Freedman AM, Schatzberg AF (1997) A placebo-controlled, double-blind, randomized trial of an extract of *Ginkgo biloba* for dementia. North Am EGb Study Group. *JAMA* 278:1327–1332.
- Le Bars PL, Kieser M, Itil KZ (2000) A 26-week analysis of a double-blind, placebo-controlled trial of the *Ginkgo biloba* extract EGb 761 in dementia. *Dement Geriatr Cogn Disord* 11:230–237.
- Lesne S, Koh MT, Kotilinek L, Kaye R, Glabe CG, Yang A, Gallagher M, Ashe KH (2006) A specific amyloid-beta protein assembly in the brain impairs memory. *Nature* 440:352–357.
- Link CD (1995) Expression of human beta-amyloid peptide in transgenic *Caenorhabditis elegans*. *Proc Natl Acad Sci USA* 92:9368–9372.
- Link CD (2005) Invertebrate models of Alzheimer's disease. *Genes Brain Behav* 4:147–156.
- Link CD, Johnson CJ, Fonte V, Paupard M, Hall DH, Styren S, Mathis CA, Klunk WE (2001) Visualization of fibrillar amyloid deposits in living, transgenic *Caenorhabditis elegans* animals using the sensitive amyloid dye, X-34. *Neurobiol Aging* 22:217–226.
- Link CD, Taft A, Kapulkin V, Duke K, Kim S, Fei Q, Wood DE, Sahagan BG (2003) Gene expression analysis in a transgenic *Caenorhabditis elegans* Alzheimer's disease model. *Neurobiol Aging* 24:397–413.
- Link CD, Fonte V, Hiester B, Yerg J, Ferguson J, Csontos S, Silverman MA, Stein GH (2006) Conversion of green fluorescent protein into a toxic, aggregation-prone protein by C-terminal addition of a short peptide. *J Biol Chem* 281:1808–1816.
- Lorenzo A, Yankner BA (1994) Beta-amyloid neurotoxicity requires fibril formation and is inhibited by Congo red. *Proc Natl Acad Sci USA* 91:12243–12247.
- Luo Y (2001) *Ginkgo biloba* neuroprotection: therapeutic implications in Alzheimer's disease. *J Alzheimers Dis* 3:401–407.
- Luo Y, Smith JV, Paramasivam V, Burdick A, Curry KJ, Buford JP, Khan I, Netzer WJ, Xu H, Butko P (2002) Inhibition of amyloid-beta aggregation and caspase-3 activation by the *Ginkgo biloba* extract EGb761. *Proc Natl Acad Sci USA* 99:12197–12202.
- Maezawa I, Hong HS, Wu HC, Battina SK, Rana S, Iwamoto T, Radke GA, Pettersson E, Martin GM, Hua DH, Jin LW (2006) A novel tricyclic pyrone compound ameliorates cell death associated with intracellular amyloid-beta oligomeric complexes. *J Neurochem* 98:57–67.
- Matsuura T, Sato T, Shingai R (2005) Interactions between *Caenorhabditis elegans* individuals during chemotactic response. *Zool Sci* 22:1095–1103.
- Mazza M, Capuano A, Bria P, Mazza S (2006) *Ginkgo biloba* and donepezil: a comparison in the treatment of Alzheimer's dementia in a randomized placebo-controlled double-blind study. *Eur J Neurol* 13:981–985.
- Mix JA, Crews Jr WD (2002) A double-blind, placebo-controlled, randomized trial of *Ginkgo biloba* extract EGb 761(R) in a sample of cognitively intact older adults: neuropsychological findings. *Hum Psychopharmacol* 17:267–277.
- Nakanishi K (1967) The ginkgolides. *Pure Appl Chem* 14:89–113.
- Nakanishi K (2005) Terpene trilactones from *Ginkgo biloba*: from ancient times to the 21st century. *Bioorg Med Chem* 13:4987–5000.
- Nuttley WM, Atkinson-Leadbetter KP, Van Der Kooy D (2002) Serotonin mediates food-odor associative learning in the nematode *Caenorhabditis elegans*. *Proc Natl Acad Sci USA* 99:12449–12454.
- Oddo S, Caccamo A, Tran L, Lambert MP, Glabe CG, Klein WL, LaFerla FM (2006) Temporal profile of amyloid-beta (Abeta) oligomerization in an in vivo model of Alzheimer disease. A link between Abeta and tau pathology. *J Biol Chem* 281:1599–1604.
- Ono K, Yamada M (2006) Antioxidant compounds have potent anti-fibrillogenic and fibril-destabilizing effects for alpha-synuclein fibrils in vitro. *J Neurochem* 97:105–115.
- Pike CJ, Walencewicz-Wasserman AJ, Kosmoski J, Cribbs DH, Glabe CG, Cotman CW (1995) Structure-activity analyses of beta-amyloid peptides: contributions of the beta 25–35 region to aggregation and neurotoxicity. *J Neurochem* 64:253–265.
- Ranganathan R, Cannon SC, Horvitz HR (2000) MOD-1 is a serotonin-gated chloride channel that modulates locomotory behaviour in *C. elegans*. *Nature* 408:470–475.
- Ranganathan R, Sawin ER, Trent C, Horvitz HR (2001) Mutations in the *Caenorhabditis elegans* serotonin reuptake transporter MOD-5 reveal serotonin-dependent and -independent activities of fluoxetine. *J Neurosci* 21:5871–5884.
- Roselli F, Tirard M, Lu J, Hutzler P, Lamberti P, Livrea P, Morabito M, Almeida OF (2005) Soluble beta-amyloid1–40 induces NMDA-dependent degradation of postsynaptic density-95 at glutamatergic synapses. *J Neurosci* 25:11061–11070.
- Sawin ER, Ranganathan R, Horvitz HR (2000) *C. elegans* locomotory rate is modulated by the environment through a dopaminergic pathway and by experience through a serotonergic pathway. *Neuron* 26:619–631.
- Schafer WR, Kenyon CJ (1995) A calcium-channel homologue required for adaptation to dopamine and serotonin in *Caenorhabditis elegans*. *Nature* 375:73–78.
- Schafer WR, Sanchez BM, Kenyon CJ (1996) Genes affecting sensitivity to serotonin in *Caenorhabditis elegans*. *Genetics* 143:1219–1230.
- Smith JV, Luo Y (2003) Elevation of oxidative free radicals in Alzheimer's disease models can be attenuated by *Ginkgo biloba* extract EGb 761. *J Alzheimers Dis* 5:287–300.
- Smith PF, MacLennan K, Darlington CL (1996) The neuroprotective properties of the *Ginkgo biloba* leaf: a review of the possible relationship to platelet-activating factor (PAF). *J Ethnopharmacol* 50:131–139.
- Stackman RW, Eckenstein F, Frei B, Kulhanek D, Nowlin J, Quinn JF (2003) Prevention of age-related spatial memory deficits in a transgenic mouse model of Alzheimer's disease by chronic *Ginkgo biloba* treatment. *Exp Neurol* 184:510–520.
- Strayer A, Wu Z, Christen Y, Link CD, Luo Y (2003) Expression of the small heat-shock protein Hsp16-2 in *Caenorhabditis elegans* is suppressed by *Ginkgo biloba* extract EGb 761. *FASEB J* 17:2305–2307.
- Urbanc B, Cruz L, Yun S, Buldyrev SV, Bitan G, Teplow DB, Stanley HE (2004) In silico study of amyloid {beta}-protein folding and oligomerization. *Proc Natl Acad Sci USA* 101:17345–17350.
- Vogensen SB, Stromgaard K, Shindou H, Jaracz S, Suehiro M, Ishii S, Shimizu T, Nakanishi K (2003) Preparation of 7-substituted ginkgolide derivatives: potent platelet activating factor (PAF) receptor antagonists. *J Med Chem* 46:601–608.

- [Walsh DM, Selkoe DJ \(2004\) Oligomers on the brain: the emerging role of soluble protein aggregates in neurodegeneration. *Protein Pept Lett* 11:213–228.](#)
- [Walsh DM, Townsend M, Podlisny MB, Shankar GM, Fadeeva JV, Agnaf OE, Hartley DM, Selkoe DJ \(2005\) Certain inhibitors of synthetic amyloid \$\beta\$ -peptide \(A \$\beta\$ \) fibrillogenesis block oligomerization of natural A \$\beta\$ and thereby rescue long-term potentiation. *J Neurosci* 25:2455–2462.](#)
- [Watanabe CM, Wolfram S, Ader P, Rimbach G, Packer L, Maguire JJ, Schultz PG, Gohil K \(2001\) The in vivo neuromodulatory effects of the herbal medicine *Ginkgo biloba*. *Proc Natl Acad Sci USA* 98:6577–6580.](#)
- [Wood W \(1988\) The nematode *Caenorhabditis elegans*. Plainview, NY: Cold Spring Harbor Laboratory.](#)
- [Wu Y, Luo Y \(2005\) Transgenic *C. elegans* as a model in Alzheimer's research. *Curr Alzheimer Res* 2:37–45.](#)
- [Yang F, Lim GP, Begum AN, Ubeda OJ, Simmons MR, Ambegaokar SS, Chen PP, Kaye R, Glabe CG, Frautschy SA, Cole GM \(2005\) Curcumin inhibits formation of amyloid beta oligomers and fibrils, binds plaques, and reduces amyloid in vivo. *J Biol Chem* 280:5892–5901.](#)
- [Yao Z, Drieu K, Papadopoulos V \(2001\) The *Ginkgo biloba* extract EGb 761 rescues the PC12 neuronal cells from beta-amyloid-induced cell death by inhibiting the formation of beta-amyloid-derived diffusible neurotoxic ligands. *Brain Res* 889:181–190.](#)
- [Zandi PP, Anthony JC, Khachaturian AS, Stone SV, Gustafson D, Tschanz JT, Norton MC, Welsh-Bohmer KA, Breitner JC \(2004\) Reduced risk of Alzheimer disease in users of antioxidant vitamin supplements: the Cache County Study. *Arch Neurol* 61:82–88.](#)
- [Zhang Y, Lu H, Bargmann CI \(2005\) Pathogenic bacteria induce aversive olfactory learning in *Caenorhabditis elegans*. *Nature* 438:179–184.](#)